

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07D 337/08, 409/10, C08G 65 /359, A61K 31 /28		(11) International Publication Number: WO 97/33882	
(21) International Application Number: PCT/US97/04076		(43) International Publication Date: 18 September 1997 (18.09.97)	
(22) International Filing Date: 11 March 1997 (11.03.97)		(74) Agents: ROEDEL, John, K., Jr., et al.; Sonniger, Powers, Leavitt & Roedel, 16th floor, One Metropolitan Square, St. Louis, MO 63102 (US).	
(30) Priority Data: 08/013,119 11 March 1996 (11.03.96) US 08/616,065 11 March 1997 (11.03.97) US		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UO), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, SI, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TO).	
(71) Applicant (for all designated States except US): G.D. SEARLE AND CO. (USUS); P.O. Box 5110, Chicago, IL 60680-9899 (US).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(75) Inventors/Applicants (for US only): REITZ, David, B. (USUS); P.O. Box 5110, Chicago, IL 60680-9899 (US); LEE, Len, F. (USUS); P.O. Box 5110, Chicago, IL 60680-9899 (US); LI, Jinglin, J. (USUS); P.O. Box 5110, Chicago, IL 60680-9899 (US); HUANG, Hame-Chih (USUS); P.O. Box 5110, Chicago, IL 60680-9899 (US); TREMONT, Samuel, J. (USUS); P.O. Box 5110, Chicago, IL 60680-9899 (US); MILLER, Raymond, E. (USUS); P.O. Box 5110, Chicago, IL 60680-9899 (US); BANERJEE, Shyamal, C. (USUS); P.O. Box 5110, Chicago, IL 60680-9899 (US).			
(54) Title: NOVEL BENZOTHIOPHINES HAVING ACTIVITY AS INHIBITORS OF ILEAL BILE ACID TRANSPORT AND TAUROCHOLATE UPTAKE			
(57) Abstract Provided are novel benzothiephines, derivatives, and analogs thereof; pharmaceutical compositions containing them; and methods of using these compounds and compositions in the prophylaxis and treatment of hyperlipidemic conditions such as those associated with atherosclerosis or hypercholesterolemia, in mammals.			

FOR THE PURPOSES OF INFORMATION ONLY
Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MF	Malawi
AN	Angola	GE	Georgia	MG	Mexico
AT	Austria	GN	Ghana	ML	Mali
AU	Australia	GR	Greece	NL	Netherlands
BA	Bosnia and Herzegovina	HR	Croatia	NO	Norway
BB	Barbados	HU	Hungary	NZ	New Zealand
BE	Belgium	IE	Ireland	PE	Peru
BF	Burkina Faso	IT	Italy	PG	Papua New Guinea
BG	Bulgaria	JP	Japan	PH	Philippines
BJ	Benin	KE	Kenya	PL	Poland
BR	Brazil	KG	Kyrgyzstan	PT	Portugal
BS	Bahamas	KH	Kampuchea	RO	Romania
BT	Bhutan	KM	Comoros	RU	Russian Federation
CA	Canada	KR	Republic of Korea	SD	Sudan
CF	Central African Republic	KZ	Kazakhstan	SE	Sweden
CG	Congo	LA	Laos	SG	Singapore
CH	Switzerland	LB	Lebanon	SI	Slovenia
CN	China	LC	Liechtenstein	SK	Slovakia
CO	Colombia	LI	Lithuania	SN	Senegal
CZ	Czech Republic	LU	Luxembourg	SV	Sweden
DE	Germany	LV	Latvia	TD	Chad
DK	Denmark	MC	Monaco	TG	Togo
EE	Estonia	MD	Republic of Moldova	TH	Thailand
ES	Spain	ME	Montenegro	TL	Turkey
FI	Finland	ML	Mali	TM	Timor-Leste
FR	France	MR	Mauritania	US	United States of America
GA	Gabon	MT	Malta	UZ	Uzbekistan
		MA	Morocco	VN	Viet Nam

NOVEL BENZOTRIEPINES HAVING ACTIVITY AS INHIBITORS
OF ILEAL BILE ACID TRANSPORT AND TAUROCHOLATE UPTAKE

This application claims the benefit of priority of
U.S. Provisional Application No. 60/013,119, filed
March 11, 1996, which is a continuation in part of U.S.
Serial No. 08/_____, filed August 21, 1995, which is
a continuation-in-part of U.S. Serial No. 08/305,526
filed September 12, 1994, both now pending.

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention relates to novel
benzothiepines, derivatives and analogs thereof,
pharmaceutical compositions containing them, and their
use in medicine, particularly in the prophylaxis and
treatment of hyperlipidemic conditions such as is
associated with atherosclerosis or
hypercholesterolemia, in mammals.

Description of Related Art

It is well-settled that hyperlipidemic conditions
associated with elevated concentrations of total
cholesterol and low-density lipoprotein cholesterol are
major risk factors for coronary heart disease and
particularly atherosclerosis. Interfering with the
circulation of bile acids within the lumen of the
intestinal tract is found to reduce the levels of serum
cholesterol in a causal relationship. Epidemiological
data has accumulated which indicates such reduction
leads to an improvement in the disease state of

atherosclerosis. Steedronsky, in "Interaction of bile
acids and cholesterol with nonsystemic agents having
hypocholesterolemic properties," *Biochimica et
Biophysica Acta*, 1210 (1994) 255-267 discusses the
biochemistry, physiology and known active agents
surrounding bile acids and cholesterol.

Pathophysiologic alterations are shown to be
consistent with interruption of the enterohepatic
circulation of bile acids in humans by Heubi, J.E., et
al. See "Primary Bile Acid Malabsorption: Defective
in Vitro Ileal Active Bile Acid Transport",
Gastroenterology, 1982:83:804-11.

In fact, cholestyramine binds the bile acids in
the intestinal tract, thereby interfering with their
normal enterohepatic circulation (Reihner, E. et al, in
"Regulation of hepatic cholesterol metabolism in
humans: stimulatory effects of cholestyramine on FMG-
CoA reductase activity and low density lipoprotein
receptor expression in gallstone patients", *Journal of
Lipid Research*, Volume 31, 1990, 2219-2226 and Suckling
et al, "Cholesterol lowering and bile acid excretion in
the hamster with cholestyramine treatment",
Atherosclerosis, 89(1991) 183-190). This results in an

increase in liver bile acid synthesis by the liver
using cholesterol as well as an upregulation of the
liver LDL receptors which enhances clearance of
cholesterol and decreases serum LDL cholesterol levels.

In another approach to the reduction of
recirculation of bile acids, the ileal bile acid
transport system is a putative pharmaceutical target
for the treatment of hypercholesterolemia based on an
interruption of the enterohepatic circulation with

specific transport inhibitors (Kramer, et al.

"Intestinal Bile Acid Absorption" The Journal of Biological Chemistry, Vol. 268, No. 24, Issue of August 25, pp. 18035-18046, 1993).

In a series of patent applications, eg Canadian Patent Application Nos. 2,025,294; 2,078,588;

2,085,782; and 2,085,830; and EP Application Nos. 0 379 161; 0 549 967; 0 559 064; and 0 563 731, Hoechst

Aktiengesellschaft discloses polymers of various

naturally occurring constituents of the enterohepatic circulation system and their derivatives, including bile acid, which inhibit the physiological bile acid transport with the goal of reducing the LDL cholesterol level sufficiently to be effective as pharmaceuticals and, in particular for use as hypocholesterolemic agents.

In vitro bile acid transportinhibition is

disclosed to show hypolipidemic activity in The

Wellcome Foundation Limited disclosure of the world

patent application number WO 93/16055 for

"Hypolipidemic Benzothiazepine Compounds"

Selected benzothiepienes are disclosed in world

patent application number WO93/321146 for numerous uses including fatty acid metabolism and coronary vascular diseases.

Other selected benzothiepienes are known for use as hypolipaeamic and hypocholesterolaemic agents,

especially for the treatment or prevention of

atherosclerosis as disclosed by application Nos. EP

508425, FR 2661676, and WO 92/18462, each of which is limited by an amide bonded to the carbon adjacent the phenyl ring of the fused bicyclo benzothiepine ring.


The above references show continuing efforts to find safe, effective agents for the prophylaxis and treatment of hyperlipidemic diseases and their usefulness as hypocholesterolemic agents.

Additionally selected benzothiepienes are disclosed for use in various disease states not within the present invention utility. These are EP 568 898A as abstracted by Derwent Abstract No. 93-351589; WO 89/1477/A as abstracted in Derwent Abstract No. 89-370688; U.S. 3,520,891 abstracted in Derwent 50701R-B; US 3,287,370, US 3,389,144; US 3,694,446 abstracted in Derwent Abstr. No. 65860T-B and WO 92/18462.

The present invention furthers such efforts by providing novel benzothiepienes, pharmaceutical compositions, and methods of use therefor.

SUMMARY OF THE INVENTION

Accordingly, among its various aspects, the present invention provides compounds of formula (I):



wherein:

q is an integer from 1 to 4;

n is an integer from 0 to 2;

R¹ and R² are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, arylalkyl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl,

wherein alkyl, alkenyl, alkynyl, haloalkyl, arylalkyl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more substituent selected from the group consisting of OR⁹, NR⁹R¹⁰, NR⁹R^{9A}, SR⁹, SR^{9A}, PR⁹R^{9A}, S(O)R⁹, SO₂R⁹, SO₃R⁹, CO₂R⁹, CN, halogen, oxo, and CONR⁹R¹⁰, wherein alkyl, alkenyl, alkynyl, alkylaryl, alkoxy, alkoxyalkyl, polyalkyl, aryl, and cycloalkyl

5

optionally have one or more carbons replaced by O, NR⁹, NR⁹R¹⁰, S, SO, SO₂, S⁺R^{9A}, P⁺R⁹R¹⁰, or phenylene, wherein R⁹, R¹⁰, and R^{9A} are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, alkylammoniumalkyl, and arylalkyl; or

R¹ and R² taken together with the carbon to which they are attached form C₁-C₆ cycloalkylidene;

R³ and R⁴ are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, acyloxy, aryl, heterocycle, OR⁹, NR⁹R¹⁰, SR⁹, S(O)R⁹, SO₂R⁹, and SO₃R⁹, wherein R⁹ and R¹⁰ are as defined above; or

R³ and R⁴ together form =O, =NOR¹¹, =S, =NNR¹¹R¹², =NR⁹, or =CR¹¹R¹²,

wherein R¹¹ and R¹² are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl, heterocycle, carboxyalkyl, carboalkoxyalkyl, cycloalkyl, cyanoalkyl, OR⁹, NR⁹R¹⁰, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, CO₂R⁹, CN, halogen, oxo, and CONR⁹R¹⁰, wherein R⁹ and R¹⁰ are as defined above, provided that both R³ and R⁴ cannot be OH, NH₂, and SH, or R¹¹ and R¹² together with the nitrogen or carbon atom to which they are attached form a cyclic ring; R⁵ and R⁶ are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl,

6

cycloalkyl, heterocycle, quaternary heterocycle, OR^9 , SR^9 , $S(O)R^9$, SO_2R^9 , and SO_3R^9 ,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, and quaternary heteroaryl can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary

heteroaryl, halogen, oxo, OR^{13} , $NR^{13}R^{14}$, SR^{13} , $S(O)R^{13}$, SO_2R^{13} , SO_3R^{13} , $NR^{13}OR^{14}$, $NR^{13}NR^{14}R^{15}$, NO_2 , CO_2R^{13} , CN , OM , SO_2OM , $SO_2NR^{13}R^{14}$, $C(O)NR^{13}R^{14}$, $C(O)OM$, COR^{13} , $P(O)R^{13}R^{14}$, $P^+R^{13}R^{14}R^{15}A^-$, $P(OR^u)OR^u$, $SR^uR^uA^-$, and $N^+R^{13}R^{14}R^{15}A^-$,

wherein:

A^- is a pharmaceutically acceptable anion and M is a pharmaceutically acceptable cation,

said alkyl, alkenyl, alkynyl, polyalkyl,

polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of OR^7 ,

NR^7R^8 , SR^7 , $S(O)R^7$, SO_2R^7 , SO_3R^7 , CO_2R^7 , CN , oxo,

$CONR^7R^8$, $N^+R^7R^8R^9A^-$, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary

heterocycle, quaternary heteroaryl, $P(O)R^7R^8$,

$P^+R^7R^8R^9A^-$, and $P(O)(OR^7)OR^8$, and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle

can optionally have one or more carbons replaced by O , NR^7 , $N^+R^7R^8A^-$, S , SO , SO_2 , $S^+R^7A^-$, PR^7 , $P(O)R^7$, $P^+R^7R^8A^-$, or phenylene, and R^{13} , R^{14} , and R^{15} are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, polyalkyl, aryl, arylalkyl, cycloalkyl, heterocycle, heterocycle, quaternary heterocycle, quaternary heteroaryl, and quaternary heteroarylalkyl.

wherein alkyl, alkenyl, alkynyl, arylalkyl, heterocycle, and polyalkyl optionally have one or more carbons replaced by O , NR^9 , $N^+R^9R^{10}A^-$, S , SO , SO_2 ,

$S^+R^9A^-$, PR^9 , $P^+R^9R^{10}A^-$, $P(O)R^9$, phenylene, carbohydrate, amino acid, peptide, or polypeptide, and

R^{13} , R^{14} , and R^{15} are optionally substituted with one or more groups selected from the group consisting of sulfoalkyl, quaternary heterocycle, quaternary

heteroaryl, OR^9 , NR^9R^{10} , $N^+R^9R^{11}R^{12}A^-$, SR^9 , $S(O)R^9$, SO_2R^9 , SO_3R^9 , oxo, CO_2R^9 , CN , halogen, $CONR^9R^{10}$, SO_2OM , $SO_2NR^9R^{10}$, $PO(OR^{16})OR^{17}$, $P^+R^9R^{10}A^-$, $S^+R^9A^-$, and $C(O)OM$,

wherein R^{16} and R^{17} are independently selected from the substituents constituting R^9 and M , and p is 0 or 1; or

R^{14} and R^{15} , together with the nitrogen atom to which they are attached, form a cyclic ring;

R^7 and R^8 are independently selected from the group consisting of hydrogen and alkyl; and

one or more R^X are independently selected from the group consisting of H , alkyl, alkenyl, alkynyl,

polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heterocycle, polyether, quaternary heterocycle, quaternary heteroaryl, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, S(O)₂R¹³, SO₃R¹³, S⁺R¹³R¹⁴-, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, NR¹³C(O)R¹⁴, C(O)NR¹³R¹⁴, NR¹⁴C(O)R¹³, C(O)OM, COR¹³, OR¹⁸, S(O)NR¹⁸, NR¹³R¹⁸, NR¹⁸OR¹⁴, N⁺R⁹R¹¹R¹²-, P⁺R⁹R¹¹R¹²-, amino acid, peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary heteroaryl can be further substituted with OR⁹, NR⁹R¹⁰, N⁺R⁹R¹¹R¹²-, SR⁹, S(O)R⁹, SO₂R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰, SO₂OM, SO₂NR⁹R¹⁰, PO(OR¹⁰)OR¹¹, P⁺R⁹R¹¹R¹²-, S⁺R⁹R¹¹-, or C(O)OM, and

wherein R¹⁸ is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heterocycle, and alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heterocycle, alkyl quaternary heterocycle, and quaternary heteroaryl optionally are substituted with one or more substituent selected from the group consisting of OR⁹, NR⁹R¹⁰, N⁺R⁹R¹¹R¹²-, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰, SO₂R⁹, SO₂NR⁹R¹⁰, PO(OR¹⁶)OR¹⁷, and C(O)OM,

wherein in R^x, one or more carbons are optionally replaced by O, NR¹³, N⁺R¹³R¹⁴-, S, SO, SO₂, S⁺R¹³R¹⁴-, PR¹³, P(O)R¹³, P⁺R¹³R¹⁴-, phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl,

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR⁹, N⁺R⁹R¹⁰-, S, SO, SO₂, S⁺R⁹R¹⁰-, PR⁹, P⁺R⁹R¹⁰-, or P(O)R⁹;

wherein quaternary heterocycle and quaternary heteroaryl are optionally substituted with one or more groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, SO₃R¹³, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴, P⁺R¹³R¹⁴R¹⁵-, P(OR¹⁰)OR¹¹, S⁺R⁹R¹¹R¹²-, and N⁺R⁹R¹¹R¹²-, provided that both R⁵ and R⁶ cannot be hydrogen,

provided that when R⁵ is OH, R¹, R², R³, R⁴, R⁷ and R⁸ cannot be all hydrogen;

provided that when R¹ or R¹ is phenyl, only one of R¹ or R¹ is H;

provided that when q = 1 and R¹ is styryl,

anilido, or anilino, only one of R¹ or R¹ is alkyl; or

a pharmaceutically acceptable salt, solvate, or produg thereof.

Preferably, R⁵ and R⁶ can independently be

selected from the group consisting of H, aryl,

heterocycle, quaternary heterocycle, and quaternary heteroaryl.

5

wherein said aryl, heteroaryl, quaternary

heterocycle, and quaternary heteroaryl can be

substituted with one or more substituent groups

independently selected from the group consisting of

alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl,

haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen,

oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, SO₃R¹³,

NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM,

SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴,

P⁺R¹³R¹⁴R¹⁵A⁻, P(OR¹³)OR¹⁴, S⁺R¹³R¹⁴A⁻, and N⁺R¹³R¹⁴R¹⁵A⁻,

polyether, aryl, haloalkyl, cycloalkyl, alkynyl, polyalkyl,

can optionally have one or more carbons replaced by O,

NR⁷, N⁺R⁷R⁸A⁻, S, SO, SO₂, S⁺R⁷A⁻, PR⁷, P(O)R⁷,

P⁺R⁷R⁸A⁻, or phenylene,

20

wherein said alkyl, alkenyl, alkynyl, polyalkyl,

polyether, aryl, haloalkyl, cycloalkyl, and heterocycle

can be further substituted with one or more substituent

groups selected from the group consisting of OR⁷,

NR⁷R⁸, SR⁷, S(O)R⁷, SO₂R⁷, SO₃R⁷, CO₂R⁷, CN, oxo,

CONR⁷R⁸, N⁺R⁷R⁸A⁻, alkyl, alkenyl, alkynyl, aryl,

cycloalkyl, heterocycle, arylalkyl, quaternary

heterocycle, quaternary heteroaryl, P(O)R⁷R⁸, P⁺R⁷R⁸A⁻,

and P(O)(OR⁷)OR⁸.

25

//

More preferably, R' or R' has the formula:



5

wherein:

t is an integer from 0 to 5;

Ar is selected from the group consisting of

phenyl, thiophenyl, pyridyl, piperazinyl, piperonyl,

pyrrolyl, naphthyl, furanyl, anthracenyl, quinolinyl,

isoquinolinyl, quinoxalinyl, imidazolyl, pyrazolyl,

oxazolyl, isoxazolyl, pyrimidinyl, thiazolyl,

triazolyl, isothiazolyl, indolyl, benzimidazolyl,

benzoxazolyl, benzothiazolyl, and benzoisothiazolyl;

and

15

one or more R^y are independently selected from the

group consisting of H, alkyl, alkenyl, alkynyl, aryl,

cycloalkyl, heterocycle, quaternary heterocycle, OR⁹,

SR⁹, S(O)R⁹, SO₂R⁹, and SO₃R⁹,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl,

and heterocycle can be substituted with one or more

substituent groups independently selected from the

group consisting of alkyl, alkenyl, alkynyl, polyalkyl,

polyether, aryl, haloalkyl, cycloalkyl, heterocycle,

arylalkyl, halogen, oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³,

SO₂R¹³, SO₃R¹³, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN,

OM, SO₂OM, SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³,

P(O)R¹³R¹⁴, P⁺R¹³R¹⁴R¹⁵A⁻, P(OR¹³)OR¹⁴, S⁺R¹³R¹⁴A⁻, and


N⁺R¹³R¹⁴R¹⁵A⁻,

22

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of OR^7 , NR^7R^8 , SR^7 , $S(O)R^7$, SO_2R^7 , SO_3R^7 , CO_2R^7 , CN , oxo , $CONR^7R^8$, $N^+R^7R^8R^9A^-$, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, $P(O)R^7R^8$, $P^+R^7R^8A^-$, and $P(O)(OR^7)OR^8$, and


wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O , NR^7 , $N^+R^7R^8A^-$, S , SO , SO_2 , $S^+R^7A^-$, PR^7 , $P(O)R^7$, $P^+R^7R^8A^-$, or phenylene.

Most preferably, R^5 or R^6 has the formula (II):




(II)

The invention is further directed to a compound selected from among:




and



wherein R^{10} is selected from the group

consisting of alkane diyl, alkene diyl, alkynes diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, peptide, and polypeptide, wherein alkane diyl, alkene diyl, alkynes diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, peptide, and polypeptide can optionally have one or more carbon atoms replaced by O , NR^7 , $N^+R^7R^8$, S , SO , SO_2 , $S^+R^7R^8$, PR^7 , $P^+R^7R^8$, phenylene, heterocycle, quaternary heterocycle, quaternary heteroaryl, or aryl.

wherein alkane diyl, alkene diyl, alkynes diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, peptide, and polypeptide can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo , OR^{13} , $NR^{13}R^{14}$, SR^{13} , $S(O)R^{13}$, SO_2R^{13} , SO_3R^{13} , $NR^{13}OR^{14}$, $NR^{13}NR^{14}R^{15}$, NO_2 , CO_2R^{13} , CN , OM , SO_2OM , $SO_2NR^{13}R^{14}$, $C(O)NR^{13}R^{14}$, $C(O)OM$, COR^{13} , $P(O)R^{13}R^{14}$, $P^+R^{13}R^{14}R^{15}A^-$, $P(OR^{13})OR^{14}$, $S^+R^{13}R^{14}A^-$, and $N^+R^{13}R^{14}R^{15}A^-$.



In any of the dimeric or multimeric structures discussed immediately above, benzothiepine compounds of the present invention can be used alone or in various combinations.

In any of the compounds of the present invention, R^1 and R^2 can be ethyl/butyl or butyl/butyl.

In another aspect, the present invention provides a pharmaceutical composition for the prophylaxis or treatment of a disease or condition for which a bile acid transport inhibitor is indicated, such as a hyperlipidemic condition, for example, atherosclerosis. Such compositions comprise any of the compounds disclosed above, alone or in combination, in an amount effective to reduce bile acid levels in the blood, or to reduce transport thereof across digestive system membranes, and a pharmaceutically acceptable carrier, excipient, or diluent.

In a further aspect, the present invention also provides a method of treating a disease or condition in

mammals, including humans, for which a bile acid transport inhibitor is indicated, comprising administering to a patient in need thereof a compound of the present invention in an effective amount in unit dosage form or in divided doses.

In yet a further aspect, the present invention also provides processes for the preparation of compounds of the present invention.

Further scope of the applicability of the present invention will become apparent from the detailed description provided below. However, it should be understood that the following detailed description and examples, while indicating preferred embodiments of the invention, are given by way of illustration only since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

DETAILED DESCRIPTION OF THE INVENTION

The following detailed description is provided to aid those skilled in the art in practicing the present invention. Even so, this detailed description should not be construed to unduly limit the present invention as modifications and variations in the embodiments discussed herein can be made by those of ordinary skill in the art without departing from the spirit or scope of the present inventive discovery.

The contents of each of the references cited herein, including the contents of the references cited within these primary references, are herein incorporated by reference in their entirety.

Definitions

In order to aid the reader in understanding the following detailed description, the following definitions are provided:

5 "Alkyl", "alkenyl", and "alkynyl" unless otherwise noted are each straight chain or branched chain hydrocarbons of from one to twenty carbons for alkyl or two to twenty carbons for alkenyl and alkynyl in the present invention and therefore mean, for example, methyl, ethyl, propyl, butyl, pentyl or hexyl and ethenyl, propenyl, butenyl, pentenyl, or hexenyl and ethynyl, propynyl, butynyl, pentynyl, or hexynyl respectively and isomers thereof.

10 "Aryl" means a fully unsaturated mono- or multi-ring carbocycle, including, but not limited to, substituted or unsubstituted phenyl, naphthyl, or anthracenyl.

15 "Heterocycle" means a saturated or unsaturated mono- or multi-ring carbocycle wherein one or more carbon atoms can be replaced by N, S, P, or O. This includes, for example, the following structures:



20 wherein Z, Z', Z" or Z''' is C, S, P, O, or N, with the proviso that one of Z, Z', Z" or Z''' is other than carbon, but is not O or S when attached to another Z

atom by a double bond or when attached to another O or S atom. Furthermore, the optional substituents are understood to be attached to Z, Z', Z" or Z''' only when each is C.

5 The term "heteroaryl" means a fully unsaturated heterocycle.

In either "heterocycle" or "heteroaryl," the point of attachment to the molecule of interest can be at the heteroatom or elsewhere within the ring.

10 The term "quaternary heterocycle" means a heterocycle in which one or more of the heteroatoms, for example, O, N, S, or P, has such a number of bonds that it is positively charged. The point of attachment of the quaternary heterocycle to the molecule of interest can be at a heteroatom or elsewhere.

15 The term "quaternary heteroaryl" means a heteroaryl in which one or more of the heteroatoms, for example, O, N, S, or P, has such a number of bonds that it is positively charged. The point of attachment of the quaternary heteroaryl to the molecule of interest can be at a heteroatom or elsewhere.

The term "halogen" means a fluoro, chloro, bromo or iodo group.

25 The term "haloalkyl" means alkyl substituted with one or more halogens.

The term "cycloalkyl" means a mono- or multi-ringed carbocycle wherein each ring contains three to ten carbon atoms, and wherein any ring can contain one or more double or triple bonds.

30 The term "diyl" means a diradical moiety wherein said moiety has two points of attachment to molecules of interest.

The term "oxo" means a doubly bonded oxygen.

The term "polyalkyl" means a branched or straight hydrocarbon chain having a molecular weight up to about 20,000, more preferably up to about 10,000, most preferably up to about 5,000.

The term "polyether" means a polyalkyl wherein one or more carbons are replaced by oxygen, wherein the polyether has a molecular weight up to about 20,000, more preferably up to about 10,000, most preferably up to about 5,000.

The term "polyalkoxy" means a polymer of alkylene oxides, wherein the polyalkoxy has a molecular weight up to about 20,000, more preferably up to about 10,000, most preferably up to about 5,000.

The term "cycloalkylidene" means a mono- or multi-ringed carbocycle wherein a carbon within the ring structure is doubly bonded to an atom which is not within the ring structures.

The term "carbohydrate" means a mono-, di-, tri-, or polysaccharide wherein the polysaccharide can have a molecular weight of up to about 20,000, for example, hydroxypropyl-methylcellulose or chitosan.

The term "peptide" means polyamino acid containing up to about 100 amino acid units.

The term "polypeptide" means polyamino acid containing from about 100 amino acid units to about 1000 amino acid units, more preferably from about 100 amino acid units to about 750 amino acid units, most preferably from about 100 amino acid units to about 500 amino acid units.

The term "alkylammoniumalkyl" means a NH_3^+ group or a mono-, di- or tri-substituted amino group, any of

which is bonded to an alkyl wherein said alkyl is bonded to the molecule of interest.

The term "triazolyl" includes all positional isomers. In all other heterocycles and heteroaryl's which contain more than one ring heteroatom and for which isomers are possible, such isomers are included in the definition of said heterocycles and heteroaryl's.

The term "sulfoalkyl" means an alkyl group to which a sulfonate group is bonded, wherein said alkyl is bonded to the molecule of interest.

The term "active compound" means a compound of the present invention which inhibits transport of bile acids.

When used in combination, for example "alkylaryl" or "arylalkyl," the individual terms listed above have the meaning indicated above.

The term "a bile acid transport inhibitor" means a compound capable of inhibiting absorption of bile acids from the intestine into the circulatory system of a mammal, such as a human. This includes increasing the fecal excretion of bile acids, as well as reducing the blood plasma or serum concentrations of cholesterol and cholesterol ester, and more specifically, reducing LDL and VLDL cholesterol. Conditions or diseases which benefit from the prophylaxis or treatment by bile acid transport inhibition include, for example, a hyperlipidemic condition such as atherosclerosis.

Compounds

The compounds of the present invention can have at least two asymmetrical carbon atoms, and therefore include racemates and stereoisomers, such as

diastereomers and enantiomers, in both pure form and in admixture. Such stereoisomers can be prepared using conventional techniques, either by reacting enantiomeric starting materials, or by separating isomers of compounds of the present invention.

Isomers may include geometric isomers, for example *cis* isomers or *trans* isomers across a double bond. All such isomers are contemplated among the compounds of the present invention.

The compounds of the present invention also include tautomers.

The compounds of the present invention as discussed below include their salts, solvates and prodrugs.

Compound Syntheses


The starting materials for use in the preparation of the compounds of the invention are known or can be prepared by conventional methods known to a skilled person or in an analogous manner to processes described in the art.

Generally, the compounds of the present invention can be prepared by the procedures described below.


For example, as shown in Scheme I, reaction of aldehyde II with formaldehyde and sodium hydroxide yields the hydroxyaldehyde III which is converted to mesylate IV with methanesulfonyl chloride and triethylamine similar to the procedure described in Chem. Ber. 98, 728-734 (1965). Reaction of mesylate IV with thiophenol V, prepared by the procedure described in WO 93/16055, in the presence of triethylamine yields keto-aldehyde VI which can be cyclized with the

reagent, prepared from zinc and titanium trichloride in refluxing ethylene glycol dimethyl ether (DME), to give a mixture of 2,3-dihydrobenzothiepine VII and two racemic stereoisomers of benzothiepin-(5H)-4-one VIII when R' and R' are nonequivalent. Oxidation of VII with 3 equivalents of m-chloro-perbenzoic acid (MCPBA) gives isomeric sulfone-epoxides IX which upon hydrogenation with palladium on carbon as the catalyst yield a mixture of four racemic stereoisomers of 4-hydroxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxides X and two racemic stereoisomers of 2,3,4,5-tetrahydrobenzothiepine-1,1-dioxides XI when R' and R' are nonequivalent.


Optically active compounds of the present invention can be prepared by using optically active starting material III or by resolution of compounds X with optical resolution agents well known in the art as described in *J. Org. Chem.*, 39, 3904 (1974), *ibid.*, 42, 2781 (1977), and *ibid.*, 44, 4891 (1979).




Alternatively, keto-aldehyde VI where R¹ is H can be prepared by reaction of thiophenol V with a 2-substituted acrolein.



Benzothiepin-(5H)-4-one VIII can be oxidized with MCPBA to give the benzothiepin-(5H)-4-one-1,1-dioxide XII which can be reduced with sodium borohydride to give four racemic stereoisomers of X. The two stereoisomers of X, Xa and Xb, having the OH group and R' on the opposite sides of the benzothiepine ring can be converted to the other two isomers of X, Xc and Xd, having the OH group and R' on the same side of the benzothiepine ring by reaction in methylene chloride with 40-50% sodium hydroxide in the presence of a phase transfer catalyst (PTC). The transformation can also be carried out with potassium t-butoxide in THF.





The compounds of the present invention where R' is OR, NRR' and S(O)₂R and R' is hydroxy can be prepared by reaction of epoxide IX where R' is H with thiol, alcohol, and amine in the presence of a base.




Another route to Xc and Xd of the present invention is shown in Scheme 2. Compound VI is oxidized to compound XIII with two equivalent of m-chloroperbenzoic acid. Hydrogenolysis of compound XIV with palladium on carbon yields compound XIV which can be cyclized with either potassium t-butoxide or sodium hydroxide under phase transfer conditions to a mixture of Xc and Xd. Separation of Xc and Xd can be accomplished by either HPLC or fractional crystallization.


The thiophenols XVIII and V used in the present invention can also be prepared according to the Scheme 3. Alkylation of phenol XV with an arylmethyl chloride in a nonpolar solvent according to the procedure in *J. Chem. Soc.*, 2431-2432 (1958) gives the ortho substituted phenol XVI. The phenol XVI can be converted to the thiophenol XVIII via the thiocarbamate XVII by the procedure described in *J. Org. Chem.*, 31, 3980 (1966). The phenol XVI is first reacted with dimethyl thiocarbamoyl chloride and triethylamine to give thiocarbamate XVII which is thermally rearranged at 200-300 °C, and the rearranged product is hydrolyzed with sodium hydroxide to yield the thiophenol XVIII. Similarly, Thiophenol V can also be prepared from 2-acylphenol XIX via the intermediate thiocarbamate XX.



Scheme 4 shows another route to benzothiepine-1,1-dioxides Xc and Xd starting from the thiophenol XVIII. Compound XVIII can be reacted with mesylate IV to give the sulfide-aldehyde XXI. Oxidation of XXI with two equivalents of MCPBA yields the sulfone-aldehyde XIV which can be cyclized with potassium t-butoxide to a mixture of Xc and Xd. Cyclization of sulfide-aldehyde with potassium t-butoxide also gives a mixture of benzothiepine XXIIc and XXIIId.




In Scheme 6, reduction of the sulfone-aldehyde XIV with hydrogen followed by reductive alkylation of the resulting amino derivative with hydrogen and an aldehyde catalyzed by palladium on carbon in the same reaction vessel yields the substituted amine derivative



XXVIII. Cyclization of XXVIII with potassium *t*-butoxide yields a mixture of substituted amino derivatives of this invention XIXc and XIXd.

Scheme 7 describes one of the methods of introducing a substituent to the aryl ring at the 5-position of benzochispiene. Iodination of 5-phenyl derivative XXX with iodine catalyzed by mercuric triflate gives the iodo derivative XXXI, which upon palladium-catalyzed carbonylation in an alcohol yields the carboxylate XXXII. Hydrolysis of the carboxylate



and derivatization of the resulting acid to acid derivatives are well known in the art.

Abbreviations used in the foregoing description have the following meanings:

THF---tetrahydrofuran

PTC---phase transfer catalyst

Aliquat 336---methyltricaprylammonium chloride

MCPBA---m-chloroperbenzoic acid

Celite---a brand of diatomaceous earth filtering aid

DMF---dimethylformamide

DME---ethylene glycol dimethyl ether

BOC---t-butoxycarbonyl group

R' and R' can be selected from among substituted

and unsubstituted C₁ to C₁₀ alkyl wherein the

substituent(s) can be selected from among

alkylcarbonyl, alkoxy, hydroxy, and nitrogen-containing

heterocycles joined to the C₁ to C₁₀ alkyl through an

ether linkage. Substituents at the 3-carbon can

include ethyl, n-propyl, n-butyl, n-pentyl, isobutyl,

isopropyl, -CH₂C(=O)C₂H₅, -CH₂OC₂H₅, and -CH₂O-(4-

picoline). Ethyl, n-propyl, n-butyl, and isobutyl are

preferred. In certain particularly preferred

compounds of the present invention, substituents R' and

R' are identical, for example n-butyl/n-butyl, so that

the compound is achiral at the 3-carbon. Eliminating

optical isomerism at the 3-carbon simplifies the

selection, synthesis, separation, and quality control

of the compound used as an ileal bile acid transport

inhibitor. In both compounds having a chiral 3-carbon

and those having an achiral 3-carbon, substituents (R')


on the benzo- ring can include hydrogen, aryl, alkyl,

hydroxy, halo, alkoxy, alkylthio, alkylsulfinyl,

alkylsulfonyl, haloalkyl, haloalkoxy, (N)-hydroxy-carbonylalkyl amine, haloalkylthio, haloalkylsulfinyl, haloalkylsulfonyl, amino, N-alkylamino, N,N-dialkylamino, (N)-alkoxycarbamoyl, (N)-aryloxycarbamoyl, (N)-aralkyloxycarbamoyl, trialkylammonium (especially with a halide counterion), (N)-amido, (N)-alkylamido, -N-alkylamido, -N,N-dialkylamido, (N)-haloalkylamido, (N)-sulfonamido, (N)-alkylsulfonamido, (N)-haloalkylsulfonamido, carboxyalkyl-amino, trialkylammonium salt, (N)-carbamic acid, alkyl or benzyl ester, N-acylamino, hydroxylamine, haloacylamino, carboxylate, thiophene a trialkyl ammonium salt having a carboxylic acid or hydroxy substituent on one or more of the alkyl substituents, an alkylene bridge having a quaternary ammonium salt substituted thereon, $-\{O(CH_2)_x\}-X$ where x is 2 to 12, w is 2 or 3 and X is a halo or a quaternary ammonium salt, and (N)-nitrogen containing heterocycle wherein the nitrogen of said heterocycle is optionally quaternized. Among the preferred species which may constitute R' are methyl, ethyl, isopropyl, t-butyl, hydroxy, methoxy, ethoxy, isopropoxy, methylthio, iodo, bromo, fluoro, methylsulfinyl, methylsulfonyl, ethylthio, amino, hydroxylamine, N-methylamino, N,N-dimethylamino, N,N-diethylamino, (N)-benzyloxycarbamoyl, trimethylammonium A', -NHC(=O)CH₃, -NHC(=O)C₂H₅, -NHC(=O)C₃H₇, carboxyethylamino, (N)-morpholinyl, (N)-azetidinyl, (N)-N-methylazetidinium A', (N)-pyrrolidinyl, pyrrolyl, (N)-N-methylpyridinium A', (N)-N-methylmorpholinium A', and N,N'-methylpiperazinyl, (N)-bromomethylamido, (N)-

N-hexylamino, thiophene, -N'(CH₃)₂CO₂H I', -NCH₂CH₂CO₂H, - (N)-N'-dimethylpiperazinium I', (N)-t-butyloxycarbamoyl, (N)-methylsulfonamido, (N)-N'-methylpyrrolidinium, and -(OCH₂CH₂)₁I', where A' is a pharmaceutically acceptable anion. The benzo ring is can be mono-substituted at the 6, 7 or 8 position, or disubstituted at the 7- and -8 positions. Also included are the 6,7,8-trialkoxo compounds, for example the 6,7,8-trimethoxy compounds. A variety of other substituents can be advantageously present on the 6, 7, 8, and/or 9- positions of the benzo ring, including, for example, guanidiny, cycloalkyl, carboxylate (e.g., a 5 or 6 carbon monosaccharide), peptide, and quaternary ammonium salts linked to the ring via poly(oxyalkylene) linkages, e.g., -(OCH₂CH₂)₂-N⁺R¹R²R³A', where x is 2 to 10. Exemplary compounds are those set forth below in Table 1.

TABLE 1
Alternative compounds #3 (Family F101.xxx.yyy) *



Prefix (yyy.xxx.yyy)	Cpds (yyy.xxx.yyy)	R ¹ -R ²	R ³	R ⁴ q
F101.001	01	n-propyl	Ph-	7-methyl
	02	n-propyl	Ph-	7-ethyl
	03	n-propyl	Ph-	7-iso-propyl
	04	n-propyl	Ph-	7-tert-butyl
	05	n-propyl	Ph-	7-OH
	06	n-propyl	Ph-	7-OCH ₃
	07	n-propyl	Ph-	7-O(1-iso-propyl)
	08	n-propyl	Ph-	7-SCH ₃
	09	n-propyl	Ph-	7-SOCH ₃
	10	n-propyl	Ph-	7-SO ₂ CH ₃

* General Notes

In the description of the substituents (N) indicates that a nitrogen bearing substituent is bonded to the ring structure via the nitrogen atom.

Similarly, 2-thiophene indicates a bond in the 2 position of the thiophene ring. A similar convention is used for other heterocyclic substituents.

Abbreviations and Definitions
NH-CBZ is defined as -HNC(=O)OCH₂Ph

11	n-propyl	Ph-	7-SCH ₂ CH ₃
12	n-propyl	Ph-	7-NH ₂
13	n-propyl	Ph-	7-NHOH
14	n-propyl	Ph-	7-NHCH ₃
15	n-propyl	Ph-	7-N(CH ₃) ₂
16	n-propyl	Ph-	7-N ⁺ (CH ₃) ₃ , I ⁻
17	n-propyl	Ph-	7-NHC(-O)CH ₃
18	n-propyl	Ph-	7-N(CH ₂ CH ₃) ₂
19	n-propyl	Ph-	7-NMeCH ₂ CO ₂ H
20	n-propyl	Ph-	7-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
21	n-propyl	Ph-	7-(N)-morpholine
22	n-propyl	Ph-	7-(N)-azetidine
23	n-propyl	Ph-	7-(N)-N-methylazetidinium, I ⁻
24	n-propyl	Ph-	7-(N)-pyrrolidine
25	n-propyl	Ph-	7-(N)-N-methyl-pyrrolidinium, I ⁻
26	n-propyl	Ph-	7-(N)-N-methyl-morpholinium, I ⁻
27	n-propyl	Ph-	7-(N)-N'-methylpiperazine
28	n-propyl	Ph-	7-(N)-N'-dimethylpiperazinium, I ⁻
29	n-propyl	Ph-	7-NH-CBZ
30	n-propyl	Ph-	7-NHC(O)C ₆ H ₁₁
31	n-propyl	Ph-	7-NHC(O)CH ₂ Br
32	n-propyl	Ph-	7-NH-C(NH)NH ₂
33	n-propyl	Ph-	7-(2)-thiophane
34	n-propyl	Ph-	8-methyl
35	n-propyl	Ph-	8-ethyl
36	n-propyl	Ph-	8-iso-propyl
37	n-propyl	Ph-	8-tert-butyl
38	n-propyl	Ph-	8-OH
39	n-propyl	Ph-	8-OCH ₃
40	n-propyl	Ph-	8-O(1-iso-propyl)
41	n-propyl	Ph-	8-SCH ₃
42	n-propyl	Ph-	8-SOCH ₃
43	n-propyl	Ph-	8-SO ₂ CH ₃
44	n-propyl	Ph-	8-SCH ₂ CH ₃
45	n-propyl	Ph-	8-NH ₂
46	n-propyl	Ph-	8-NHOH
47	n-propyl	Ph-	8-NHCH ₃
48	n-propyl	Ph-	8-N(CH ₃) ₂
49	n-propyl	Ph-	8-N ⁺ (CH ₃) ₃ , I ⁻
50	n-propyl	Ph-	8-NHC(-O)CH ₃
51	n-propyl	Ph-	8-N(CH ₂ CH ₃) ₂
52	n-propyl	Ph-	8-NMeCH ₂ CO ₂ H

53	n-propyl	Ph-	8-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
54	n-propyl	Ph-	8-(N)-morpholine
55	n-propyl	Ph-	8-(N)-azetidine
56	n-propyl	Ph-	8-(N)-N-methylazetidinium, I ⁺
57	n-propyl	Ph-	8-(N)-pyrrolidine
58	n-propyl	Ph-	8-(N)-N-methyl-pyrrolidinium, I ⁺
59	n-propyl	Ph-	8-(N)-N-methyl-morpholinium, I ⁺
60	n-propyl	Ph-	8-(N)-N'-methylpiperazine
61	n-propyl	Ph-	8-(N)-N'-dimethylpiperazinium, I ⁺
62	n-propyl	Ph-	8-NH-CBz
63	n-propyl	Ph-	8-NHC(O)C ₆ H ₁₁
64	n-propyl	Ph-	8-NHC(O)CH ₂ Br
65	n-propyl	Ph-	8-NH-C(NH)NH ₂
66	n-propyl	Ph-	8-(2)-chlorophene
67	n-propyl	Ph-	9-methyl
68	n-propyl	Ph-	9-ethyl
69	n-propyl	Ph-	9-iso-propyl
70	n-propyl	Ph-	9-tert-butyl
71	n-propyl	Ph-	9-OH
72	n-propyl	Ph-	9-OCH ₃
73	n-propyl	Ph-	9-O(1-iso-propyl)
74	n-propyl	Ph-	9-SCH ₃
75	n-propyl	Ph-	9-SOCH ₃
76	n-propyl	Ph-	9-SO ₂ CH ₃
77	n-propyl	Ph-	9-SCH ₂ CH ₃
78	n-propyl	Ph-	9-NH ₂
79	n-propyl	Ph-	9-NHOH
80	n-propyl	Ph-	9-NHCH ₃
81	n-propyl	Ph-	9-N(CH ₃) ₂
82	n-propyl	Ph-	9-N ⁺ (CH ₃) ₃ , I ⁻
83	n-propyl	Ph-	9-NHC(=O)CH ₃
84	n-propyl	Ph-	9-N(CH ₂ CH ₃) ₂
85	n-propyl	Ph-	9-NMeCH ₂ CO ₂ H
86	n-propyl	Ph-	9-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
87	n-propyl	Ph-	9-(N)-azetidine
88	n-propyl	Ph-	9-(N)-morpholine
89	n-propyl	Ph-	9-(N)-N-methylazetidinium, I ⁺
90	n-propyl	Ph-	9-(N)-pyrrolidine
91	n-propyl	Ph-	9-(N)-N-methyl-pyrrolidinium, I ⁺
92	n-propyl	Ph-	9-(N)-N-methyl-morpholinium, I ⁺
93	n-propyl	Ph-	9-(N)-N'-methylpiperazine
93	n-propyl	Ph-	9-(N)-N'-dimethylpiperazinium, I ⁺
95	n-propyl	Ph-	9-NH-CBz

41

Prefix	Cpd#	R ¹ -R ²	R ³	(R ³) _q
(FTT, XXX, YYY)				
F101.002	01	n-butyl	Ph-	7-methyl
	02	n-butyl	Ph-	7-ethyl
	03	n-butyl	Ph-	7-iso-propyl
	04	n-butyl	Ph-	7-tert-butyl
	05	n-butyl	Ph-	7-OH
	06	n-butyl	Ph-	7-OCH ₃
	07	n-butyl	Ph-	7-O(1-iso-propyl)
	08	n-butyl	Ph-	7-SCH ₃
	09	n-butyl	Ph-	7-SOCH ₃
	10	n-butyl	Ph-	7-SO ₂ CH ₃
	11	n-butyl	Ph-	7-SCH ₂ CH ₃
	12	n-butyl	Ph-	7-NH ₂
	13	n-butyl	Ph-	7-NHOH
	14	n-butyl	Ph-	7-NHCH ₃
	15	n-butyl	Ph-	7-N(CH ₃) ₂
	16	n-butyl	Ph-	7-N ⁺ (CH ₃) ₃ , I ⁻
	17	n-butyl	Ph-	7-NHC(=O)CH ₃
	18	n-butyl	Ph-	7-N(CH ₂ CH ₃) ₂
	19	n-butyl	Ph-	7-NMeCH ₂ CO ₂ H
	20	n-butyl	Ph-	7-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
	21	n-butyl	Ph-	7-(N)-morpholine
	22	n-butyl	Ph-	7-(N)-azetidine
	23	n-butyl	Ph-	7-(N)-N-methylazetidinium, I ⁺
	24	n-butyl	Ph-	7-(N)-pyrrolidine
	25	n-butyl	Ph-	7-(N)-N-methyl-pyrrolidinium, I ⁺
	26	n-butyl	Ph-	7-(N)-N-methyl-morpholinium, I ⁺
	27	n-butyl	Ph-	7-(N)-N'-methylpiperazine
	28	n-butyl	Ph-	7-(N)-N'-dimethylpiperazinium, I ⁺
	29	n-butyl	Ph-	7-NH-CBz
	30	n-butyl	Ph-	7-NHC(O)C ₆ H ₁₁
	31	n-butyl	Ph-	7-NHC(O)CH ₂ Br

42

32	n-butyl	Ph- 7-NH-C(NH)NH ₂	74	n-butyl	Ph- 9-SCH ₃
33	n-butyl	Ph- 7-(2)-thiophene	75	n-butyl	Ph- 9-SOCH ₃
34	n-butyl	Ph- 8-methyl	76	n-butyl	Ph- 9-SO ₂ CH ₃
35	n-butyl	Ph- 8-ethyl	77	n-butyl	Ph- 9-SCH ₂ CH ₃
36	n-butyl	Ph- 8-iso-propyl	78	n-butyl	Ph- 9-NH ₂
37	n-butyl	Ph- 8-tert-butyl	79	n-butyl	Ph- 9-NHOH
38	n-butyl	Ph- 8-OH	80	n-butyl	Ph- 9-NHCH ₃
39	n-butyl	Ph- 8-OCH ₃	81	n-butyl	Ph- 9-N(CH ₃) ₂
40	n-butyl	Ph- 8-O(iso-propyl)	82	n-butyl	Ph- 9-N ⁺ (CH ₃) ₃ , I ⁻
41	n-butyl	Ph- 8-SCH ₃	83	n-butyl	Ph- 9-NHC(=O)CH ₃
42	n-butyl	Ph- 8-SOCH ₃	84	n-butyl	Ph- 9-N(CH ₂ CH ₃) ₂
43	n-butyl	Ph- 8-SO ₂ CH ₃	85	n-butyl	Ph- 9-NMeCH ₂ CO ₂ H
44	n-butyl	Ph- 8-SCH ₂ CH ₃	86	n-butyl	Ph- 9-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
45	n-butyl	Ph- 8-NH ₂	87	n-butyl	Ph- 9-(N)-morpholine
46	n-butyl	Ph- 8-NHOH	88	n-butyl	Ph- 9-(N)-azetidine
47	n-butyl	Ph- 8-NHCH ₃	89	n-butyl	Ph- 9-(N)-N-methylazetidinium, I ⁻
48	n-butyl	Ph- 8-N(CH ₃) ₂	90	n-butyl	Ph- 9-(N)-pyrrolidine
49	n-butyl	Ph- 8-N ⁺ (CH ₃) ₃ , I ⁻	91	n-butyl	Ph- 9-(N)-N-methyl-pyrrolidinium, I ⁻
50	n-butyl	Ph- 8-NHC(=O)CH ₃	92	n-butyl	Ph- 9-(N)-N-methyl-morpholinium, I ⁻
51	n-butyl	Ph- 8-N(CH ₂ CH ₃) ₂	93	n-butyl	Ph- 9-(N)-N'-methylpiperazine
52	n-butyl	Ph- 8-NMeCH ₂ CO ₂ H	94	n-butyl	Ph- 9-(N)-N'-dimethylpiperazinium, I ⁻
53	n-butyl	Ph- 8-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻	95	n-butyl	Ph- 9-NH-Cbz
54	n-butyl	Ph- 8-(N)-morpholine	96	n-butyl	Ph- 9-NHC(O)C ₅ H ₁₁
55	n-butyl	Ph- 8-(N)-azetidine	97	n-butyl	Ph- 9-NHC(O)CH ₂ Br
56	n-butyl	Ph- 8-(N)-N-methylazetidinium, I ⁻	98	n-butyl	Ph- 9-NH-C(NH)NH ₂
57	n-butyl	Ph- 8-(N)-pyrrolidine	99	n-butyl	Ph- 9-(2)-thiophene
58	n-butyl	Ph- 8-(N)-N-methyl-pyrrolidinium, I ⁻	100	n-butyl	Ph- 7-OCH ₃ , 8-OCH ₃
59	n-butyl	Ph- 8-(N)-N-methyl-morpholinium, I ⁻	101	n-butyl	Ph- 7-SCH ₃ , 8-OCH ₃
60	n-butyl	Ph- 8-(N)-N'-methylpiperazine	102	n-butyl	Ph- 7-SCH ₃ , 8-SCH ₃
61	n-butyl	Ph- 8-(N)-N'-dimethylpiperazinium, I ⁻	103	n-butyl	Ph- 6-OCH ₃ , 7-OCH ₃ , 8-OCH ₃
62	n-butyl	Ph- 8-NH-Cbz			
63	n-butyl	Ph- 8-NHC(O)C ₅ H ₁₁			
64	n-butyl	Ph- 8-NHC(O)CH ₂ Br			
65	n-butyl	Ph- 8-NH-C(NH)NH ₂			
66	n-butyl	Ph- 8-(2)-thiophene			
67	n-butyl	Ph- 9-methyl			
68	n-butyl	Ph- 9-ethyl			
69	n-butyl	Ph- 9-iso-propyl			
70	n-butyl	Ph- 9-tert-butyl			
71	n-butyl	Ph- 9-OH			
72	n-butyl	Ph- 9-OCH ₃			
73	n-butyl	Ph- 9-O(iso-propyl)			

10	n-pentyl	Ph- 7-SO ₂ CH ₃	52	n-pentyl	Ph- 8-NMeCH ₂ CO ₂ H
11	n-pentyl	Ph- 7-SCH ₂ CH ₃	53	n-pentyl	Ph- 8-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
12	n-pentyl	Ph- 7-NH ₂	54	n-pentyl	Ph- 8-(N)-morpholine
13	n-pentyl	Ph- 7-NHOH	55	n-pentyl	Ph- 8-(N)-azetidine
14	n-pentyl	Ph- 7-NHCH ₃	56	n-pentyl	Ph- 8-(N)-N-methylazetidinium, I ⁺
15	n-pentyl	Ph- 7-N(CH ₃) ₂	57	n-pentyl	Ph- 8-(N)-pyrrolidine
16	n-pentyl	Ph- 7-N ⁺ (CH ₃) ₃ , I ⁻	58	n-pentyl	Ph- 8-(N)-N-methyl-pyrrolidinium, I ⁺
17	n-pentyl	Ph- 7-NHC(=O)CH ₃	59	n-pentyl	Ph- 8-(N)-N-methyl-morpholinium, I ⁺
18	n-pentyl	Ph- 7-N(CH ₂ CH ₃) ₂	60	n-pentyl	Ph- 8-(N)-N'-methylpiperazine
19	n-pentyl	Ph- 7-NMeCH ₂ CO ₂ H	61	n-pentyl	Ph- 8-(N)-N'-dimethylpiperazinium, I ⁺
20	n-pentyl	Ph- 7-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻	62	n-pentyl	Ph- 8-NH-CBZ
21	n-pentyl	Ph- 7-(N)-morpholine	63	n-pentyl	Ph- 8-NHC(=O)C ₆ H ₁₁
22	n-pentyl	Ph- 7-(N)-azetidine	64	n-pentyl	Ph- 8-NHC(=O)CH ₂ Br
23	n-pentyl	Ph- 7-(N)-N-methylazetidinium, I ⁺	65	n-pentyl	Ph- 8-NH-C(NH) ₂ NH ₂
24	n-pentyl	Ph- 7-(N)-pyrrolidine	66	n-pentyl	Ph- 8-(2)-thiophene
25	n-pentyl	Ph- 7-(N)-N-methyl-pyrrolidinium, I ⁺	67	n-pentyl	Ph- 9-methyl
26	n-pentyl	Ph- 7-(N)-N-methyl-morpholinium, I ⁺	68	n-pentyl	Ph- 9-ethyl
27	n-pentyl	Ph- 7-(N)-N'-methylpiperazine	69	n-pentyl	Ph- 9-iso-propyl
28	n-pentyl	Ph- 7-(N)-N'-methylpiperazine	70	n-pentyl	Ph- 9-tert-butyl
29	n-pentyl	Ph- 7-(N)-N'-dimethylpiperazinium, I ⁺	71	n-pentyl	Ph- 9-OH
30	n-pentyl	Ph- 7-NH-CBZ	72	n-pentyl	Ph- 9-OCH ₃
31	n-pentyl	Ph- 7-NHC(=O)C ₆ H ₁₁	73	n-pentyl	Ph- 9-O(=O)-propyl
32	n-pentyl	Ph- 7-NHC(=O)CH ₂ Br	74	n-pentyl	Ph- 9-SCH ₃
33	n-pentyl	Ph- 7-NH-C(NH) ₂ NH ₂	75	n-pentyl	Ph- 9-SOCH ₃
		Ph- 7-(2)-thiophene	76	n-pentyl	Ph- 9-SO ₂ CH ₃
34	n-pentyl	Ph- 8-methyl	77	n-pentyl	Ph- 9-SCH ₂ CH ₃
35	n-pentyl	Ph- 8-ethyl	78	n-pentyl	Ph- 9-NH ₂
36	n-pentyl	Ph- 8-iso-propyl	79	n-pentyl	Ph- 9-NHOH
37	n-pentyl	Ph- 8-tert-butyl	80	n-pentyl	Ph- 9-NHCH ₃
38	n-pentyl	Ph- 8-OH	81	n-pentyl	Ph- 9-N(CH ₃) ₂
39	n-pentyl	Ph- 8-OCH ₃	82	n-pentyl	Ph- 9-N ⁺ (CH ₃) ₃ , I ⁻
40	n-pentyl	Ph- 8-O(=O)-propyl	83	n-pentyl	Ph- 9-NHC(=O)CH ₃
41	n-pentyl	Ph- 8-SCH ₃	84	n-pentyl	Ph- 9-N(CH ₂ CH ₃) ₂
42	n-pentyl	Ph- 8-SOCH ₃	85	n-pentyl	Ph- 9-NMeCH ₂ CO ₂ H
43	n-pentyl	Ph- 8-SO ₂ CH ₃	86	n-pentyl	Ph- 9-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁺
44	n-pentyl	Ph- 8-SCH ₂ CH ₃	87	n-pentyl	Ph- 9-(N)-morpholine
45	n-pentyl	Ph- 8-NH ₂	88	n-pentyl	Ph- 9-(N)-azetidine
46	n-pentyl	Ph- 8-NHOH	89	n-pentyl	Ph- 9-(N)-N-methylazetidinium, I ⁺
47	n-pentyl	Ph- 8-NHCH ₃	90	n-pentyl	Ph- 9-(N)-pyrrolidine
48	n-pentyl	Ph- 8-N(CH ₃) ₂	91	n-pentyl	Ph- 9-(N)-N-methyl-pyrrolidinium, I ⁺
49	n-pentyl	Ph- 8-N ⁺ (CH ₃) ₃ , I ⁻	92	n-pentyl	Ph- 9-(N)-N-methyl-morpholinium, I ⁺
50	n-pentyl	Ph- 8-NHC(=O)CH ₃	93	n-pentyl	Ph- 9-(N)-N'-methylpiperazine
51	n-pentyl	Ph- 8-N(CH ₂ CH ₃) ₂	93	n-pentyl	Ph- 9-(N)-N'-dimethylpiperazinium, I ⁺

Prefix (FTT, xxx, yyy)	Cpd#	R ¹ -R ²	R ³	(R ³) _q
F101.004	01	n-hexyl	Ph- 7-methyl	
	02	n-hexyl	Ph- 7-ethyl	
	03	n-hexyl	Ph- 7-iso-propyl	
	04	n-hexyl	Ph- 7-tert-butyl	
	05	n-hexyl	Ph- 7-OH	
	06	n-hexyl	Ph- 7-OCH ₃	
	07	n-hexyl	Ph- 7-O(1-iso-propyl)	
	08	n-hexyl	Ph- 7-SCH ₃	
	09	n-hexyl	Ph- 7-SO ₂ CH ₃	
	10	n-hexyl	Ph- 7-SCH ₂ CH ₃	
	11	n-hexyl	Ph- 7-NH ₂	
	12	n-hexyl	Ph- 7-NHOH	
	13	n-hexyl	Ph- 7-NHCH ₃	
	14	n-hexyl	Ph- 7-N(CH ₃) ₂	
	15	n-hexyl	Ph- 7-N ⁺ (CH ₃) ₃ , I ⁻	
	16	n-hexyl	Ph- 7-NHC(=O)CH ₃	
	17	n-hexyl	Ph- 7-NHC(=O)CH ₂ CH ₃	
	18	n-hexyl	Ph- 7-NHC(=O)CH ₂ CO ₂ H	
	19	n-hexyl	Ph- 7-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻	
	20	n-hexyl	Ph- 7-(N)-morpholine	
	21	n-hexyl	Ph- 7-(N)-azetidine	
	22	n-hexyl	Ph- 7-(N)-N-methylazetidinium, I ⁻	
	23	n-hexyl	Ph- 7-(N)-pyrrolidine	
	24	n-hexyl	Ph- 7-(N)-N-methyl-pyrrolidinium, I ⁻	
	25	n-hexyl	Ph- 7-(N)-N-methyl-morpholinium, I ⁻	
	26	n-hexyl	Ph- 7-(N)-N'-methylpiperazine	
	27	n-hexyl	Ph- 7-(N)-N'-dimethylpiperazinium, I ⁻	
	28	n-hexyl	Ph- 7-NH-CBZ	
	29	n-hexyl	Ph- 7-NHC(O)C ₅ H ₁₁	
	30	n-hexyl		

47

31	n-hexyl	Ph- 7-NHC(O)CH ₂ Br
32	n-hexyl	Ph- 7-NH-C(NH)NH ₂
33	n-hexyl	Ph- 7-(2)-thiophene
34	n-hexyl	Ph- 8-methyl
35	n-hexyl	Ph- 8-ethyl
36	n-hexyl	Ph- 8-iso-propyl
37	n-hexyl	Ph- 8-tert-butyl
38	n-hexyl	Ph- 8-OH
39	n-hexyl	Ph- 8-OCH ₃
40	n-hexyl	Ph- 8-O(1-iso-propyl)
41	n-hexyl	Ph- 8-SCH ₃
42	n-hexyl	Ph- 8-SOCH ₃
43	n-hexyl	Ph- 8-SO ₂ CH ₃
44	n-hexyl	Ph- 8-SCH ₂ CH ₃
45	n-hexyl	Ph- 8-NH ₂
46	n-hexyl	Ph- 8-NHOH
47	n-hexyl	Ph- 8-NHCH ₃
48	n-hexyl	Ph- 8-N(CH ₃) ₂
49	n-hexyl	Ph- 8-N ⁺ (CH ₃) ₃ , I ⁻
50	n-hexyl	Ph- 8-NHC(=O)CH ₃
51	n-hexyl	Ph- 8-N(CH ₂ CH ₃) ₂
52	n-hexyl	Ph- 8-NHC(=O)CH ₂ CO ₂ H
53	n-hexyl	Ph- 8-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
54	n-hexyl	Ph- 8-(N)-morpholine
55	n-hexyl	Ph- 8-(N)-azetidine
56	n-hexyl	Ph- 8-(N)-N-methylazetidinium, I ⁻
57	n-hexyl	Ph- 8-(N)-pyrrolidine
58	n-hexyl	Ph- 8-(N)-N-methyl-pyrrolidinium, I ⁻
59	n-hexyl	Ph- 8-(N)-N-methyl-morpholinium, I ⁻
60	n-hexyl	Ph- 8-(N)-N'-methylpiperazine
61	n-hexyl	Ph- 8-(N)-N'-dimethylpiperazinium, I ⁻
62	n-hexyl	Ph- 8-NH-CBZ
63	n-hexyl	Ph- 8-NHC(O)C ₅ H ₁₁
64	n-hexyl	Ph- 8-NHC(O)CH ₂ Br
65	n-hexyl	Ph- 8-NH-C(NH)NH ₂
66	n-hexyl	Ph- 8-(2)-thiophene
67	n-hexyl	Ph- 9-methyl
68	n-hexyl	Ph- 9-ethyl
69	n-hexyl	Ph- 9-iso-propyl
70	n-hexyl	Ph- 9-tert-butyl
71	n-hexyl	Ph- 9-OH
72	n-hexyl	Ph- 9-OCH ₃

48

73	n-hexyl	Ph-	9-O-(iso-propyl)	09	iso-propyl	Ph-	7-SOCH ₃
74	n-hexyl	Ph-	9-SCH ₃	10	iso-propyl	Ph-	7-SO ₂ CH ₃
75	n-hexyl	Ph-	9-SOCH ₃	11	iso-propyl	Ph-	7-SCH ₂ CH ₃
76	n-hexyl	Ph-	9-SO ₂ CH ₃	12	iso-propyl	Ph-	7-NH ₂
77	n-hexyl	Ph-	9-SCH ₂ CH ₃	13	iso-propyl	Ph-	7-NHOH
78	n-hexyl	Ph-	9-NH ₂	14	iso-propyl	Ph-	7-NHCH ₃
79	n-hexyl	Ph-	9-NHOH	15	iso-propyl	Ph-	7-N(CH ₃) ₂
80	n-hexyl	Ph-	9-NHCH ₃	16	iso-propyl	Ph-	7-N ⁺ (CH ₃) ₃ , I ⁻
81	n-hexyl	Ph-	9-N(CH ₃) ₂	17	iso-propyl	Ph-	7-NHC(=O)CH ₃
82	n-hexyl	Ph-	9-N ⁺ (CH ₃) ₃ , I ⁻	18	iso-propyl	Ph-	7-N(CH ₂ CH ₃) ₂
83	n-hexyl	Ph-	9-NHC(=O)CH ₃	19	iso-propyl	Ph-	7-NMeCH ₂ CO ₂ H
84	n-hexyl	Ph-	9-N(CH ₂ CH ₃) ₂	20	iso-propyl	Ph-	7-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
85	n-hexyl	Ph-	9-NMeCH ₂ CO ₂ H	21	iso-propyl	Ph-	7-(N)-morpholine
86	n-hexyl	Ph-	9-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻	22	iso-propyl	Ph-	7-(N)-azetidine
87	n-hexyl	Ph-	9-(N)-morpholine	23	iso-propyl	Ph-	7-(N)-N-methylazetidinium, I ⁺
88	n-hexyl	Ph-	9-(N)-azetidine	24	iso-propyl	Ph-	7-(N)-pyrrolidine
89	n-hexyl	Ph-	9-(N)-N-methylazetidinium, I ⁺	25	iso-propyl	Ph-	7-(N)-N-methyl-pyrrolidinium, I ⁺
90	n-hexyl	Ph-	9-(N)-pyrrolidine	26	iso-propyl	Ph-	7-(N)-N-methyl-morpholinium, I ⁺
91	n-hexyl	Ph-	9-(N)-N-methyl-pyrrolidinium, I ⁺	27	iso-propyl	Ph-	7-(N)-N-methyl-piperazine
92	n-hexyl	Ph-	9-(N)-N-methyl-morpholinium, I ⁺	28	iso-propyl	Ph-	7-(N)-N'-dimethylpiperazinium, I ⁺
93	n-hexyl	Ph-	9-(N)-N'-methylpiperazine	29	iso-propyl	Ph-	7-NH-CBZ
94	n-hexyl	Ph-	9-(N)-N'-dimethylpiperazinium, I ⁺	30	iso-propyl	Ph-	7-NHC(=O)C ₆ H ₁₁
95	n-hexyl	Ph-	9-NH-CBZ	31	iso-propyl	Ph-	7-NHC(=O)CH ₂ Br
96	n-hexyl	Ph-	9-NHC(=O)C ₆ H ₁₁	32	iso-propyl	Ph-	7-NH-C(NH) ₂ NH ₂
97	n-hexyl	Ph-	9-NHC(=O)CH ₂ Br	33	iso-propyl	Ph-	7-(2)-chlorophene
98	n-hexyl	Ph-	9-NH-C(NH) ₂ NH ₂				
99	n-hexyl	Ph-	9-(2)-chlorophene				
100	n-hexyl	Ph-	7-OCH ₃ , 8-OCH ₃	34	iso-propyl	Ph-	8-methyl
101	n-hexyl	Ph-	7-SCH ₃ , 8-OCH ₃	35	iso-propyl	Ph-	8-ethyl
102	n-hexyl	Ph-	7-SCH ₃ , 8-SCH ₃	36	iso-propyl	Ph-	8-iso-propyl
103	n-hexyl	Ph-	6-OCH ₃ , 7-OCH ₃ , 8-OCH ₃	37	iso-propyl	Ph-	8-tert-butyl
				38	iso-propyl	Ph-	8-OH
				39	iso-propyl	Ph-	8-OCH ₃
				40	iso-propyl	Ph-	8-O(1-iso-propyl)
				41	iso-propyl	Ph-	8-SCH ₃
				42	iso-propyl	Ph-	8-SOCH ₃
				43	iso-propyl	Ph-	8-SO ₂ CH ₃
				44	iso-propyl	Ph-	8-SCH ₂ CH ₃
				45	iso-propyl	Ph-	8-NH ₂
				46	iso-propyl	Ph-	8-NHOH
				47	iso-propyl	Ph-	8-NHCH ₃
				48	iso-propyl	Ph-	8-N(CH ₃) ₂
				49	iso-propyl	Ph-	8-N ⁺ (CH ₃) ₃ , I ⁻
				50	iso-propyl	Ph-	8-NHC(=O)CH ₃

51	iso-propyl	Ph-	8-N(CH ₂ CH ₃) ₂
52	iso-propyl	Ph-	8-NMeCH ₂ CO ₂ H
53	iso-propyl	Ph-	8-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
54	iso-propyl	Ph-	8-(N)-morpholine
55	iso-propyl	Ph-	8-(N)-azetidine
56	iso-propyl	Ph-	8-(N)-N-methylazetidinium, I ⁻
57	iso-propyl	Ph-	8-(N)-pyrrolidine
58	iso-propyl	Ph-	8-(N)-N-methyl-pyrrolidinium, I ⁻
59	iso-propyl	Ph-	8-(N)-N-methyl-morpholinium, I ⁻
60	iso-propyl	Ph-	8-(N)-N'-methylpiperazine
61	iso-propyl	Ph-	8-(N)-N'-dimethylpiperazinium, I ⁻
62	iso-propyl	Ph-	8-NH-CB ₂
63	iso-propyl	Ph-	8-NHC(O)C ₆ H ₁₁
64	iso-propyl	Ph-	8-NHC(O)CH ₂ Br
65	iso-propyl	Ph-	8-NH-C(NH) ₂ NH ₂
66	iso-propyl	Ph-	8-(2)-thiophene
67	iso-propyl	Ph-	9-methyl
68	iso-propyl	Ph-	9-ethyl
69	iso-propyl	Ph-	9-iso-propyl
70	iso-propyl	Ph-	9-tert-butyl
71	iso-propyl	Ph-	9-OH
72	iso-propyl	Ph-	9-OCH ₃
73	iso-propyl	Ph-	9-O(iso-propyl)
74	iso-propyl	Ph-	9-SCH ₃
75	iso-propyl	Ph-	9-SOCH ₃
76	iso-propyl	Ph-	9-SO ₂ CH ₃
77	iso-propyl	Ph-	9-SCH ₂ CH ₃
78	iso-propyl	Ph-	9-NH ₂
79	iso-propyl	Ph-	9-NHOH
80	iso-propyl	Ph-	9-NHCH ₃
81	iso-propyl	Ph-	9-N(CH ₃) ₂
82	iso-propyl	Ph-	9-N ⁺ (CH ₃) ₃ , I ⁻
83	iso-propyl	Ph-	9-NHC(=O)CH ₃
84	iso-propyl	Ph-	9-N(CH ₂ CH ₃) ₂
85	iso-propyl	Ph-	9-NMeCH ₂ CO ₂ H
86	iso-propyl	Ph-	9-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
87	iso-propyl	Ph-	9-(N)-morpholine
88	iso-propyl	Ph-	9-(N)-azetidine
89	iso-propyl	Ph-	9-(N)-N-methylazetidinium, I ⁻
90	iso-propyl	Ph-	9-(N)-pyrrolidine
91	iso-propyl	Ph-	9-(N)-N-methyl-pyrrolidinium, I ⁻
92	iso-propyl	Ph-	9-(N)-N-methyl-morpholinium, I ⁻
93	iso-propyl	Ph-	9-(N)-N'-methylpiperazine

Prefix	Cod#	R ¹ -R ²	R ⁵	(R ⁴) _q
(ITT. doc. YYY)				
F101.006				
93	iso-propyl	Ph-	9-(N)-N'-dimethylpiperazinium, I ⁻	
95	iso-propyl	Ph-	9-NK-CB2	
96	iso-propyl	Ph-	9-NHC(O)C ₃ H ₁₁	
97	iso-propyl	Ph-	9-NHC(O)CH ₂ Br	
98	iso-propyl	Ph-	9-NH-C(NH)NH ₂	
99	iso-propyl	Ph-	9-(2)-thiophene	
100	iso-propyl	Ph-	7-OCH ₃ , 8-OCH ₃	
101	iso-propyl	Ph-	7-SCH ₃ , 8-OCH ₃	
102	iso-propyl	Ph-	7-SCH ₃ , 8-SCH ₃	
103	iso-propyl	Ph-	6-OCH ₃ , 7-OCH ₃ , 8-OCH ₃	
01	iso-butyl	Ph-	7-methyl	
02	iso-butyl	Ph-	7-ethyl	
03	iso-butyl	Ph-	7-iso-propyl	
04	iso-butyl	Ph-	7-tert-butyl	
05	iso-butyl	Ph-	7-QH	
06	iso-butyl	Ph-	7-OC ₃ H ₇	
07	iso-butyl	Ph-	7-O(1iso-propyl)	
08	iso-butyl	Ph-	7-SCH ₃	
09	iso-butyl	Ph-	7-SOCH ₃	
10	iso-butyl	Ph-	7-SO ₂ CH ₃	
11	iso-butyl	Ph-	7-SCH ₂ CH ₃	
12	iso-butyl	Ph-	7-NH ₂	
13	iso-butyl	Ph-	7-NHOH	
14	iso-butyl	Ph-	7-NHCH ₃	
15	iso-butyl	Ph-	7-N(CH ₃) ₂	
16	iso-butyl	Ph-	7-N ⁺ (CH ₃) ₃ , I ⁻	
17	iso-butyl	Ph-	7-NHC(=O)CH ₃	
18	iso-butyl	Ph-	7-N(CH ₂ CH ₃) ₂	
19	iso-butyl	Ph-	7-NMeCH ₂ CO ₂ H	
20	iso-butyl	Ph-	7-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻	
21	iso-butyl	Ph-	7-(N)-morpholine	
22	iso-butyl	Ph-	7-(N)-azetidine	
23	iso-butyl	Ph-	7-(N)-N-methylazetidinium, I ⁻	
24	iso-butyl	Ph-	7-(N)-pyrrolidine	
25	iso-butyl	Ph-	7-(N)-N-methyl-pyrrolidinium, I ⁻	
26	iso-butyl	Ph-	7-(N)-N-methyl-morpholinium, I ⁻	
27	iso-butyl	Ph-	7-(N)-N'-methylpiperazine	
28	iso-butyl	Ph-	7-(N)-N'-dimethylpiperazinium, I ⁻	
29	iso-butyl	Ph-	7-NH-CB2	

30	iso-butyl	Ph-	7-NHC(O)C ₆ H ₁₁
31	iso-butyl	Ph-	7-NHC(O)CH ₂ Br
32	iso-butyl	Ph-	7-NH-C(NH)NH ₂
33	iso-butyl	Ph-	7-(2)-chlorophene
34	iso-butyl	Ph-	8-methyl
35	iso-butyl	Ph-	8-ethyl
36	iso-butyl	Ph-	8-iso-propyl
37	iso-butyl	Ph-	8-tert-butyl
38	iso-butyl	Ph-	8-OH
39	iso-butyl	Ph-	8-OCH ₃
40	iso-butyl	Ph-	8-O(1so-propyl)
41	iso-butyl	Ph-	8-SCH ₃
42	iso-butyl	Ph-	8-SOCH ₃
43	iso-butyl	Ph-	8-SO ₂ CH ₃
44	iso-butyl	Ph-	8-SCH ₂ CH ₃
45	iso-butyl	Ph-	8-NH ₂
46	iso-butyl	Ph-	8-NHCH ₃
47	iso-butyl	Ph-	8-NHCH ₂ CH ₃
48	iso-butyl	Ph-	8-N(CH ₃) ₂
49	iso-butyl	Ph-	8-N ⁺ (CH ₃) ₃ , I ⁻
50	iso-butyl	Ph-	8-NHC(=O)CH ₃
51	iso-butyl	Ph-	8-N(CH ₂ CH ₃) ₂
52	iso-butyl	Ph-	8-NHCCH ₂ CO ₂ H
53	iso-butyl	Ph-	8-N ⁺ (Me)CH ₂ CO ₂ H, I ⁻
54	iso-butyl	Ph-	8-(N)-morpholine
55	iso-butyl	Ph-	8-(N)-azetidine
56	iso-butyl	Ph-	8-(N)-N-methylazetidine, I ⁻
57	iso-butyl	Ph-	8-(N)-pyrrolidine
58	iso-butyl	Ph-	8-(N)-N-methyl-pyrrolidine, I ⁻
59	iso-butyl	Ph-	8-(N)-N-methyl-morpholine, I ⁻
60	iso-butyl	Ph-	8-(N)-N-methylpiperazine
61	iso-butyl	Ph-	8-(N)-N'-dimethylpiperazine, I ⁻
62	iso-butyl	Ph-	8-NH-CBz
63	iso-butyl	Ph-	8-NHC(O)C ₆ H ₁₁
64	iso-butyl	Ph-	8-NHC(O)CH ₂ Br
65	iso-butyl	Ph-	8-NH-C(NH)NH ₂
66	iso-butyl	Ph-	8-(2)-chlorophene
67	iso-butyl	Ph-	9-methyl
68	iso-butyl	Ph-	9-ethyl
69	iso-butyl	Ph-	9-iso-propyl
70	iso-butyl	Ph-	9-tert-butyl
71	iso-butyl	Ph-	9-OH

53

72	iso-butyl	Ph-	9-OCH ₃
73	iso-butyl	Ph-	9-O(1so-propyl)
74	iso-butyl	Ph-	9-SCH ₃
75	iso-butyl	Ph-	9-SOCH ₃
76	iso-butyl	Ph-	9-SO ₂ CH ₃
77	iso-butyl	Ph-	9-SCH ₂ CH ₃
78	iso-butyl	Ph-	9-NH ₂
79	iso-butyl	Ph-	9-NHCH ₃
80	iso-butyl	Ph-	9-NHCH ₂ CH ₃
81	iso-butyl	Ph-	9-N(CH ₃) ₂
82	iso-butyl	Ph-	9-N ⁺ (CH ₃) ₃ , I ⁻
83	iso-butyl	Ph-	9-NHC(=O)CH ₃
84	iso-butyl	Ph-	9-N(CH ₂ CH ₃) ₂
85	iso-butyl	Ph-	9-NHCCH ₂ CO ₂ H
86	iso-butyl	Ph-	9-N ⁺ (Me)CH ₂ CO ₂ H, I ⁻
87	iso-butyl	Ph-	9-(N)-morpholine
88	iso-butyl	Ph-	9-(N)-azetidine
89	iso-butyl	Ph-	9-(N)-N-methylazetidine, I ⁻
90	iso-butyl	Ph-	9-(N)-pyrrolidine
91	iso-butyl	Ph-	9-(N)-N-methyl-pyrrolidine, I ⁻
92	iso-butyl	Ph-	9-(N)-N-methyl-morpholine, I ⁻
93	iso-butyl	Ph-	9-(N)-N-methylpiperazine
94	iso-butyl	Ph-	9-(N)-N'-dimethylpiperazine, I ⁻
95	iso-butyl	Ph-	9-NH-CBz
96	iso-butyl	Ph-	9-NHC(O)C ₆ H ₁₁
97	iso-butyl	Ph-	9-NHC(O)CH ₂ Br
98	iso-butyl	Ph-	9-NH-C(NH)NH ₂
99	iso-butyl	Ph-	9-(2)-chlorophene
100	iso-butyl	Ph-	7-OCH ₃ , 8-OCH ₃
101	iso-butyl	Ph-	7-SCH ₃ , 8-OCH ₃
102	iso-butyl	Ph-	7-SCH ₃ , 8-SCH ₃
103	iso-butyl	Ph-	6-OCH ₃ , 7-OCH ₃ , 8-OCH ₃

Pract. F101.007	Qp#	R ¹ -R ²	R ⁵	(R ³) _q
01	01	1so-pentyl	Ph-	7-methyl
02	02	1so-pentyl	Ph-	7-ethyl
03	03	1so-pentyl	Ph-	7-iso-propyl
04	04	1so-pentyl	Ph-	7-tert-butyl
05	05	1so-pentyl	Ph-	7-OH
06	06	1so-pentyl	Ph-	7-OCH ₃
07	07	1so-pentyl	Ph-	7-O(1so-propyl)

54

08	Iso-pentyl	Ph-	7-SCH ₃	50	Iso-pentyl	Ph-	8-NHC(=O)CH ₃
09	Iso-pentyl	Ph-	7-SOCH ₃	51	Iso-pentyl	Ph-	8-N(CH ₂ CH ₃) ₂
10	Iso-pentyl	Ph-	7-SO ₂ CH ₃	52	Iso-pentyl	Ph-	8-NMeCH ₂ CO ₂ H
11	Iso-pentyl	Ph-	7-SCH ₂ CH ₃	53	Iso-pentyl	Ph-	8-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
12	Iso-pentyl	Ph-	7-NH ₂	54	Iso-pentyl	Ph-	8-(N)-morpholine
13	Iso-pentyl	Ph-	7-NHOH	55	Iso-pentyl	Ph-	8-(N)-azetidine
14	Iso-pentyl	Ph-	7-NHCH ₃	56	Iso-pentyl	Ph-	8-(N)-N-methylazetidinyl, I ⁻
15	Iso-pentyl	Ph-	7-N(CH ₃) ₂	57	Iso-pentyl	Ph-	8-(N)-pyrrolidine
16	Iso-pentyl	Ph-	7-N ⁺ (CH ₃) ₃ , I ⁻	58	Iso-pentyl	Ph-	8-(N)-N-methyl-pyrrolidinyl, I ⁻
17	Iso-pentyl	Ph-	7-NHC(=O)CH ₃	59	Iso-pentyl	Ph-	8-(N)-N-methyl-morpholinyl, I ⁻
18	Iso-pentyl	Ph-	7-N(CH ₂ CH ₃) ₂	60	Iso-pentyl	Ph-	8-(N)-N'-methylpiperazine
19	Iso-pentyl	Ph-	7-NMeCH ₂ CO ₂ H	61	Iso-pentyl	Ph-	8-(N)-N'-dimethylpiperazinyl, I ⁻
20	Iso-pentyl	Ph-	7-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻	62	Iso-pentyl	Ph-	8-NH-CBZ
21	Iso-pentyl	Ph-	7-(N)-morpholine	63	Iso-pentyl	Ph-	8-NHC(O)C ₅ H ₁₁
22	Iso-pentyl	Ph-	7-(N)-azetidine	64	Iso-pentyl	Ph-	8-NHC(O)CH ₂ Br
23	Iso-pentyl	Ph-	7-(N)-N-methylazetidinyl, I ⁻	65	Iso-pentyl	Ph-	8-NH-C(NH)NH ₂
24	Iso-pentyl	Ph-	7-(N)-pyrrolidine	66	Iso-pentyl	Ph-	8-(2)-thiophene
25	Iso-pentyl	Ph-	7-(N)-N-methyl-pyrrolidinyl, I ⁻	67	Iso-pentyl	Ph-	9-methyl
26	Iso-pentyl	Ph-	7-(N)-N-methyl-morpholinyl, I ⁻	68	Iso-pentyl	Ph-	9-ethyl
27	Iso-pentyl	Ph-	7-(N)-N'-methylpiperazine	69	Iso-pentyl	Ph-	9-iso-propyl
28	Iso-pentyl	Ph-	7-(N)-N'-dimethylpiperazinyl, I ⁻	70	Iso-pentyl	Ph-	9-tert-butyl
29	Iso-pentyl	Ph-	7-NH-CBZ	71	Iso-pentyl	Ph-	9-OH
30	Iso-pentyl	Ph-	7-NHC(O)C ₅ H ₁₁	72	Iso-pentyl	Ph-	9-OCH ₃
31	Iso-pentyl	Ph-	7-NHC(O)CH ₂ Br	73	Iso-pentyl	Ph-	9-O(1-iso-propyl)
32	Iso-pentyl	Ph-	7-NH-C(NH)NH ₂	74	Iso-pentyl	Ph-	9-SCH ₃
33	Iso-pentyl	Ph-	7-(2)-thiophene	75	Iso-pentyl	Ph-	9-SOCH ₃
34	Iso-pentyl	Ph-	8-methyl	76	Iso-pentyl	Ph-	9-SO ₂ CH ₃
35	Iso-pentyl	Ph-	8-ethyl	77	Iso-pentyl	Ph-	9-SCH ₂ CH ₃
36	Iso-pentyl	Ph-	8-iso-propyl	78	Iso-pentyl	Ph-	9-NH ₂
37	Iso-pentyl	Ph-	8-tert-butyl	79	Iso-pentyl	Ph-	9-NHOH
38	Iso-pentyl	Ph-	8-OH	80	Iso-pentyl	Ph-	9-NHCH ₃
39	Iso-pentyl	Ph-	8-OCH ₃	81	Iso-pentyl	Ph-	9-N(CH ₃) ₂
40	Iso-pentyl	Ph-	8-O(1-iso-propyl)	82	Iso-pentyl	Ph-	9-N ⁺ (CH ₃) ₃ , I ⁻
41	Iso-pentyl	Ph-	8-SCH ₃	83	Iso-pentyl	Ph-	9-NHC(=O)CH ₃
42	Iso-pentyl	Ph-	8-SOCH ₃	84	Iso-pentyl	Ph-	9-N(CH ₂ CH ₃) ₂
43	Iso-pentyl	Ph-	8-SO ₂ CH ₃	85	Iso-pentyl	Ph-	9-NMeCH ₂ CO ₂ H
44	Iso-pentyl	Ph-	8-SCH ₂ CH ₃	86	Iso-pentyl	Ph-	9-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
45	Iso-pentyl	Ph-	8-NH ₂	87	Iso-pentyl	Ph-	9-(N)-morpholine
46	Iso-pentyl	Ph-	8-NHOH	88	Iso-pentyl	Ph-	9-(N)-azetidine
47	Iso-pentyl	Ph-	8-NHCH ₃	89	Iso-pentyl	Ph-	9-(N)-N-methylazetidinyl, I ⁻
48	Iso-pentyl	Ph-	8-N(CH ₃) ₂	90	Iso-pentyl	Ph-	9-(N)-pyrrolidine
49	Iso-pentyl	Ph-	8-N ⁺ (CH ₃) ₃ , I ⁻	91	Iso-pentyl	Ph-	9-(N)-N-methyl-pyrrolidinyl, I ⁻
				92	Iso-pentyl	Ph-	9-(N)-N-methyl-morpholinyl, I ⁻

Prefix	Cpd#	R ¹ =R ²	R ³	(R ³) _q
<u>(III, xxx, xxy)</u>				
F101.008	01	CH ₂ C(=O)C ₂ H ₅	Ph-	7-methyl
	02	CH ₂ C(=O)C ₂ H ₅	Ph-	7-ethyl
	03	CH ₂ C(=O)C ₂ H ₅	Ph-	7-iso-propyl
	04	CH ₂ C(=O)C ₂ H ₅	Ph-	7-tert-butyl
	05	CH ₂ C(=O)C ₂ H ₅	Ph-	7-OR
	06	CH ₂ C(=O)C ₂ H ₅	Ph-	7-OCH ₃
	07	CH ₂ C(=O)C ₂ H ₅	Ph-	7-O(1-iso-propyl)
	08	CH ₂ C(=O)C ₂ H ₅	Ph-	7-SCH ₃
	09	CH ₂ C(=O)C ₂ H ₅	Ph-	7-SOCH ₃
	10	CH ₂ C(=O)C ₂ H ₅	Ph-	7-SO ₂ CH ₃
	11	CH ₂ C(=O)C ₂ H ₅	Ph-	7-SCH ₂ CH ₃
	12	CH ₂ C(=O)C ₂ H ₅	Ph-	7-NH ₂
	13	CH ₂ C(=O)C ₂ H ₅	Ph-	7-NHOH
	14	CH ₂ C(=O)C ₂ H ₅	Ph-	7-NHCH ₃
	15	CH ₂ C(=O)C ₂ H ₅	Ph-	7-N(CH ₃) ₂
	16	CH ₂ C(=O)C ₂ H ₅	Ph-	7-N ⁺ (CH ₃) ₃ , I ⁻
	17	CH ₂ C(=O)C ₂ H ₅	Ph-	7-NHC(=O)CH ₃
	18	CH ₂ C(=O)C ₂ H ₅	Ph-	7-N(CH ₂ CH ₃) ₂
	19	CH ₂ C(=O)C ₂ H ₅	Ph-	7-NHCCH ₂ CO ₂ H
	20	CH ₂ C(=O)C ₂ H ₅	Ph-	7-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
	21	CH ₂ C(=O)C ₂ H ₅	Ph-	7-(N)-morpholine
	22	CH ₂ C(=O)C ₂ H ₅	Ph-	7-(N)-azetidine
	23	CH ₂ C(=O)C ₂ H ₅	Ph-	7-(N)-N-methylazetidinium, I ⁺
	24	CH ₂ C(=O)C ₂ H ₅	Ph-	7-(N)-pyrrolidine
	25	CH ₂ C(=O)C ₂ H ₅	Ph-	7-(N)-N-methyl-pyrrolidinium, I ⁺
	26	CH ₂ C(=O)C ₂ H ₅	Ph-	7-(N)-N-methyl-morpholinium, I ⁺
	27	CH ₂ C(=O)C ₂ H ₅	Ph-	7-(N)-N'-methylpiperazine
	28	CH ₂ C(=O)C ₂ H ₅	Ph-	7-(N)-N'-dimethylpiperazinium, I ⁺
	29	CH ₂ C(=O)C ₂ H ₅	Ph-	7-NH-CBZ
	30	CH ₂ C(=O)C ₂ H ₅	Ph-	7-NHC(O)C ₅ H ₁₁
	31	CH ₂ C(=O)C ₂ H ₅	Ph-	7-NHC(O)CH ₂ Br
	32	CH ₂ C(=O)C ₂ H ₅	Ph-	7-NH-C(NH) ₂ NH ₂
	33	CH ₂ C(=O)C ₂ H ₅	Ph-	7-(2)-chlorophene
	34	CH ₂ C(=O)C ₂ H ₅	Ph-	8-methyl
	35	CH ₂ C(=O)C ₂ H ₅	Ph-	8-ethyl
	36	CH ₂ C(=O)C ₂ H ₅	Ph-	8-iso-propyl
	37	CH ₂ C(=O)C ₂ H ₅	Ph-	8-tert-butyl
	38	CH ₂ C(=O)C ₂ H ₅	Ph-	8-OH
	39	CH ₂ C(=O)C ₂ H ₅	Ph-	8-OCH ₃
	40	CH ₂ C(=O)C ₂ H ₅	Ph-	8-O(1-iso-propyl)
	41	CH ₂ C(=O)C ₂ H ₅	Ph-	8-SCH ₃
	42	CH ₂ C(=O)C ₂ H ₅	Ph-	8-SOCH ₃
	43	CH ₂ C(=O)C ₂ H ₅	Ph-	8-SO ₂ CH ₃
	44	CH ₂ C(=O)C ₂ H ₅	Ph-	8-SCH ₂ CH ₃
	45	CH ₂ C(=O)C ₂ H ₅	Ph-	8-NH ₂
	46	CH ₂ C(=O)C ₂ H ₅	Ph-	8-NHOH
	47	CH ₂ C(=O)C ₂ H ₅	Ph-	8-NHCH ₃
	48	CH ₂ C(=O)C ₂ H ₅	Ph-	8-N(CH ₃) ₂
	49	CH ₂ C(=O)C ₂ H ₅	Ph-	8-N ⁺ (CH ₃) ₃ , I ⁺
	50	CH ₂ C(=O)C ₂ H ₅	Ph-	8-NHC(=O)CH ₃
	51	CH ₂ C(=O)C ₂ H ₅	Ph-	8-N(CH ₂ CH ₃) ₂
	52	CH ₂ C(=O)C ₂ H ₅	Ph-	8-NHCCH ₂ CO ₂ H
	53	CH ₂ C(=O)C ₂ H ₅	Ph-	8-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁺
	54	CH ₂ C(=O)C ₂ H ₅	Ph-	8-(N)-morpholine
	55	CH ₂ C(=O)C ₂ H ₅	Ph-	8-(N)-azetidine
	56	CH ₂ C(=O)C ₂ H ₅	Ph-	8-(N)-N-methylazetidinium, I ⁺
	57	CH ₂ C(=O)C ₂ H ₅	Ph-	8-(N)-pyrrolidine
	58	CH ₂ C(=O)C ₂ H ₅	Ph-	8-(N)-N-methyl-pyrrolidinium, I ⁺
	59	CH ₂ C(=O)C ₂ H ₅	Ph-	8-(N)-N-methyl-morpholinium, I ⁺
	60	CH ₂ C(=O)C ₂ H ₅	Ph-	8-(N)-N'-methylpiperazine
	61	CH ₂ C(=O)C ₂ H ₅	Ph-	8-(N)-N'-dimethylpiperazinium, I ⁺
	62	CH ₂ C(=O)C ₂ H ₅	Ph-	8-NH-CBZ
	63	CH ₂ C(=O)C ₂ H ₅	Ph-	8-NHC(O)C ₅ H ₁₁
	64	CH ₂ C(=O)C ₂ H ₅	Ph-	8-NHC(O)CH ₂ Br
	65	CH ₂ C(=O)C ₂ H ₅	Ph-	8-NH-C(NH) ₂ NH ₂
	66	CH ₂ C(=O)C ₂ H ₅	Ph-	8-(2)-chlorophene
	67	CH ₂ C(=O)C ₂ H ₅	Ph-	9-methyl
	68	CH ₂ C(=O)C ₂ H ₅	Ph-	9-ethyl
	69	CH ₂ C(=O)C ₂ H ₅	Ph-	9-iso-propyl

Pat. No.	Cpd No.	R ¹ -m ²	R ⁵	(R ⁵) _q
70	100	CH ₂ C(=O)C ₂ H ₅	Ph-	9-tert-butyl
71	101	CH ₂ C(=O)C ₂ H ₅	Ph-	9-OH
72	102	CH ₂ C(=O)C ₂ H ₅	Ph-	9-OCH ₃
73	103	CH ₂ C(=O)C ₂ H ₅	Ph-	9-O(iso-propyl)
74		CH ₂ C(=O)C ₂ H ₅	Ph-	9-SCH ₃
75		CH ₂ C(=O)C ₂ H ₅	Ph-	9-SOCH ₃
76		CH ₂ C(=O)C ₂ H ₅	Ph-	9-SO ₂ CH ₃
77		CH ₂ C(=O)C ₂ H ₅	Ph-	9-SCH ₂ CH ₃
78		CH ₂ C(=O)C ₂ H ₅	Ph-	9-NH ₂
79		CH ₂ C(=O)C ₂ H ₅	Ph-	9-NHOH
80		CH ₂ C(=O)C ₂ H ₅	Ph-	9-NHCH ₃
81		CH ₂ C(=O)C ₂ H ₅	Ph-	9-N(CH ₃) ₂
82		CH ₂ C(=O)C ₂ H ₅	Ph-	9-N ⁺ (CH ₃) ₃ , I ⁻
83		CH ₂ C(=O)C ₂ H ₅	Ph-	9-NHC(=O)CH ₃
84		CH ₂ C(=O)C ₂ H ₅	Ph-	9-N(CH ₂ CH ₃) ₂
85		CH ₂ C(=O)C ₂ H ₅	Ph-	9-NMeCH ₂ CO ₂ H
86		CH ₂ C(=O)C ₂ H ₅	Ph-	9-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
87		CH ₂ C(=O)C ₂ H ₅	Ph-	9-(N)-morpholine
88		CH ₂ C(=O)C ₂ H ₅	Ph-	9-(N)-acetidine
89		CH ₂ C(=O)C ₂ H ₅	Ph-	9-[N]-N-methylacetidinidinium, I ⁻
90		CH ₂ C(=O)C ₂ H ₅	Ph-	9-(N)-pyrrolidine
91		CH ₂ C(=O)C ₂ H ₅	Ph-	9-(N)-N-methyl-pyrrolidinidinium, I ⁻
92		CH ₂ C(=O)C ₂ H ₅	Ph-	9-(N)-N-methyl-morpholinidinium, I ⁻
93		CH ₂ C(=O)C ₂ H ₅	Ph-	9-(N)-N'-methylpiperazine
94		CH ₂ C(=O)C ₂ H ₅	Ph-	9-(N)-N'-dimethylpiperazinidinium, I ⁻
95		CH ₂ C(=O)C ₂ H ₅	Ph-	9-NH-CBZ
96		CH ₂ C(=O)C ₂ H ₅	Ph-	9-NHC(=O)C ₅ H ₁₁
97		CH ₂ C(=O)C ₂ H ₅	Ph-	9-NHC(=O)CH ₂ Br
98		CH ₂ C(=O)C ₂ H ₅	Ph-	9-NH-C(NH)NH ₂
99		CH ₂ C(=O)C ₂ H ₅	Ph-	9-(2)-thiophene
100		CH ₂ C(=O)C ₂ H ₅	Ph-	7-OCH ₃ , 8-OCH ₃
101		CH ₂ C(=O)C ₂ H ₅	Ph-	7-SCH ₃ , 8-OCH ₃
102		CH ₂ C(=O)C ₂ H ₅	Ph-	7-SCH ₃ , 8-SCH ₃
103		CH ₂ C(=O)C ₂ H ₅	Ph-	6-OCH ₃ , 7-OCH ₃ , 8-OCH ₃

05	CH ₃ OC ₂ H ₅	Ph-	7-OH
06	CH ₃ OC ₂ H ₅	Ph-	7-OCH ₃
07	CH ₃ OC ₂ H ₅	Ph-	7-O(iso-propyl)
08	CH ₃ OC ₂ H ₅	Ph-	7-SCH ₃
09	CH ₃ OC ₂ H ₅	Ph-	7-SOCH ₃
10	CH ₃ OC ₂ H ₅	Ph-	7-SO ₂ CH ₃
11	CH ₃ OC ₂ H ₅	Ph-	7-SCH ₂ CH ₃
12	CH ₃ OC ₂ H ₅	Ph-	7-NH ₂
13	CH ₃ OC ₂ H ₅	Ph-	7-NHOH
14	CH ₃ OC ₂ H ₅	Ph-	7-NHCH ₃
15	CH ₃ OC ₂ H ₅	Ph-	7-N(CH ₃) ₂
16	CH ₃ OC ₂ H ₅	Ph-	7-N ⁺ (CH ₃) ₃ , I ⁻
17	CH ₃ OC ₂ H ₅	Ph-	7-NHC(=O)CH ₃
18	CH ₃ OC ₂ H ₅	Ph-	7-N(CH ₂ CH ₃) ₂
19	CH ₃ OC ₂ H ₅	Ph-	7-NMeCH ₂ CO ₂ H
20	CH ₃ OC ₂ H ₅	Ph-	7-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
21	CH ₃ OC ₂ H ₅	Ph-	7-(N)-morpholine
22	CH ₃ OC ₂ H ₅	Ph-	7-(N)-acetidine
23	CH ₃ OC ₂ H ₅	Ph-	7-(N)-N-methylacetidinidinium, I ⁻
24	CH ₃ OC ₂ H ₅	Ph-	7-(N)-pyrrolidine
25	CH ₃ OC ₂ H ₅	Ph-	7-(N)-N-methyl-pyrrolidinidinium, I ⁻
26	CH ₃ OC ₂ H ₅	Ph-	7-(N)-N-methyl-morpholinidinium, I ⁻
27	CH ₃ OC ₂ H ₅	Ph-	7-(N)-N'-methylpiperazine
28	CH ₃ OC ₂ H ₅	Ph-	7-(N)-N'-dimethylpiperazinidinium, I ⁻
29	CH ₃ OC ₂ H ₅	Ph-	7-NH-CBZ
30	CH ₃ OC ₂ H ₅	Ph-	7-NHC(=O)C ₅ H ₁₁
31	CH ₃ OC ₂ H ₅	Ph-	7-NHC(=O)CH ₂ Br
32	CH ₃ OC ₂ H ₅	Ph-	7-NH-C(NH)NH ₂
33	CH ₃ OC ₂ H ₅	Ph-	7-(2)-thiophene
34	CH ₃ OC ₂ H ₅	Ph-	8-methyl
35	CH ₃ OC ₂ H ₅	Ph-	8-ethyl
36	CH ₃ OC ₂ H ₅	Ph-	8-iso-propyl
37	CH ₃ OC ₂ H ₅	Ph-	8-tert-butyl
38	CH ₃ OC ₂ H ₅	Ph-	8-OH
39	CH ₃ OC ₂ H ₅	Ph-	8-OCH ₃
40	CH ₃ OC ₂ H ₅	Ph-	8-O(iso-propyl)
41	CH ₃ OC ₂ H ₅	Ph-	8-SCH ₃
42	CH ₃ OC ₂ H ₅	Ph-	8-SOCH ₃
43	CH ₃ OC ₂ H ₅	Ph-	8-SO ₂ CH ₃
44	CH ₃ OC ₂ H ₅	Ph-	8-SCH ₂ CH ₃
45	CH ₃ OC ₂ H ₅	Ph-	8-NH ₂
46	CH ₃ OC ₂ H ₅	Ph-	8-NHOH

47	CH ₃ OC ₂ H ₅	Ph- 8-NHCH ₃	Ph- 8-NHCH ₃
48	CH ₃ OC ₂ H ₅	Ph- 8-N(CH ₃) ₂	Ph- 8-N(CH ₃) ₂
49	CH ₃ OC ₂ H ₅	Ph- 8-N ⁺ (CH ₃) ₃ , I ⁻	Ph- 8-N ⁺ (CH ₃) ₃ , I ⁻
50	CH ₃ OC ₂ H ₅	Ph- 8-NHC(-O)CH ₃	Ph- 8-NHC(-O)CH ₃
51	CH ₃ OC ₂ H ₅	Ph- 8-N(CH ₂ CH ₃) ₂	Ph- 8-N(CH ₂ CH ₃) ₂
52	CH ₃ OC ₂ H ₅	Ph- 8-NHCCH ₂ CO ₂ H	Ph- 8-NHCCH ₂ CO ₂ H
53	CH ₃ OC ₂ H ₅	Ph- 8-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻	Ph- 8-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
54	CH ₃ OC ₂ H ₅	Ph- 8-(N)-morpholine	Ph- 8-(N)-morpholine
55	CH ₃ OC ₂ H ₅	Ph- 8-(N)-azetidine	Ph- 8-(N)-azetidine
56	CH ₃ OC ₂ H ₅	Ph- 8-(N)-N-methylazetidinylum, I ⁻	Ph- 8-(N)-N-methylazetidinylum, I ⁻
57	CH ₃ OC ₂ H ₅	Ph- 8-(N)-pyrrolidine	Ph- 8-(N)-pyrrolidine
58	CH ₃ OC ₂ H ₅	Ph- 8-(N)-N-methyl-pyrrolidinylum, I ⁻	Ph- 8-(N)-N-methyl-pyrrolidinylum, I ⁻
59	CH ₃ OC ₂ H ₅	Ph- 8-(N)-N-methyl-morpholinylum, I ⁻	Ph- 8-(N)-N-methyl-morpholinylum, I ⁻
60	CH ₃ OC ₂ H ₅	Ph- 8-(N)-N'-methylpiperazine	Ph- 8-(N)-N'-methylpiperazine
61	CH ₃ OC ₂ H ₅	Ph- 8-(N)-N'-dimethylpiperazinylum, I ⁻	Ph- 8-(N)-N'-dimethylpiperazinylum, I ⁻
62	CH ₃ OC ₂ H ₅	Ph- 8-NH-CBz	Ph- 8-NH-CBz
63	CH ₃ OC ₂ H ₅	Ph- 8-NHC(O)C ₆ H ₁₁	Ph- 8-NHC(O)C ₆ H ₁₁
64	CH ₃ OC ₂ H ₅	Ph- 8-NHC(O)CH ₂ Br	Ph- 8-NHC(O)CH ₂ Br
65	CH ₃ OC ₂ H ₅	Ph- 8-NH-C(NH)NH ₂	Ph- 8-NH-C(NH)NH ₂
66	CH ₃ OC ₂ H ₅	Ph- 8-(2)-chlorophene	Ph- 8-(2)-chlorophene
67	CH ₃ OC ₂ H ₅	Ph- 9-methyl	Ph- 9-methyl
68	CH ₃ OC ₂ H ₅	Ph- 9-ethyl	Ph- 9-ethyl
69	CH ₃ OC ₂ H ₅	Ph- 9-iso-propyl	Ph- 9-iso-propyl
70	CH ₃ OC ₂ H ₅	Ph- 9-tert-butyl	Ph- 9-tert-butyl
71	CH ₃ OC ₂ H ₅	Ph- 9-OH	Ph- 9-OH
72	CH ₃ OC ₂ H ₅	Ph- 9-OCH ₃	Ph- 9-OCH ₃
73	CH ₃ OC ₂ H ₅	Ph- 9-O(1-iso-propyl)	Ph- 9-O(1-iso-propyl)
74	CH ₃ OC ₂ H ₅	Ph- 9-SCH ₃	Ph- 9-SCH ₃
75	CH ₃ OC ₂ H ₅	Ph- 9-SOCH ₃	Ph- 9-SOCH ₃
76	CH ₃ OC ₂ H ₅	Ph- 9-SO ₂ CH ₃	Ph- 9-SO ₂ CH ₃
77	CH ₃ OC ₂ H ₅	Ph- 9-SCH ₂ CH ₃	Ph- 9-SCH ₂ CH ₃
78	CH ₃ OC ₂ H ₅	Ph- 9-NH ₂	Ph- 9-NH ₂
79	CH ₃ OC ₂ H ₅	Ph- 9-NHOH	Ph- 9-NHOH
80	CH ₃ OC ₂ H ₅	Ph- 9-NHCH ₃	Ph- 9-NHCH ₃
81	CH ₃ OC ₂ H ₅	Ph- 9-N(CH ₃) ₂	Ph- 9-N(CH ₃) ₂
82	CH ₃ OC ₂ H ₅	Ph- 9-N ⁺ (CH ₃) ₃ , I ⁻	Ph- 9-N ⁺ (CH ₃) ₃ , I ⁻
83	CH ₃ OC ₂ H ₅	Ph- 9-NHC(-O)CH ₃	Ph- 9-NHC(-O)CH ₃
84	CH ₃ OC ₂ H ₅	Ph- 9-N(CH ₂ CH ₃) ₂	Ph- 9-N(CH ₂ CH ₃) ₂
85	CH ₃ OC ₂ H ₅	Ph- 9-NMeCH ₂ CO ₂ H	Ph- 9-NMeCH ₂ CO ₂ H
86	CH ₃ OC ₂ H ₅	Ph- 9-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻	Ph- 9-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
87	CH ₃ OC ₂ H ₅	Ph- 9-(N)-morpholine	Ph- 9-(N)-morpholine

61

Pretilt	Cpd#	R ¹ , R ²	R ³	R ⁴
(FTT, X-ray, VFT)				
F101.010	01	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-methyl
	02	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-ethyl
	03	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-iso-propyl
	04	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-tert-butyl
	05	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-OH
	06	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-OCH ₃
	07	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-O(1-iso-propyl)
	08	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-SCH ₃
	09	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-SOCH ₃
	10	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-SO ₂ CH ₃
	11	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-SCH ₂ CH ₃
	12	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-NH ₂
	13	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-NHOH
	14	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-NHCH ₃
	15	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-N(CH ₃) ₂
	16	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-N ⁺ (CH ₃) ₃ , I ⁻
	17	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-NHC(-O)CH ₃
	18	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-N(CH ₂ CH ₃) ₂
	19	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-NMeCH ₂ CO ₂ H
	20	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
	21	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-(N)-morpholine
	22	CH ₃ CH(OH)C ₂ H ₅	Ph-	7-(N)-azetidine


62

23	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-(N)-N-methylazetidinium, I ⁻	65	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-NH-C(NH)NH ₂
24	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-(N)-pyrrolidine	66	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-(2)-thiophene
25	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-(N)-N-methyl-pyrrolidinium, I ⁻	67	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-methyl
26	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-(N)-N-methyl-morpholinium, I ⁻	68	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-ethyl
27	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-(N)-N'-methylpiperazine	69	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-iso-propyl
28	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-(N)-N'-dimethylpiperazinium, I ⁻	70	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-tert-butyl
29	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-NH-CBZ	71	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-OH
30	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-NHC(O)C ₅ H ₁₁	72	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-OCH ₃
31	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-NHC(O)CH ₂ Br	73	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-O(1-iso-propyl)
32	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-NH-C(NH)NH ₂	74	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-SCH ₃
33	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-(2)-thiophene	75	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-SOCH ₃
34	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-methyl	76	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-SO ₂ CH ₃
35	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-ethyl	77	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-SCH ₂ CH ₃
36	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-iso-propyl	78	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-NH ₂
37	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-tert-butyl	79	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-NHCH ₃
38	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-OH	80	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-NHCH ₃
39	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-OCH ₃	81	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-N(CH ₃) ₂
40	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-O(1-iso-propyl)	82	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-N ⁺ (CH ₃) ₃ , I ⁻
41	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-SCH ₃	83	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-NHC(=O)CH ₃
42	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-SOCH ₃	84	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-N(CH ₂ CH ₃) ₂
43	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-SO ₂ CH ₃	85	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-NMeCH ₂ CO ₂ H
44	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-SCH ₂ CH ₃	86	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
45	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-NH ₂	87	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-(N)-morpholine
46	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-NHCH ₃	88	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-(N)-azetidine
47	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-N(CH ₃) ₂	89	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-(N)-N-methylazetidinium, I ⁻
48	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-N ⁺ (CH ₃) ₃ , I ⁻	90	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-(N)-pyrrolidine
49	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-NHC(=O)CH ₃	91	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-(N)-N-methyl-pyrrolidinium, I ⁻
50	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-N(CH ₂ CH ₃) ₂	92	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-(N)-N-methyl-morpholinium, I ⁻
51	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-NMeCH ₂ CO ₂ H	93	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-(N)-N'-methylpiperazine
52	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻	94	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-(N)-N'-dimethylpiperazinium, I ⁻
53	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-(N)-morpholine	95	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-NH-CBZ
54	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-(N)-azetidine	96	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-NHC(O)C ₅ H ₁₁
55	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-(N)-N-methylazetidinium, I ⁻	97	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-NHC(O)CH ₂ Br
56	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-(N)-pyrrolidine	98	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-NH-C(NH)NH ₂
57	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-(N)-N-methyl-pyrrolidinium, I ⁻	99	CH ₂ CH(OH)C ₂ H ₅	Ph-	9-(2)-thiophene
58	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-(N)-N'-methyl-morpholinium, I ⁻	100	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-OCH ₃ , 8-OCH ₃
59	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-(N)-N'-methylpiperazine	101	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-SCH ₃ , 8-OCH ₃
60	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-(N)-N'-dimethylpiperazinium, I ⁻	102	CH ₂ CH(OH)C ₂ H ₅	Ph-	7-SCH ₃ , 8-SCH ₃
61	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-NH-CBZ	103	CH ₂ CH(OH)C ₂ H ₅	Ph-	6-OCH ₃ , 7-OCH ₃ , 8-OCH ₃
62	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-NHC(O)C ₅ H ₁₁				
63	CH ₂ CH(OH)C ₂ H ₅	Ph-	8-NHC(O)CH ₂ Br				
64	CH ₂ CH(OH)C ₂ H ₅	Ph-					


Prefix		Q ₁ #	R ₁ -n ²	R ₅	(R ₂) ^q
F101.011		(TFT, XXX, YYY)			
01	CH ₂ O-(4-picoline)	Ph-	7-methyl		
02	CH ₂ O-(4-picoline)	Ph-	7-ethyl		
03	CH ₂ O-(4-picoline)	Ph-	7-iso-propyl		
04	CH ₂ O-(4-picoline)	Ph-	7-tert-butyl		
05	CH ₂ O-(4-picoline)	Ph-	7-OH		
06	CH ₂ O-(4-picoline)	Ph-	7-OCH ₃		
07	CH ₂ O-(4-picoline)	Ph-	7-O(1-iso-propyl)		
08	CH ₂ O-(4-picoline)	Ph-	7-SCH ₃		
09	CH ₂ O-(4-picoline)	Ph-	7-SOCH ₃		
10	CH ₂ O-(4-picoline)	Ph-	7-SO ₂ CH ₃		
11	CH ₂ O-(4-picoline)	Ph-	7-SCH ₂ CH ₃		
12	CH ₂ O-(4-picoline)	Ph-	7-NH ₂		
13	CH ₂ O-(4-picoline)	Ph-	7-NHOH		
14	CH ₂ O-(4-picoline)	Ph-	7-NHCH ₃		
15	CH ₂ O-(4-picoline)	Ph-	7-N(CH ₃) ₂		
16	CH ₂ O-(4-picoline)	Ph-	7-N ⁺ (CH ₃) ₃ , I ⁻		
17	CH ₂ O-(4-picoline)	Ph-	7-NHC(=O)CH ₃		
18	CH ₂ O-(4-picoline)	Ph-	7-N(CH ₂ CH ₃) ₂		
19	CH ₂ O-(4-picoline)	Ph-	7-NMeCH ₂ CO ₂ H		
20	CH ₂ O-(4-picoline)	Ph-	7-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻		
21	CH ₂ O-(4-picoline)	Ph-	7-(N)-morpholine		
22	CH ₂ O-(4-picoline)	Ph-	7-(N)-azetidine		
23	CH ₂ O-(4-picoline)	Ph-	7-(N)-N-methylazetidinum, I ⁺		
24	CH ₂ O-(4-picoline)	Ph-	7-(N)-pyrrolidine		
25	CH ₂ O-(4-picoline)	Ph-	7-(N)-N-methyl-pyrrolidinum, I ⁺		
26	CH ₂ O-(4-picoline)	Ph-	7-(N)-N-methyl-morpholinum, I ⁺		
27	CH ₂ O-(4-picoline)	Ph-	7-(N)-N'-methylpiperazine		
28	CH ₂ O-(4-picoline)	Ph-	7-(N)-N'-dimethylpiperazinum, I ⁺		
29	CH ₂ O-(4-picoline)	Ph-	7-NH-CBZ		
30	CH ₂ O-(4-picoline)	Ph-	7-NHC(O)C ₅ H ₁₁		
31	CH ₂ O-(4-picoline)	Ph-	7-NHC(O)CH ₂ Br		
32	CH ₂ O-(4-picoline)	Ph-	7-NH-C(=NH)NH ₂		
33	CH ₂ O-(4-picoline)	Ph-	7-(2)-thiophene		
34	CH ₂ O-(4-picoline)	Ph-	8-methyl		
35	CH ₂ O-(4-picoline)	Ph-	8-ethyl		
36	CH ₂ O-(4-picoline)	Ph-	8-iso-propyl		
37	CH ₂ O-(4-picoline)	Ph-	8-tert-butyl		
38	CH ₂ O-(4-picoline)	Ph-	8-OH		
39	CH ₂ O-(4-picoline)	Ph-	8-OCH ₃		
40	CH ₂ O-(4-picoline)	Ph-	8-O(1-iso-propyl)		
41	CH ₂ O-(4-picoline)	Ph-	8-SCH ₃		
42	CH ₂ O-(4-picoline)	Ph-	8-SOCH ₃		
43	CH ₂ O-(4-picoline)	Ph-	8-SO ₂ CH ₃		
44	CH ₂ O-(4-picoline)	Ph-	8-SCH ₂ CH ₃		
45	CH ₂ O-(4-picoline)	Ph-	8-NH ₂		
46	CH ₂ O-(4-picoline)	Ph-	8-NHOH		
47	CH ₂ O-(4-picoline)	Ph-	8-NHCH ₃		
48	CH ₂ O-(4-picoline)	Ph-	8-N(CH ₃) ₂		
49	CH ₂ O-(4-picoline)	Ph-	8-N ⁺ (CH ₃) ₃ , I ⁻		
50	CH ₂ O-(4-picoline)	Ph-	8-NHC(=O)CH ₃		
51	CH ₂ O-(4-picoline)	Ph-	8-N(CH ₂ CH ₃) ₂		
52	CH ₂ O-(4-picoline)	Ph-	8-NMeCH ₂ CO ₂ H		
53	CH ₂ O-(4-picoline)	Ph-	8-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻		
54	CH ₂ O-(4-picoline)	Ph-	8-(N)-azetidine		
55	CH ₂ O-(4-picoline)	Ph-	8-(N)-N-methylazetidinum, I ⁺		
56	CH ₂ O-(4-picoline)	Ph-	8-(N)-pyrrolidine		
57	CH ₂ O-(4-picoline)	Ph-	8-(N)-N-methyl-pyrrolidinum, I ⁺		
58	CH ₂ O-(4-picoline)	Ph-	8-(N)-N'-methylpiperazine		
59	CH ₂ O-(4-picoline)	Ph-	8-(N)-N'-dimethylpiperazinum, I ⁺		
60	CH ₂ O-(4-picoline)	Ph-	8-(N)-N'-methylpiperazine		
61	CH ₂ O-(4-picoline)	Ph-	8-(N)-N'-dimethylpiperazinum, I ⁺		
62	CH ₂ O-(4-picoline)	Ph-	8-NH-CBZ		
63	CH ₂ O-(4-picoline)	Ph-	8-NHC(O)C ₅ H ₁₁		
64	CH ₂ O-(4-picoline)	Ph-	8-NHC(O)CH ₂ Br		
65	CH ₂ O-(4-picoline)	Ph-	8-NH-C(=NH)NH ₂		
66	CH ₂ O-(4-picoline)	Ph-	8-(2)-thiophene		
67	CH ₂ O-(4-picoline)	Ph-	9-methyl		
68	CH ₂ O-(4-picoline)	Ph-	9-ethyl		
69	CH ₂ O-(4-picoline)	Ph-	9-iso-propyl		
70	CH ₂ O-(4-picoline)	Ph-	9-tert-butyl		
71	CH ₂ O-(4-picoline)	Ph-	9-OH		
72	CH ₂ O-(4-picoline)	Ph-	9-OCH ₃		
73	CH ₂ O-(4-picoline)	Ph-	9-O(1-iso-propyl)		
74	CH ₂ O-(4-picoline)	Ph-	9-SCH ₃		
75	CH ₂ O-(4-picoline)	Ph-	9-SOCH ₃		
76	CH ₂ O-(4-picoline)	Ph-	9-SO ₂ CH ₃		
77	CH ₂ O-(4-picoline)	Ph-	9-SCH ₂ CH ₃		
78	CH ₂ O-(4-picoline)	Ph-	9-NH ₂		
79	CH ₂ O-(4-picoline)	Ph-	9-NHOH		
80	CH ₂ O-(4-picoline)	Ph-	9-NHCH ₃		
81	CH ₂ O-(4-picoline)	Ph-	9-N(CH ₃) ₂		




82	CH ₃ O-(4-picoline)	Ph-	9-N ⁺ (CH ₃) ₃ , I ⁻
83	CH ₃ O-(4-picoline)	Ph-	9-NHC(=O)CH ₃
84	CH ₃ O-(4-picoline)	Ph-	9-N(CH ₂ CH ₃) ₂
85	CH ₃ O-(4-picoline)	Ph-	9-NMeCH ₂ CO ₂ H
86	CH ₃ O-(4-picoline)	Ph-	9-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁻
87	CH ₃ O-(4-picoline)	Ph-	9-(N)-morpholine
88	CH ₃ O-(4-picoline)	Ph-	9-(N)-acetidine
89	CH ₃ O-(4-picoline)	Ph-	9-(N)-methylazetidine, I ⁻
90	CH ₃ O-(4-picoline)	Ph-	9-(N)-pyrrolidine
91	CH ₃ O-(4-picoline)	Ph-	9-(N)-N-methyl-pyrrolidinium, I ⁻
92	CH ₃ O-(4-picoline)	Ph-	9-(N)-N-methyl-morpholinium, I ⁻
93	CH ₃ O-(4-picoline)	Ph-	9-(N)-N'-methylpiperazine
94	CH ₃ O-(4-picoline)	Ph-	9-(N)-N'-dimethylpiperazinium, I ⁻
95	CH ₃ O-(4-picoline)	Ph-	9-NH-CBz
96	CH ₃ O-(4-picoline)	Ph-	9-NHC(O)C ₅ H ₁₁
97	CH ₃ O-(4-picoline)	Ph-	9-NHC(O)CH ₂ Br
98	CH ₃ O-(4-picoline)	Ph-	9-NH-C(NH) ₂ NH ₂
99	CH ₃ O-(4-picoline)	Ph-	9-(2)-thiophene
100	CH ₃ O-(4-picoline)	Ph-	7-OCH ₃ , 8-OCH ₃
101	CH ₃ O-(4-picoline)	Ph-	7-SCH ₃ , 8-OCH ₃
102	CH ₃ O-(4-picoline)	Ph-	7-SCH ₃ , 8-SCH ₃
103	CH ₃ O-(4-picoline)	Ph-	6-OCH ₃ , 7-OCH ₃ , 8-OCH ₃






Additional Structures of the Present Invention










Compound Number	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸
101	ethyl	n-butyl	OH	H		phenyl		
102	ethyl	n-butyl	OH	H		phenyl		
103	n-butyl	ethyl	OH	H		phenyl		
104	ethyl	n-butyl	OH	H		phenyl		
105	ethyl	n-butyl	OH	H		phenyl		
106	ethyl	n-butyl	OH	H		phenyl		
107	n-butyl	ethyl	OH	H		phenyl		
108	ethyl	n-butyl	OH	H		phenyl		
109	ethyl	n-butyl	OH	H		phenyl		
110	ethyl	n-butyl	OH	H		phenyl		
111	n-butyl	ethyl	OH	H		phenyl		



































112	ethyl	n-butyl	OH	H		H	7-amino
113	ethyl	n-butyl	OH	H	4-hydroxyphenyl	H	7-amino
114	ethyl	n-butyl	OH	H	4-methoxyphenyl	H	7-amino
115	n-butyl	ethyl	OH	H	4-methoxyphenyl	H	7-(O-benzylcarbamato)
116	ethyl	n-butyl	OH	H	4-methoxyphenyl	H	7-(O-benzylcarbamato)
117	n-butyl	ethyl	OH	H	phenyl	H	7-(O-benzylcarbamato)
118	ethyl	n-butyl	OH	H	phenyl	H	7-(O-benzylcarbamato)
119	ethyl	n-butyl	OH	H	phenyl	H	7-(O-tert-butylcarbamato)
120	n-butyl	ethyl	OH	H	phenyl	H	7-(O-benzylcarbamato)
121	ethyl	n-butyl	OH	H	phenyl	H	7-amino
122	n-butyl	ethyl	OH	H	phenyl	H	7-amino
123	ethyl	n-butyl	OH	H	phenyl	H	7-hexylamino
124	n-butyl	ethyl	OH	H	phenyl	H	7-(hexylamino)
125	ethyl	n-butyl	OH	H	phenyl	H	1
126	n-butyl	ethyl	OH	H	4-fluorophenyl	H	 at the 8-position
127	n-butyl	ethyl	OH	H	4-fluorophenyl	H	7-(O-benzylcarbamato)
128	ethyl	n-butyl	OH	H	4-fluorophenyl	H	7-amino
129	ethyl	n-butyl	OH	H	4-fluorophenyl	H	7-(O-benzylcarbamato)
131	ethyl	n-butyl	OH	H	4-fluorophenyl	H	7-amino
							 at the 7-position

132	ethyl	n-butyl	OH	H	phenyl	H	 at the 8-position
133	ethyl	n-butyl	OH	H	phenyl	H	8-(hexyloxy)
134	ethyl	n-butyl	OH	H	phenyl	H	 at the 8-position
135	ethyl	n-butyl	OH	H	phenyl	H	 at the 8-position
136	ethyl	n-butyl	OH	H	phenyl	H	8-hydroxy
137	n-butyl	ethyl	OH	H	phenyl	H	 at the 7-position
138	n-butyl	ethyl	OH	H	phenyl	H	8-acetoxy
139	n-butyl	ethyl	OH	H	phenyl	H	 at the 7-position
140							
141							
142	ethyl	n-butyl	H	OH	H	3-methoxy-phenyl	7-methylmercapto
143	ethyl	n-butyl	OH	H	3-methoxyphenyl	H	7-methylmercapto
144	ethyl	n-butyl	OH	H	4-fluorophenyl	H	7-(N-azetidinyl)
262	ethyl	n-butyl	OH	H	3-methoxyphenyl	H	7-methoxy
263	ethyl	n-butyl	H	OH	H	3-methoxy-	7-methoxy






296	ethyl	n-butyl	OH	H	H		H	7-amino
1000	ethyl	n-butyl	OH	H	H		H	7-dimethylamino
1001	ethyl	n-butyl	OH	H	H		H	7-dimethylamino
1002	ethyl	n-butyl	OH	H	H		H	7-dimethylamino
1003	ethyl	n-butyl	OH	H	H		H	7-dimethylamino
1004	ethyl	n-butyl	OH	H	H		H	7-dimethylamino
1005	n-butyl	n-butyl	OH	H	H		H	7-dimethylamino

72

264	ethyl	n-butyl	OH	H	H		phenyl	7-methoxy
265	ethyl	n-butyl	OH	H	H		3-nitrophenyl	7-methoxy
266	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
267	ethyl	n-butyl	OH	H	H		3-methoxyphenyl	7-methoxy
268	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
269	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
270	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
271	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
272	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
273	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
274	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
275	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
276	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
277	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
278	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
279	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
280	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
281	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
282	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
283	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
284	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
285	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
286	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
287	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
288	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
289	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
290	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
291	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
292	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
293	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
294	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy
295	ethyl	n-butyl	OH	H	H		4-fluorophenyl	7-methoxy

72







73

1006	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1007	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1008	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1009	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1010	n-butyl	n-butyl	OH	H	3-fluoro-4-methoxyphenyl	H	7-dimethylamino
1011	n-butyl	n-butyl	OH	H	3-fluoro-4-(5-triethylammoniumpentloxy)phenyl, trifluoroacetate salt	H	7-dimethylamino
1012	n-butyl	n-butyl	OH	H	4-hydroxyphenyl	H	7-dimethylamino; 9-methoxy
1013	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1014	n-butyl	n-butyl	OH	H	4-methoxyphenyl	H	7-dimethylamino; 9-methoxy

WO 97/33882

PCT/US97/04076

74

1015	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1016	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1017	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1018	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1019	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1020	n-butyl	n-butyl	OH	H		H	7-dimethylamino

WO 97/33882

PCT/US97/04076

9t

1024	n-butyl	n-butyl	OH	H		7-dimethylamino
1025	n-butyl	n-butyl	OH	H		7-dimethylamino
1026	n-butyl	n-butyl	OH	H		7-dimethylamino
1027	n-butyl	n-butyl	OH	H		7-dimethylamino
1028	n-butyl	n-butyl	OH	H		7-dimethylamino

5t

1021	n-butyl	n-butyl	OH	H		7-dimethylamino
1022	n-butyl	n-butyl	OH	H		7-dimethylamino
1023	n-butyl	n-butyl	OH	H		7-dimethylamino

PCT/US97/04076

WO 97/33882






PCT/US97/04076

WO 97/33882








1029	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1030	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1031	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1032	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1033	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1034	n-butyl	n-butyl	OH	H		H	7-dimethylamino

1035	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1036	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1037	n-butyl	n-butyl	OH	H	4-hydroxyphenyl	H	7-dimethylamino
1038	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1039	n-butyl	n-butyl	OH	H	phenyl	H	7-dimethylamino
1040	n-butyl	n-butyl	OH	H		H	7-dimethylamino






???How does this differ from 73281?

1048	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1049	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1050	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1051	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1052	n-butyl	n-butyl	OH	H		H	7-dimethylamino

33






1041	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1042	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1043	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1044	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1045	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1046	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1047	n-butyl	n-butyl	OH	H		H	7-dimethylamino

34

1053	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1054	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1055	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1056	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1057	n-butyl	n-butyl	OH	H		H	7-dimethylamino








WO 97/33882







PCT/US97/04076





1058	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1059	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1060	ethyl	n-butyl	OH	H	3-fluoro-4-methoxyphenyl	H	7-methylamino
1061	n-butyl	n-butyl	OH	H		H	7-methylamino
1062	n-butyl	n-butyl	OH	H		H	7-methylamino
1063	n-butyl	n-butyl	OH	H		H	7-methylamino

WO 97/33882

PCT/US97/04076






1070	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1071	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1072	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1073	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1074	ethyl	n-butyl	OH	H		H	7-dimethylamino
1075	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1076	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1077	n-butyl	n-butyl	OH	H		H	7-dimethylamino

1064	n-butyl	n-butyl	OH	H		H	7-methylamino
1065	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1066	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1067	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1068	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1069	n-butyl	n-butyl	OH	H		H	7-dimethylamino

1078	ethyl	n-butyl	OH	H		H	7-dimethylamino
1079	ethyl	n-butyl	OH	H		H	7-dimethylamino
1080	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1081	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1082	n-butyl	n-butyl	OH	H		H	7-dimethylamino










WO 97/33882





PCT/US97/04076






1083	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1084	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1085	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1086	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1087	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1088	ethyl	n-butyl	OH	H		H	7-dimethylamino
1089	ethyl	n-butyl	OH	H		H	7-dimethylamino
1090	n-butyl	n-butyl	OH	H		H	7-dimethylamino






WO 97/33882





PCT/US97/04076





1095	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1096	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1097	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1098	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1099	ethyl	n-butyl	OH	H		H	7-dimethylamino
1100	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1101	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1102	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1103	n-butyl	n-butyl	OH	H		H	7-dimethylamino


1091	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1092	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1093	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1094	n-butyl	n-butyl	OH	H		H	7-dimethylamino




1104	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1105	n-butyl	n-butyl	OH	H	5-piperonyl	H	7-dimethylamino
1106	n-butyl	n-butyl	OH	H	3-hydroxyphenyl	H	7-dimethylamino
1107	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1108	n-butyl	n-butyl	OH	H	3-pyridyl	H	7-dimethylamino
1109	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1110	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1111	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1112	n-butyl	n-butyl	OH	H	4-pyridyl	H	7-dimethylamino

1113	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1114	n-butyl	n-butyl	OH	H	3-methoxyphenyl	H	7-methylamino
1115	n-butyl	n-butyl	OH	H	4-fluorophenyl	H	7-dimethylamino
1116	ethyl	n-butyl	OH	H	3-tolyl	H	7-dimethylamino
1117	ethyl	n-butyl	OH	H		H	7-dimethylamino
1118	ethyl	n-butyl	OH	H	3-fluoro-4-hydroxyphenyl	H	7-dimethylamino
1119	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1120	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1121	n-butyl	n-butyl	OH	H		H	7-dimethylamino

1135	n-butyl	n-butyl	OH	H	3,4-dimethoxyphenyl	7-dimethylamino
1136	n-butyl	n-butyl	OH	H		7-dimethylamino
1137	n-butyl	n-butyl	OH	H	4-fluorophenyl	9-(2,2-dimethylhydrazino)
1138	n-butyl	n-butyl	OH	H		7-dimethylamino
1139	n-butyl	n-butyl	OH	H	3,4-difluorophenyl	7-dimethylamino
1140	n-butyl	n-butyl	OH	H	3-methoxyphenyl	7-(2,2-dimethylhydrazino)
1141	n-butyl	n-butyl	OH	H	4-fluorophenyl	7-dimethylamino
1142	n-butyl	n-butyl	OH	H		7-dimethylamino
1143	n-butyl	n-butyl	H	OH	3-fluoro-4-methoxyphenyl	7-dimethylamino
1144	n-butyl	n-butyl	OH	H	5-pyridon	7-dimethylamino
1145	n-butyl	n-butyl	OH	H	4-methoxyphenyl	9-dimethylamino
1146	n-butyl	n-butyl	OH	H		7-dimethylamino
1147	n-butyl	n-butyl	OH	H	3-methoxyphenyl	7-diethylamino
1148	n-butyl	n-butyl	OH	H	4-fluorophenyl	7-dimethylsulfonyl, fluoroide salt
1149	n-butyl	n-butyl	OH	H	4-fluorophenyl	7-ethylamino
1150	n-butyl	n-butyl	OH	H	3-methoxyphenyl	7-ethylamino
1151	n-butyl	n-butyl	OH	H	3-fluoro-4-methoxyphenyl	7-dimethylamino
1152	n-butyl	n-butyl	OH	H	phenyl	7-(ethoxymethyl) methylamino
1153	n-butyl	n-butyl	OH	H	4-fluorophenyl	7-methylamino
1154	n-butyl	n-butyl	OH	H	3-methoxyphenyl	5-methoxy
1155	n-butyl	n-butyl	OH	H	4-fluorophenyl	7-methyl

1122	n-butyl	n-butyl	OH	H		7-dimethylamino
1123	n-butyl	n-butyl	OH	H	phenyl	7-dimethylamino
1125	n-butyl	n-butyl	OH	H	3-methoxyphenyl	7-dimethylamino
1126	n-butyl	n-butyl	OH	H	3-chloro-4-methoxyphenyl	7-dimethylamino
1127	n-butyl	n-butyl	OH	H		7-dimethylamino
1128	n-butyl	n-butyl	OH	H	3-fluoro-4-hydroxyphenyl	7-dimethylamino
1129	n-butyl	n-butyl	OH	H	4-fluorophenyl	9-dimethylamino
1130	n-butyl	n-butyl	OH	H	3-chloro-4-fluorophenyl	7-dimethylamino
1131	n-butyl	n-butyl	OH	H	4-methoxyphenyl	7-dimethylamino
1132	n-butyl	n-butyl	OH	H		7-dimethylamino
1133	n-butyl	n-butyl	OH	H	4-cyanomethylphenyl	7-dimethylamino
1134	ethyl	n-butyl	OH	H		7-dimethylamino

1156	n-butyl	n-butyl	OH	H	4-fluorophenyl	H	7-methylmercapto
1157	n-butyl	n-butyl	OH	H	4-fluorophenyl	H	7-fluoro;
1158	n-butyl	n-butyl	OH	H	4-pyridinyl, hydrochloride salt	H	9-dimethylamino
1159	n-butyl	ethyl	OH	H	phenyl	H	7-methoxy
1160	n-butyl	n-butyl	OH	H	4-fluorophenyl	H	7-dimethylamino
1161	n-butyl	n-butyl	OH	H	3,5-dichloro-4-methoxyphenyl	H	7-diethylamino
1162	n-butyl	n-butyl	OH	H	phenyl	H	7-dimethylamino
1163	n-butyl	n-butyl	OH	H	3-(dimethylamino)phenyl	H	7-dimethylamino
1164	n-butyl	n-butyl	OH	H	4-pyridinyl	H	7-methoxy
1165	n-butyl	n-butyl	OH	H	3-fluoro-4-methoxyphenyl	H	7-trimethylammonium iodide
1166	n-butyl	n-butyl	OH	H	3-hydroxyphenyl	H	7-trimethylammonium iodide
1167	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1168	n-butyl	n-butyl	OH	H	4-hydroxyphenyl	H	7-trimethylammonium iodide
1169	n-butyl	n-butyl	OH	H	phenyl	H	8-dimethylamino
1170	n-butyl	n-butyl	OH	H	3-methoxyphenyl	H	7-ethylpropylamino
1171	n-butyl	n-butyl	OH	H	4-(trifluoromethylsulfonyloxy)phenyl	H	7-dimethylamino
1172	n-butyl	n-butyl	OH	H	4-pyridinyl	H	7-methoxy
1173	n-butyl	n-butyl	OH	H	4-fluorophenyl	H	7-ethylpropylamino
1174	ethyl	n-butyl	OH	H	3-methoxyphenyl	H	7-phenyl
1175	ethyl	n-butyl	OH	H	3-methoxyphenyl	H	7-methylsulfonyl
1176	n-butyl	n-butyl	OH	H	4-fluorophenyl	H	9-fluoro
1177	n-butyl	n-butyl	OH	H	3-methoxyphenyl	H	7-butylmethylamino
1178	n-butyl	n-butyl	OH	H	3-(trifluoromethylsulfonyloxy)phenyl	H	7-dimethylamino
1179	n-butyl	n-butyl	OH	H	phenyl	H	8-methoxy
1180	n-butyl	n-butyl	OH	H	phenyl	H	7-trimethylammonium iodide
1181	n-butyl	n-butyl	OH	H	4-fluorophenyl	H	7-butylmethylamino
1182	n-butyl	n-butyl	OH	H	4-(dimethylamino)phenyl	H	7-methoxy
1183	n-butyl	n-butyl	OH	H	3-methoxyphenyl	H	7-fluoro
1184	n-butyl	n-butyl	OH	H	4-fluorophenyl	H	7-fluoro;
1185	n-butyl	n-butyl	OH	H	4-fluorophenyl	H	9-fluoro
1186	n-butyl	n-butyl	OH	H	phenyl	H	7-fluoro
1187	n-butyl	n-butyl	OH	H	4-fluorophenyl	H	7-fluoro;
1188	n-butyl	n-butyl	OH	H	4-methoxyphenyl	H	9-fluoro
1189	n-butyl	n-butyl	OH	H	3,4-difluorophenyl	H	7-methyl
1190	n-butyl	n-butyl	OH	H	2-bromophenyl	H	7-trimethylammonium iodide
1191	n-butyl	n-butyl	OH	H	4-(dimethylamino)phenyl	H	7-trimethylammonium iodide
1192	n-butyl	n-butyl	OH	H	3-(dimethylamino)phenyl	H	7-bromo
1193	n-butyl	n-butyl	OH	H	4-(2-(2-methylpropyl)phenyl)	H	7-hydroxy
							7-dimethylamino

1194	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1195	n-butyl	n-butyl	OH	H	4-methoxyphenyl	H	7-(4'-methylpiperazin-1-yl)
1196	n-butyl	n-butyl	OH	H		H	7-methoxy
1197	n-butyl	ethyl	R3 + R4 = oxo	R3 + R4 = oxo	phenyl	H	7-(N-methylformamido)
1198	n-butyl	n-butyl	OH	H	4-(pyridinyl-N-oxide)	H	7-methoxy
1199	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1200	n-butyl	n-butyl	H	OH	H	phenyl	7-dimethylamino
1201	n-butyl	n-butyl	OH	H	H	H	7-methyl

97

1234	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1235	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1236	n-butyl	n-butyl	OH	H	4-(bromomethyl)phenyl	H	7-dimethylamino
1237	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1238	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1239	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1240	n-butyl	n-butyl	OH	H	4-methoxy-3-methylphenyl	H	7-dimethylamino
1241	n-butyl	n-butyl	OH	H	3-(dimethylaminomethyl)phenyl	H	7-dimethylamino
1242	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1243	n-butyl	n-butyl	OH	H		H	7-dimethylamino

WO 97/33882

PCT/US97/04076

98

1244	n-butyl	n-butyl	OH	H	3-methoxyphenyl	H	7-(1'-methylhydrazido)
1245	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1246	n-butyl	n-butyl	OH	H	3-(bromomethyl)phenyl	H	7-dimethylamino
1247	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1248	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1249	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1250	n-butyl	n-butyl	OH	H	3-(dimethylamino)phenyl	H	7-dimethylamino
1251	n-butyl	n-butyl	OH	H	1-naphthyl	H	7-dimethylamino
1252	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1253	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1254	n-butyl	n-butyl	OH	H		H	7-dimethylamino

WO 97/33882

PCT/US97/04076

1270	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1271	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1272	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1273	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1274	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1275	n-butyl	n-butyl	OH	H		7-dimethylamino	H

001

1255	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1256	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1257	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1258	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1259	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1260	ethyl	n-butyl	OH	H		7-dimethylamino	phenyl
1261	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1262	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1263	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1264	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1265	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1266	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1267	ethyl	n-butyl	OH	H		7-carboxy, methyl ester	H
1268	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1269	n-butyl	n-butyl	OH	H		7-dimethylamino	H

bb







1286	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1287	n-butyl	ethyl	OH	H		H	7-dimethylamino
1288	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1289	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1290	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1291	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1292	n-butyl	n-butyl	OH	H		H	7-dimethylamino

102

1276	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1277	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1278	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1279	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1280	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1281	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1282	ethyl	n-butyl	OH	H		H	7-trimethylammonium iodide
1283	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1284	n-butyl	n-butyl	OH	H		H	9-ethylamino
1285	n-butyl	ethyl	OH	H		H	7-dimethylamino

101





103

1293	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1294	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1295	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1296	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1297	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1298	n-butyl	n-butyl	OH	H		H	7-dimethylamino

WO 97/33882







PCT/US97/04076

104











1299	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1300	n-butyl	ethyl	H	OH	H	phenyl	7-dimethylamino
1301	n-butyl	n-butyl	OH	H	3-methoxyphenyl	H	7-trimethylammonium iodide
1302	n-butyl	n-butyl	OH	H	3-hydroxyphenyl	H	9-hydroxy
1303	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1304	n-butyl	n-butyl	OH	H	3-methoxyphenyl	H	7-tert-butylamino
1305	n-butyl	n-butyl	OH	H	4-fluorophenyl	H	9-methylamino
1306	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1307	n-butyl	n-butyl	OH	H	4-methoxyphenyl	H	9-(4'-morpholino)
1308	ethyl	n-butyl	OH	H		H	7-dimethylamino
1309	n-butyl	n-butyl	OH	H	4-methoxyphenyl	H	9-fluoro
1310	ethyl	n-butyl	OH	H	phenyl	H	7-amino
1311	n-butyl	ethyl	OH	H	phenyl	H	7-(hydroxylamino)
1312	n-butyl	ethyl	OH	H	phenyl	H	8-hexyloxy
1313	n-butyl	ethyl	OH	H	phenyl	H	8-ethoxy
1314	ethyl	n-butyl	OH	H	phenyl	H	7-(hydroxylamino)
1315	ethyl	n-butyl	OH	H	phenyl	H	7-(hexyloxy)

WO 97/33882

PCT/US97/04076






1325	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1327	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1328	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1329	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1330	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1331	n-butyl	n-butyl	OH	H		7-dimethylamino	H

106

1316	n-butyl	ethyl	OH	H		7-dimethylamino	H
1317	n-butyl	ethyl	OH	H		7-dimethylamino	H
1318	ethyl	n-butyl	OH	H		7-dimethylamino	H
1319	ethyl	n-butyl	OH	H		7-dimethylamino	H
1320	ethyl	n-butyl	OH	H		7-dimethylamino	H
1321	n-butyl	ethyl	OH	H		7-dimethylamino	H
1322	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1323	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1324	n-butyl	n-butyl	OH	H		7-dimethylamino	H
1325	n-butyl	n-butyl	OH	H		7-dimethylamino	H

107





101

1332	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1333	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1334	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1335	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1336	n-butyl	n-butyl	OH	H		H	7-dimethylamino

WO 97/33882





PCT/US97/04076







101

1337	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1338	n-butyl	n-butyl	OH	H	4-methoxyphenyl	H	7-(4'-methoxypiperazinyl)
1339	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1340	n-butyl	ethyl	OH	H	5-piperonyl	H	7-methyl
1341	n-butyl	n-butyl	acetoxy	H	3-methoxyphenyl	H	7-dimethylamino
1342	n-butyl	n-butyl	OH	H	5-piperonyl	H	7-(4'-fluorophenyl)
1343	ethyl	n-butyl	OH	H	phenyl	H	7-amino
1344	n-butyl	n-butyl	OH	H	3-fluoro-4-methoxyphenyl	H	7-dimethylamino
1345	ethyl	n-butyl	OH	H	phenyl	H	7-trimethylammonium iodide
1346	ethyl	n-butyl	OH	H	phenyl	H	 at the 8-position
1347	n-butyl	n-butyl	OH	H	3-fluoro-4-methoxyphenyl	H	7-dimethylamino
1348	isobutyl	isobutyl	OH	H	phenyl	H	7-dimethylamino
1349	ethyl	n-butyl	OH	H	phenyl	H	7-dimethylamino
1350	n-butyl	n-butyl	OH	H	3-fluoro-4-methoxyphenyl	H	7-trimethylammonium iodide
1351	n-butyl	n-butyl	OH	H		H	7-dimethylamino






WO 97/33882

PCT/US97/04076

1358	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1359	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1360	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1361	n-butyl	n-butyl	OH	H		H	7-dimethylamino

1352	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1353	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1354	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1355	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1356	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1357	n-butyl	n-butyl	OH	H		H	7-dimethylamino





111

1362	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1363	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1364	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1365	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1366	n-butyl	n-butyl	OH	H		H	7-dimethylamino

WO 97/33882







PCT/US97/04076

112






1367	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1368	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1369	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1370	n-butyl	n-butyl	OH	H		H	7-dimethylamino

WO 97/33882

PCT/US97/04076






1376	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1377	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1378	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1379	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1380	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1381	n-butyl	n-butyl	OH	H		H	7-dimethylamino

111

1371	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1372	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1373	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1374	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1375	n-butyl	n-butyl	OH	H		H	7-dimethylamino

113






115

1382	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1383	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1384	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1385	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1386	n-butyl	n-butyl	OH	H		H	7-dimethylamino

WO 97/33882






PCT/US97/04076

116






1387	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1388	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1389	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1390	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1391	n-butyl	n-butyl	OH	H		H	7-dimethylamino

WO 97/33882






PCT/US97/04076






1397	n-butyl	n-butyl	OH	H		7-dimethylamino
1398	n-butyl	n-butyl	OH	H		7-dimethylamino
1399	n-butyl	n-butyl	OH	H		7-dimethylamino
1400	n-butyl	n-butyl	OH	H		7-dimethylamino
1401	n-butyl	n-butyl	OH	H		7-dimethylamino

811

1392	n-butyl	n-butyl	OH	H		7-dimethylamino
1393	n-butyl	n-butyl	OH	H		7-dimethylamino
1394	n-butyl	n-butyl	OH	H		7-dimethylamino
1395	n-butyl	n-butyl	OH	H		7-dimethylamino
1396	n-butyl	n-butyl	OH	H		7-dimethylamino

411

1402	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1403	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1404	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1405	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1406	n-butyl	n-butyl	OH	H		H	7-dimethylamino






1407	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1408	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1409	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1410	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1411	n-butyl	n-butyl	OH	H		H	7-dimethylamino






1417	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1418	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1419	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1420	n-butyl	n-butyl	OH	H		H	7-dimethylamino

122

1412	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1413	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1414	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1415	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1416	n-butyl	n-butyl	OH	H		H	7-dimethylamino

121

1421	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1422	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1423	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1424	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1425	n-butyl	n-butyl	OH	H		H	7-dimethylamino

1426	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1427	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1428	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1429	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1430	n-butyl	n-butyl	OH	H		H	7-dimethylamino







7-dimethylamino	H		H	OH	n-butyl	n-butyl	1436
7-dimethylamino	H		H	OH	n-butyl	n-butyl	1437
7-dimethylamino	H		H	OH	n-butyl	n-butyl	1438
7-dimethylamino	H		H	OH	n-butyl	n-butyl	1439
7-dimethylamino	H		H	OH	n-butyl	n-butyl	1440
7-dimethylamino	H		H	OH	n-butyl	n-butyl	1441

126

7-dimethylamino	H		H	OH	n-butyl	n-butyl	1431
7-dimethylamino	H		H	OH	n-butyl	n-butyl	1432
7-dimethylamino	H		H	OH	n-butyl	n-butyl	1433
7-dimethylamino	H		H	OH	n-butyl	n-butyl	1434
7-dimethylamino	H		H	OH	n-butyl	n-butyl	1435
7-dimethylamino	H		H	OH	n-butyl	n-butyl	1436

125




129

1442	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1443	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1444	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1445	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1446	n-butyl	n-butyl	OH	H		H	7-methoxy; 8-methoxy
1447	n-butyl	n-butyl	OH	H		H	7-dimethylamino

WO 97/33832


PCT/US97/04076

128


1448	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1449	n-butyl	n-butyl	OH	H		H	7-dimethylamino
1450	n-butyl	n-butyl	OH	H	phenyl	H	7-dimethylamino
1451	n-butyl	n-butyl	OH	H		H	7-dimethylamino

WO 97/33832


PCT/US97/04076




PEG = 3400 molecular weight polyethylene glycol polymer chain















PEG = 3400 molecular weight polyethylene glycol polymer chain















PEG = 3400 molecular weight polyethylene glycol polymer chain


























141



142



143

In further compounds of the present invention, R³ and R⁴ are independently selected from among hydrogen and ring-carbon substituted or unsubstituted aryl, thiophene, pyridine, pyrrole, thiazole, imidazole, pyrazole, pyrimidine, morpholine, N-alkylpyridinium, N-alkylpiperazinium, N-alkylmorpholinium, or furan in which the substituent(s) are selected from among halo, hydroxyl, trihaloalkyl, alkoxy, amino, N-alkylamino, N,N-dialkylamino, quaternary ammonium salts, a C₁ to C₄ alkylene bridge having a quaternary ammonium salt substituted thereon, alkoxycarbonyl, aryloxy, carbonyl, alkylcarbonyloxy and arylcarbonyloxy, (O,O)-dioxoalkylene, -[O(CH₂)_x]X where x is 2 to 12, w is 2 or 3 and X comprises halo or a quaternary ammonium salt, thiophene, pyridine, pyrrole, thiazole, imidazole, pyrazole, or furan. The aryl group of R³ or R⁴ is preferably phenyl, phenylene, or benzene triyl, i.e., may be unsubstituted, mono-substituted, or di-substituted. Among the species which may constitute the substituents on the aryl ring of R³ or R⁴ are fluoro, chloro, bromo, methoxy, ethoxy, isopropoxy, trimethylammonium (preferably with an iodide or chloride counterion), methoxycarbonyl, ethoxycarbonyl, formyl, acetyl, propanoyl, (N)-hexyldimethylammonium, hexylenetriethylammonium, tri(oxyethylene)iodide, and tetra(oxyethylene)trimethyl-ammonium iodide, each substituted at the p-position, the m-position, or both of the aryl ring. Other substituents that can be present on a phenylene, benzene triyl or other aromatic ring include 3,4-dioxymethylene (5-membered ring) and

144


3,4-dioxyethylene (6- membered ring). Among compounds which have been or can be demonstrated to have desirable ileal bile acid transport inhibiting properties are those in which R' or R' is selected from phenyl, p-fluorophenyl, m-fluorophenyl, p-hydroxyphenyl, m-hydroxyphenyl, p-methoxyphenyl, m-methoxyphenyl, p-N,N-dimethylaminophenyl, m-N,N-dimethylaminophenyl, I' p-(CH₃), N'-phenyl, I' m-(CH₃), N'-phenyl, I' p-CH₃, N'-CH₃, (OCH₃CH₃), -O-phenyl, I' p-CH₃, N'-CH₃, (OCH₃CH₃), -O-phenyl, I' m-(N,N-dimethylpiperazinium) - (N')-CH₃, (OCH₃CH₃), -O-phenyl, 3-methoxy-4-fluorophenyl, thienyl-2-yl, 5-chlorothieryl-2-yl, 3,4-difluorophenyl, I' p-(N,N-dimethylpiperazinium) - (N')-CH₃, (OCH₃CH₃), -O-phenyl, 3-fluoro-4-methoxyphenyl, -4-pyridinyl, 2-pyridinyl, 3-pyridinyl, N-methyl-4-pyridinium, I' N-methyl-3-pyridinium, 3,4-dioxymethylenepheryl, 3,4-dioxyethylenepheryl, and p-methoxycarbonylphenyl. Preferred compounds include 3-ethyl-3-butyl and 3-butyl-3-butyl compounds having each of the above preferred R' substituents in combination with the R' substituents shown in Table 1. It is particularly preferred that one but not both of R' and R' is hydrogen.

It is especially preferred that R' and R' be hydrogen, that R' and R' not be hydrogen, and that R' and R' be oriented in the same direction relative to the plane of the molecule, i.e., both in a- or both in S-configuration. It is further preferred that, where R' is butyl and R' is ethyl, then R' has the same

orientation relative to the plane of the molecule as R' and R'.

Set forth in Table 1A are lists of species of R'/R', R'/R' and R'.

Table 1A : Alternative R groups



R ¹ , R ²	R ³ , R ⁴ , R ⁵	(R ⁶), R ⁷
ethyl	HO-	1-acetyl
n-propyl	Ph-	1-ethyl
n-butyl	p-F-Ph-	1-iso-propyl
n-pentyl	m-F-Ph-	1-tert-butyl
n-hexyl	p-CH ₃ O-Ph-	1-OH
iso-propyl	m-CH ₃ O-Ph-	1-OCH ₃
iso-butyl	p-(CH ₃) ₂ -Ph-	1-O(1-iso-propyl)
iso-pentyl	m-(CH ₃) ₂ -Ph-	1-SCN
CH ₃ C(=O)C ₂ H ₅	I ⁺ , p-(CH ₃) ₂ -Ph-	1-SOCH ₃
CH ₃ OC ₂ H ₅	I ⁺ , m-(CH ₃) ₂ -Ph-	1-SCNCH ₃
CH ₃ CH(OMe)C ₂ H ₅	I ⁺ , p-(CH ₃) ₂ -O-Ph-	1-NH ₂
CH ₃ O-(4-picolyl)	I ⁺ , m-(CH ₃) ₂ -O-Ph-	1-NHCH ₃
	I ⁺ , p-Ph-N-	1-N(CH ₃) ₂
	I ⁺ , m-Ph-N-	1-N(CH ₃) ₂ , I ⁺
	diacetylbipiperazine)-	1-NHC(=O)CH ₃
	(N ¹ -CH ₃ -(OCH ₂ CH ₂) ₂ -O-	1-N(CH ₃ CH ₂) ₂
	Ph-	1-NHCCH ₂ CO ₂ H
	I ⁺ , m-Ph-N-	1-N ⁺ (Me) ₂ CH ₂ CO ₂ H, I ⁺
	diacetylbipiperazine)-	1-(N)-morpholine
	(N ¹ -CH ₃ -(OCH ₂ CH ₂) ₂ -O-	1-(N)-azetidine
	Ph-	1-(N)-8-methylazetidine, I ⁺
	m-F, p-CH ₃ O-Ph-	1-(N)-pyrrolidine
	3,4-dioxymethylene-Ph	1-(N)-8-methyl-pyrrolidinum, I ⁺
	m-CH ₃ O-, p ⁺ -Ph-	1-(N)-8-methyl-morpholinum, I ⁺
	4-pyridine	1-(N)-N ⁺ -dimethylpiperazine
	8-methyl-4-pyridinium, I ⁺	1-(N)-N ⁺ -dimethylpiperazinium, I ⁺
	3-pyridine	7-NH-CH ₂
	8-methyl-3-pyridinium, I ⁺	7-NHC(=O)C ₂ H ₅
	2-pyridine	7-NHC(=O)CH ₂ Br
	p-CH ₃ OC-Ph-	7-NH-C(=O)NH ₂
	chloxy-2-yl	7-(2)-chlorophene
	3-Cl-tetrayl-3-yl	
	3,4-difluoro	
	m-F, p-CH ₃ O-Ph	

continued next page...

8-acetyl
 8-ethyl
 8-iso-propyl
 8-tert-butyl
 8-OH
 8-OC₂H₅
 8-O(1-iso-propyl)
 8-SCN
 8-SOCH₃
 8-SOCH₃
 8-SCNCH₃
 8-NH₂
 8-NHCH₃
 8-NHCH₃
 8-N(CH₃)₂
 8-N(CH₃)₂, I⁺
 8-NHC(=O)CH₃
 8-N(CH₂CH₂)₂
 8-NHCCH₂CO₂H
 8-N⁺(Me)₂CH₂CO₂H, I⁺
 8-(N)-morpholine
 8-(N)-azetidine
 8-(N)-8-methylazetidine, I⁺
 8-(N)-pyrrolidine
 8-(N)-8-methyl-pyrrolidinum, I⁺
 8-(N)-8-methyl-morpholinum, I⁺
 8-(N)-N⁺-dimethylpiperazine
 8-(N)-N⁺-dimethylpiperazinium, I⁺
 8-NH-CH₂
 8-NHC(=O)C₂H₅
 8-NHC(=O)CH₂Br
 8-NH-C(=O)NH₂
 8-(2)-chlorophene

continued next page...

The core moiety can comprise alkane diyl, alkene diyl, alkene diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carboxylate, amino acid, and peptide, polypeptide, wherein alkane diyl, alkene diyl, alkene diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carboxylate, amino acid, and peptide polypeptide, can optionally have one or more carbon replaced by O, NR¹, NR², S, SO, SO₂, SR¹, PR¹, P+R/R8, phenylene, heterocycle, quaternary heterocycle, quaternary heteroaryl, or aryl.

wherein alkane diyl, alkene diyl, alkene diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carboxylate, amino acid, peptide, and polypeptide can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴, P⁺R¹³R¹⁴R¹⁵A⁻, P(OR¹³)OR¹⁴, S⁺R¹³A⁻, and N⁺R⁹R¹¹R¹²A⁻;





wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of OR⁷, NR⁷R⁸, SR⁷, S(O)R⁷, SO₂R⁷, CO₂R⁷, CN, oxo, CONR⁷R⁸, N⁺R⁷R⁸A⁻, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, P(O)R⁷R⁸, P⁺R⁷R⁸A⁻, and P(O)(OR⁷)OR⁸, and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O,

(51)

NR⁷, N⁺R⁷R⁸A⁻, S, SO, SO₂, S⁺R⁷A⁻, PR⁷, P(O)R⁷, P⁺R⁷R⁸A⁻, or phenylene.

Exemplary core moieties include:






wherein:

R⁷ is selected from the group consisting of C and N, and

(52)

R³⁰ and R³¹ are independently selected from the group consisting of:



wherein R³⁰, R³¹, R³² and R³³ are independently selected from alkyl, alkenyl, alkylaryl, aryl, arylalkyl, cycloalkyl, heterocycle, and heterocycloalkyl.

A³⁰ is a pharmaceutically acceptable anion, and k = 1 to 10.

In compounds of Formula DIV, R³⁰, R³¹, R³² in Formulae DII and DIII, and R³³ in Formula DIII can be bonded at any of their 6-, 7-, 8-, or 9- positions to R³⁴. In compounds of Formula DIVA, it is preferred that R³⁵ comprises a phenylene moiety bonded at a m- or p-position thereof to R³⁶.

In another embodiment, a core moiety backbone, R³⁷, as discussed herein in Formulas DII and DIII can be multiply substituted with more than four pendant active benzothiepine units, i.e., R³⁸, R³⁹, R⁴⁰, R⁴¹, and R⁴² as discussed above, through multiple functional groups within the core moiety backbone. The core moiety backbone unit, R⁴³, can comprise a single core moiety unit, multimers thereof, and multimeric mixtures of the different core moiety units discussed herein, i.e., alone or in combination. The number of individual core

moiety backbone units can range from about one to about 100, preferably about one to about 80, more preferably about one to about 50, and even more preferably about one to about 25. The number of points of attachment of similar or different pendant active benzothiepine units within a single core moiety backbone unit can be in the range from about one to about 100, preferably about one to about 80, more preferably about one to about 50, and even more preferably about one to about 25. Such points of attachment can include bonds to C, S, O, N, or P within any of the groups encompassed by the definition of R⁴⁴.

The more preferred benzothiepine moieties comprising R⁴⁵, R⁴⁶, R⁴⁷ and/or R⁴⁸ conform to the preferred structures as outlined above for Formula I. The 3-carbon on each benzothiepine moiety can be achiral, and the substituents R⁴⁹, R⁵⁰, R⁵¹, R⁵², R⁵³ and R⁵⁴ can be selected from the preferred groups and combinations of substituents as discussed above. The core structures can comprise, for example, poly(oxalkylene) or oligo(oxalkylene), especially poly- or oligo(oxethylene) or poly- or oligo(oxypropylene).

Dosages, Formulations, and Routes of Administration

The ileal bile acid transport inhibitor compounds of the present invention can be administered for the prophylaxis and treatment of hyperlipidemic diseases or conditions by any means, preferably oral, that produce contact of these compounds with their site of action in the body, for example in the ileum of a mammal, e.g., a human.

For the prophylaxis or treatment of the conditions referred to above, the compounds of the present invention can be used as the compound *per se*.

Pharmaceutically acceptable salts are particularly suitable for medical applications because of their greater aqueous solubility relative to the parent compound. Such salts must clearly have a

pharmaceutically acceptable anion or cation. Suitable pharmaceutically acceptable acid addition salts of the compounds of the present invention when possible include those derived from inorganic acids, such as hydrochloric, hydrobromic, phosphoric, metaphosphoric, nitric, sulfuric, and sulfuric acids, and organic acids such as acetic, benzenesulfonic, benzoic, citric, ethanesulfonic, fumaric, gluconic, glycolic, isochloric, lactic, lactobionic, maleic, malic, methanesulfonic, succinic, toluenesulfonic, tartaric, and trifluoroacetic acids. The chloride salt is particularly preferred for medical purposes. Suitable pharmaceutically acceptable base salts include ammonium salts, alkali metal salts such as sodium and potassium salts, and alkaline earth salts such as magnesium and calcium salts.

The anions of the definition of A in the present invention are, of course, also required to be pharmaceutically acceptable and are also selected from the above list.

The compounds of the present invention can be presented with an acceptable carrier in the form of a pharmaceutical composition. The carrier must, of course, be acceptable in the sense of being compatible with the other ingredients of the composition and must not be deleterious to the recipient. The carrier can be a solid or a liquid, or both, and is preferably formulated with the compound as a unit-dose composition, for example, a tablet, which can contain from 0.05% to 95% by weight of the active compound. Other pharmacologically active substances can also be present, including other compounds of the present invention. The pharmaceutical compositions of the invention can be prepared by any of the well known techniques of pharmacy, consisting essentially of admixing the components.

These compounds can be administered by any conventional means available for use in conjunction

155

with pharmaceuticals, either as individual therapeutic compounds or as a combination of therapeutic compounds.

The amount of compound which is required to achieve the desired biological effect will, of course, depend on a number of factors such as the specific compound chosen, the use for which it is intended, the mode of administration, and the clinical condition of the recipient.

In general, a daily dose can be in the range of from about 0.3 to about 100 mg/kg bodyweight/day, preferably from about 1 mg to about 50 mg/kg bodyweight/day, more preferably from about 3 to about 10 mg/kg bodyweight/day. This total daily dose can be administered to the patient in a single dose, or in proportionate multiple subdoses. Subdoses can be administered 2 to 6 times per day. Doses can be in sustained release form effective to obtain desired results.

Orally administrable unit dose formulations, such as tablets or capsules, can contain, for example, from about 0.1 to about 100 mg of benzothiepine compound, preferably about 1 to about 75 mg of compound, more preferably from about 10 to about 50 mg of compound. In the case of pharmaceutically acceptable salts, the weights indicated above refer to the weight of the benzothiepine ion derived from the salt.

Oral delivery of an ileal bile acid transport inhibitor of the present invention can include formulations, as are well known in the art, to provide prolonged or sustained delivery of the drug to the gastrointestinal tract by any number of mechanisms. These include, but are not limited to, pH sensitive release from the dosage form based on the changing pH of the small intestine, slow erosion of a tablet or capsule, retention in the stomach based on the physical properties of the formulation, bioadhesion of the dosage form to the mucosal lining of the intestinal tract, or enzymatic release of the active drug from the dosage form. The intended effect is to extend the time

156

period over which the active drug molecule is delivered to the site of action (the ileum) by manipulation of the dosage form. Thus, enteric-coated and enteric-coated controlled release formulations are within the scope of the present invention. Suitable enteric coatings include cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropylmethylcellulose phthalate and anionic polymers of methacrylic acid and methacrylic acid methyl ester.

When administered intravenously, the dose can, for example, be in the range of from about 0.1 mg/kg body weight to about 1.0 mg/kg body weight, preferably from about 0.25 mg/kg body weight to about 0.75 mg/kg body weight, more preferably from about 0.4 mg/kg body weight to about 0.6 mg/kg body weight. This dose can be conveniently administered as an infusion of from about 10 ng/kg body weight to about 100 ng/kg body weight per minute. Infusion fluids suitable for this purpose can contain, for example, from about 0.1 ng to about 10 mg, preferably from about 1 ng to about 10 mg per milliliter. Unit doses can contain, for example, from about 1 mg to about 10 g of the compound of the present invention. Thus, ampoules for injection can contain, for example, from about 1 mg to about 100 mg.

Pharmaceutical compositions according to the present invention include those suitable for oral, rectal, topical, buccal (e.g., sublingual), and parenteral (e.g., subcutaneous, intramuscular, intradermal, or intravenous) administration, although the most suitable route in any given case will depend on the nature and severity of the condition being treated and on the nature of the particular compound which is being used. In most cases, the preferred route of administration is oral.

Pharmaceutical compositions suitable for oral administration can be presented in discrete units, such as capsules, cachets, lozenges, or tablets, each containing a predetermined amount of at least one

compound of the present invention; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water or water-in-oil emulsion. As indicated, such compositions can be prepared by any suitable method of pharmacy which includes the step of bringing into association the active compound(s) and the carrier (which can constitute one or more accessory ingredients). In general, the compositions are prepared by uniformly and intimately admixing the active compound with a liquid or finely divided solid carrier, or both, and then, if necessary, shaping the product. For example, a tablet can be prepared by compressing or molding a powder or granules of the compound, optionally with one or more accessory ingredients. Compressed tablets can be prepared by compressing, in a suitable machine, the compound in a free-flowing form, such as a powder or granules optionally mixed with a binder, lubricant, inert diluent and/or surface active/dispersing agent(s). Molded tablets can be made by molding, in a suitable machine, the powdered compound moistened with an inert liquid diluent.

Pharmaceutical compositions suitable for buccal (sub-lingual) administration include lozenges comprising a compound of the present invention in a flavored base, usually sucrose, and acacia or tragacanth, and pastilles comprising the compound in an inert base such as gelatin and glycerin or sucrose and acacia.

Pharmaceutical compositions suitable for parenteral administration conveniently comprise sterile aqueous preparations of a compound of the present invention. These preparations are preferably administered intravenously, although administration can also be effected by means of subcutaneous, intramuscular, or intradermal injection. Such preparations can conveniently be prepared by admixing the compound with water and rendering the resulting solution sterile and isotonic with the blood.

Injectable compositions according to the invention will generally contain from 0.1 to 5% w/w of a compound disclosed herein.

Pharmaceutical compositions suitable for rectal administration are preferably presented as unit-dose suppositories. These can be prepared by admixing a compound of the present invention with one or more conventional solid carriers, for example, cocoa butter, and then shaping the resulting mixture.

Pharmaceutical compositions suitable for topical application to the skin preferably take the form of an ointment, cream, lotion, paste, gel, spray, aerosol, or oil. Carriers which can be used include vaseline, lanoline, polyethylene glycols, alcohols, and combinations of two or more thereof. The active compound is generally present at a concentration of from 0.1 to 15% w/w of the composition, for example, from 0.5 to 2%.

Transdermal administration is also possible.

Pharmaceutical compositions suitable for transdermal administration can be presented as discrete patches adapted to remain in intimate contact with the epidermis of the recipient for a prolonged period of time. Such patches suitably contain a compound of the present invention in an optionally buffered, aqueous solution, dissolved and/or dispersed in an adhesive, or dispersed in a polymer. A suitable concentration of the active compound is about 1% to 35%, preferably about 3% to 15%. As one particular possibility, the compound can be delivered from the patch by electrotransport or iontophoresis, for example, as described in *Pharmaceutical Research*, 3(6), 318 (1986).

In any case, the amount of active ingredient that can be combined with carrier materials to produce a single dosage form to be administered will vary depending upon the host treated and the particular mode of administration.

The solid dosage forms for oral administration including capsules, tablets, pills, powders, and

159

granules noted above comprise one or more compounds of the present invention admixed with at least one inert diluent such as sucrose, lactose, or starch. Such dosage forms may also comprise, as in normal practice, additional substances other than inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

Liquid dosage forms for oral administration can include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions may also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or settling agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Pharmaceutically acceptable carriers encompass all the foregoing and the like.

Treatment Regimen

The dosage regimen to prevent, give relief from, or ameliorate a disease condition having hyperlipemia as an element of the disease, e.g., atherosclerosis, or

160

to protect against or treat further high cholesterol plasma or blood levels with the compounds and/or compositions of the present invention is selected in accordance with a variety of factors. These include the type, age, weight, sex, diet, and medical condition of the patient, the severity of the disease, the route of administration, pharmacological considerations such as the activity, efficacy, pharmacokinetics and toxicology profiles of the particular compound employed, whether a drug delivery system is utilized, and whether the compound is administered as part of a drug combination. Thus, the dosage regimen actually employed may vary widely and therefore deviate from the preferred dosage regimen set forth above.

Initial treatment of a patient suffering from a hyperlipidemic condition can begin with the dosages indicated above. Treatment should generally be continued as necessary over a period of several weeks to several months or years until the hyperlipidemic disease condition has been controlled or eliminated. Patients undergoing treatment with the compounds or compositions disclosed herein can be routinely monitored by, for example, measuring serum cholesterol levels by any of the methods well known in the art, to determine the effectiveness of therapy. Continuous analysis of such data permits modification of the treatment regimen during therapy so that optimal effective amounts of compounds of the present invention are administered at any point in time, and so that the duration of treatment can be determined as well. In this way, the treatment regimen/dosing schedule can be rationally modified over the course of therapy so that the lowest amount of ileal bile acid transport inhibitor of the present invention which exhibits satisfactory effectiveness is administered, and so that administration is continued only so long as is necessary to successfully treat the hyperlipidemic condition.

161

The following non-limiting examples serve to illustrate various aspects of the present invention.

EXAMPLES OF SYNTHETIC PROCEDURES

Preparation 1

2-Ethyl-2-(mesyloxymethyl)hexanal (1)

To a cold (10 °C) solution of 12.6 g (0.11 mole) of methanesulfonyl chloride and 10.3 g (0.13 mole) of triethylamine was added dropwise 15.8 g of 2-ethyl-2-(hydroxymethyl)hexanal, prepared according to the procedure described in Chem. Ber. 98, 728-734 (1965), while maintaining the reaction temperature below 30 °C. The reaction mixture was stirred at room temperature for 18 h, quenched with dilute HCl and extracted with methylene chloride. The methylene chloride extract was dried over MgSO_4 and concentrated in vacuo to give 24.4 g of brown oil.

Preparation 2

2-((2-Benzoylphenylthio)methyl)-2-ethylhexanal (2)


A mixture of 31 g (0.144 mol) of 2-mercaptobenzophenone, prepared according to the procedure described in WO 93/16055, 24.4 g (0.1 mole) of 2-ethyl-2-(mesyloxymethyl)-hexanal (1), 14.8 g (0.146 mole) of triethylamine, and 80 mL of 2-methoxyethyl ether was held at reflux for 24 h. The reaction mixture was poured into 3N HCl and extracted

162

with 300 mL of methylene chloride. The methylene chloride layer was washed with 300 mL of 10% NaOH, dried over MgSO₄ and concentrated in vacuo to remove 2-methoxyethyl ether. The residue was purified by HPLC (10% EtOAc-hexane) to give 20.5 g (58%) of 2 as an oil.

Example 1

3-Butyl-3-ethyl-5-phenyl-2,3-dihydrobenzothiepine (3), *cis*-3-Butyl-3-ethyl-5-phenyl-2,3-dihydrobenzothiepine- (5*H*) 4-one (4a) and *trans*-3-Butyl-3-ethyl-5-phenyl-2,3-dihydro-benzothiepine- (5*H*) 4-one (4b)



A mixture of 2.6 g (0.04 mole) of zinc dust, 7.2 g (0.047 mole) of TiCl₄, and 80 mL of anhydrous ethylene glycol dimethyl ether (DME) was held at reflux for 2 h. The reaction mixture was cooled to 5 °C. To the reaction mixture was added dropwise a solution of 3.54 g (0.01 mole) of 2 in 30 mL of DME in 40 min. The reaction mixture was stirred at room temperature for 16 h and then was held at reflux for 2 h and cooled before being poured into brine. The organic was extract into methylene chloride. The methylene chloride extract was dried over MgSO₄ and concentrated in vacuo. The residue was purified by HPLC (hexane) to give 1.7 g (43%) of 3 as an oil in the first fraction. The second fraction was discarded and the third fraction was further purified by HPLC (hexane) to give 0.07 g (2%) of 4a in the earlier fraction and 0.1 g (3%) of 4b in the later fraction.

Example 2

cis-3-Butyl-3-ethyl-5-phenyl-2,3-dihydrobenzothiepine- (5*H*) 4-one-1,1-dioxide (5a) and *trans*-3-Butyl-3-ethyl-5-phenyl-2,3-dihydro-benzothiepine- (5*H*) 4-one-1,1-dioxide (5b)




To a solution of 1.2 g (3.5 mmole) of 50-60% MCPBA in 20 mL of methylene chloride was added 0.59 g (1.75

163

mmole) of a mixture of 4a and 4b in 10 mL of methylene chloride. The reaction mixture was stirred for 20 h. An additional 1.2 g (1.75 mmole) of 50-60% MCPBA was added and the reaction mixture was stirred for an additional 3 h then was triturated with 50 mL of 10% NaOH. The insoluble solid was filtered. The methylene chloride layer of the filtrate was washed with brine, dried over MgSO₄, and concentrated in vacuo. The residual syrup was purified by HPLC (5% EtOAc-hexane) to give 0.2 g (30%) of 5a as an oil in the first fraction and 0.17 g (26%) of 5b as an oil in the second fraction.

Example 3

(3a, 4a, 5b) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (6a), (3a, 4b, 5a) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (6b), (3a, 4a, 5a) 3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (6c), and (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (6d)



A. Reduction of 5a and 5b with Sodium Borohydride

To a solution of 0.22 g (0.59 mmole) of 5b in 10 mL of ethanol was added 0.24 g (6.4 mmole) of sodium borohydride. The reaction mixture was stirred at room temperature for 18 h and concentrated in vacuo to remove ethanol. The residue was triturated with water and extracted with methylene chloride. The methylene chloride extract was dried over MgSO₄ and concentrated in vacuo to give 0.2 g of syrup. In a separate experiment, 0.45 g of 5a was treated with 0.44 g of sodium borohydride in 10 mL of ethanol and was worked up as described above to give 0.5 g of syrup which was identical to the 0.2 g of syrup obtained above. These two materials were combined and purified by HPLC using 10% EtOAc-hexane as eluent. The first fraction was 0.18 g (27%) of 6a as a syrup. The second fraction was 0.2 g

164

(30%) of **6b** also as a syrup. The column was then eluted with 20% EtOAc-hexane to give 0.077 g (11%) of **6c** in the third fraction as a solid. Recrystallization from hexane gave a solid, mp 179-181 °C. Finally, the column was eluted with 30% EtOAc-hexane to give 0.08 g (12%) of **6d** in the fourth fraction as a solid.

Recrystallization from hexane gave a solid, mp 160-161 °C.

B. Conversion of **6a** to **6c** and **6d** with NaOH and PTC


To a solution of 0.29 g (0.78 mmole) of **6a** in 10 mL CH_2Cl_2 , was added 9 g of 40% NaOH. The reaction mixture was stirred for 0.5 h at room temperature and was added one drop of Aliquat-336 (methyltricaprylammonium chloride) phase transfer catalyst (PTC). The mixture was stirred for 0.5 h at room temperature before being treated with 25 mL of ice-crystals then was extracted with CH_2Cl_2 (3x10 mL), dried over MgSO_4 and concentrated in vacuo to recover 0.17 g of a colorless film. The components of this mixture were separated using an HPLC and eluted with EtOAc-hexane to give 12.8 mg (4%) of 2-(2-benzylphenylsulfonylmethyl)-2-ethylhexenal in the first fraction, 30.9 mg (11%) of **6c** in the second fraction and 90.0 mg (31%) of **6d** in the third fraction.

Oxidation of **6a** to **5b**

To a solution of 0.20 g (0.52 mmole) of **6a** in 5 mL of CH_2Cl_2 , was added 0.23 g (1.0 mmole) of pyridinium chlorochromate. The reaction mixture was stirred for 2 h then was treated with additional 0.23 g of pyridinium chlorochromate and stirred overnight. The dark reaction mixture was poured into a ceramic filterfrit containing silica gel and was eluted with CH_2Cl_2 . The filtrate was concentrated in vacuo to recover 167 mg (87%) of **5b** as a colorless oil.

Example 4

3-Butyl-3-ethyl-5-phenyl-2,3-dihydrobenzothiepine-1,1-dioxide (7)



To a solution of 5.13 g (15.9 mmole) of **3** in 50 mL of CH_2Cl_2 , was added 10 g (31.9 mmole) of 50-60% MCPBA (m-chloroperoxybenzoic acid) portionwise causing a mild reflux and formation of a white solid. The reaction mixture was allowed to stir overnight under N_2 and was triturated with 25 mL of water followed by 50 mL of 10% NaOH solution. The organic was extracted into CH_2Cl_2 (4x20 mL). The CH_2Cl_2 extract was dried over MgSO_4 and evaporated to dryness to recover 4.9 g (87%) of an opaque viscous oil.

Example 5


(1a, 2b, 8a) 2-Butyl-2-ethyl-8b-phenyl-1a, 2, 3, 8b-tetrahydro-benzothiepine[4, 5-b]oxirano-4, 4-dioxide (8a)
(1a, 2a, 8a) 2-Butyl-2-ethyl-8b-phenyl-1a, 2, 3, 8b-tetrahydro-benzothiepine [4, 5-b]oxirano-4, 4-dioxide



To 1.3 g (4.03 mole) of **3** in 25 mL of CH_2Cl_2 , was added portionwise 5 g (14.1 mmole) of 50-60 % MCPBA causing a mild exotherm. The reaction mixture was stirred under N_2 overnight and was then held at reflux for 3 h. The insoluble white slurry was filtered. The filtrate was extracted with 10% potassium carbonate (3x50 mL), once with brine, dried over MgSO_4 , and concentrated in vacuo to give 1.37 g of a light yellow oil. Purification by HPLC gave 0.65 g of crystalline product. This product is a mixture of two isomers. Trituration of this crystalline product in hexane recovered 141.7 mg (10%) of a white crystalline product. This isomer was characterized by NMR and mass spectra to be the (1a, 2b, 8a) isomer **8a**. The hexane filtrate was concentrated in vacuo to give 206 mg of white film which is a mixture of 30% **8a** and 70% **8b** by ^1H NMR.

Example 6

cis-3-Butyl-3-ethyl-5-phenyl-2,3,4,5-tetrahydrobenzochlopiene-1,1-dioxide (9a), trans-3-Butyl-3-ethyl-5-phenyl-2,3,4,5-tetrahydrobenzochlopiene-1,1-dioxide (9b), and 3-Butyl-3-ethyl-4-hydroxy-5-cyclohexylidene-2,3,4,5-tetrahydrobenzochlopiene-1,1-dioxide (10)



A mixture of 0.15 g (0.4 mmole) of a 3:7 mixture of 8a and 8b was dissolved in 15 ml MeOH in a 3 oz.

Fisher/Porter vessel, then was added 0.1 g of 10% Pd/C catalyst. This mixture was hydrogenated at 70 psi H₂ for 5 h and filtered. The filtrate was evaporated to dryness in vacuo to recover 0.117 g of a colorless oil. This material was purified by HPLC eluting with EtOAc-hexane. The first fraction was 4.2 mg (3%) of 9b. The second fraction, 5.0 mg (4%), was a 50/50 mixture of 9a and 9b. The third fraction was 8.8 mg (6%) of 6a. The fourth fraction was 25.5 mg (18%) of 6b. The fifth fraction was 9.6 mg (7%) of a mixture of 6b and a product believed to be 3-butyl-3-ethyl-4,5-dihydroxy-5-phenyl-2,3,4,5-tetrahydrobenzochlopiene-1,1-dioxide based on mass spectrum. The sixth fraction was 7.5 mg (5%) of a mixture of 6d and one of the isomers of 10, 10a.


Example 7

In another experiment, a product (3.7 g) from epoxidation of 3 with excess MCPBA in refluxing CHCl₃, under air was hydrogenated in 100 mL of methanol using 1 g of 10% Pd/C catalyst and 70 psi hydrogen. The product was purified by HPLC to give 0.9 g (25%) of 9b, 0.45 g (13%) of 9a, 0.27 g (7%) of 6a, 0.51 g (14%) of 6b, 0.02 g (1%) of 6c, 0.06 g (2%) of one isomer of 10, 10a and 0.03 g (1%) of another isomer of 10, 10b.

167

Example 8


2-((2-Benzoylphenylthio)methyl)butyraldehyde (11)



To an ice bath cooled solution of 9.76 g (0.116 mole) of 2-ethylacrolein in 40 mL of dry THF was added 24.6 g (0.116 mole) of 2-mercaptobenzophenone in 40 mL of THF followed by 13 g (0.128 mole) of triethylamine. The reaction mixture was stirred at room temperature for 3 days, diluted with ether, and was washed successively with dilute HCl, brine, and 1 M potassium carbonate. The ether layer was dried over MgSO₄ and concentrated in vacuo. The residue was purified by HPLC (10% EtOAc-hexane) to give 22 g (64%) of 11 in the second fraction. An attempt to further purify this material by Kugelrohr distillation at 0.5 torr (160-190 °C) gave a fraction (12.2 g) which contained starting material indicating a reversed reaction during distillation. This material was dissolved in ether (100 mL) and was washed with 50 mL of 1 M potassium carbonate three times to give 6.0 g of a syrup which was purified by HPLC (10% EtOAc-hexane) to give 5.6 g of pure 11.

Example 9

3-Ethyl-5-phenyl-2,3-dihydrobenzochlopiene (12)



To a mixture of 2.61 g (0.04 mole) of zinc dust and 60 mL of DME was added 7.5 g (0.048 mole) of TiCl₄. The reaction mixture was held at reflux for 2 h. A solution of 2.98 g (0.01 mole) of 11 was added dropwise in 1 h. The reaction mixture was held at reflux for 18 h, cooled and poured into water. The organic was extracted into ether. The ether layer was washed with brine and filtered through Celite. The filtrate was dried over MgSO₄ and concentrated. The residual oil (2.5 g) was purified by HPLC to give 2.06 g (77%) of 12 as an oil in the second fraction.

168

Example 10


(1a, 2a, 8b) 2-Ethyl-8b-phenyl-1a, 2, 3, 8b-tetrahydro-
benzothiepine-[(4, 5-b)oxirene-4, 4-dioxide (13)]



To a solution of 1.5 g (5.64 mmole) of 12 in 25 ml of
CHCl₃, was added 6.8 g (19.4 mmole) of 50-60% MCPB
portionwise causing an exotherm and formation of a white
solid. The mixture was stirred at room temperature
overnight diluted with 100 ml methylene chloride and
washed successively with 10% K₂CO₃ (4x50 ml), water
(twice with 25 ml) and brine. The organic layer was
then dried over MgSO₄ and evaporated to dryness to
recover 1.47 g of an off white solid. ¹H NMR indicated
that only one isomer is present. This solid was
slurried in 200 ml of warm Et₂O and filtered to give
0.82 g (46%) of 13 as a white solid, mp 185-186.5 °C.

Example 11

(3a, 4b, 5a) 3-Ethyl-4-hydroxy-5-phenyl-2, 3, 4, 5-
tetrahydro-benzothiepine-1, 1-dioxide (14a), (3a, 4b, 5b)
3-Ethyl-4-hydroxy-5-phenyl-2, 3, 4, 5-
tetrahydrobenzothiepine-1, 1-dioxide (14b), and cis-3-
ethyl-5-phenyl-2, 3, 4, 5-tetrahydro-benzothiepine-1, 1-
dioxide (15)



A mixture of 0.5 g (1.6 mole) of 13, 50 ml of acetic
acid and 0.5 g of 10% Pd/C catalyst was hydrogenated
with 70 psi hydrogen for 4 h. The crude reaction slurry
was filtered and the filtrate was stirred with 150 ml
of a saturated NaHCO₃ solution followed by 89 g of
NaHCO₃ powder portionwise to neutralize the rest of
acetic acid. The mixture was extracted with methylene
chloride (4x25 ml), then the organic layer was dried
over MgSO₄ and concentrated in vacuo to give 0.44 g
(87%) of a voluminous white solid which was purified by
HPLC (EtOAc-Hexane) to give 26.8 mg (6%) of 15 in the
first fraction, 272 mg (54%) of 14a as a solid, mp 142-

169

Example 12

2-Ethyl-2-((2-Hydroxyethylphenyl)thiomethyl)hexenal
(16)



A mixture of 5.0 g (0.036 mole) of 2-mercaptobenzyl
alcohol, 6.4 g (0.032 mole) of 1, 3, 6 g (0.036 mole) of
triethylamine and 25 ml of 2-methoxyethyl ether was
held at reflux for 7 h. Additional 1.1 g of
mercaptobenzyl alcohol and 0.72 g of triethylamine was
added to the reaction mixture and the mixture was held
at reflux for additional 16 h. The reaction mixture was
cooled and poured into 6N HCl and extracted with
methylene chloride. The methylene chloride extract was
washed twice with 10% NaOH, dried over MgSO₄ and
concentrated in vacuo to give 9.6 g of residue.
Purification by HPLC (20% EtOAc-hexane) gave 3.7 g
(41%) of 16 as an oil.

Example 13


2-Ethyl-2-((2-formylphenyl)thiomethyl)hexenal (17)



A mixture of 3.7 g of 16, 5.6 g (0.026 mole) of
pyridinium chlorochromate, 2 g of Celite and 30 mL of
methylene chloride was stirred for 18 h and filtered
through a bed of silica gel. The silica gel was eluted
with methylene chloride. The combined methylene
chloride eluant was purified by HPLC (20% EtOAc-hexane)
to give 2.4 g (66%) of an oil.

Example 14

3-Butyl-3-ethyl-2, 3-dihydrobenzothiepine (18)



A mixture of 2.6 g (0.04 mole) of zinc dust, 7.2 g
(0.047 mole) of TiCl₄, and 50 mL of DME was held at

170

reflux for 2 h and cooled to room temperature. To this mixture was added 2.4 g (8.6 mmole) of 17 in 20 mL of DME in 10 min. The reaction mixture was stirred at room temperature for 2 h and held at reflux for 1 h then was let standing at room temperature over weekend. The reaction mixture was poured into dilute HCl and was stirred with methylene chloride. The methylene chloride-water mixture was filtered through Celite. The methylene chloride layer was washed with brine, dried over MgSO_4 , and concentrated in vacuo to give 3.0 g of a residue. Purification by HPLC gave 0.41 g (20%) of 18 as an oil in the early fraction.

Example 15

(1a, 2a, 8ba) 2-Butyl-2-ethyl-1a,2,3,8b-tetrahydro-benzothiepine(4,5-b)oxirane-6,4-dioxide (19a) and (1ac, 2b, 8ba) 2-Butyl-2-ethyl-8b-phenyl-1a,2,3,8b-tetrahydro-benzothiepine(4,5-b)oxirane-6,4-dioxide (19b)



To a solution of 0.4 g of 0.4 g (1.6 mmole) of 18 in 30 mL of methylene chloride was added 2.2 g (3.2 mmole) of 50-60% MCPBA. The reaction mixture was stirred for 2 h and concentrated in vacuo. The residue was dissolved in 30 mL of CHCl_3 and was held at reflux for 18 h under N_2 . The reaction mixture was stirred with 100 mL of 10% NaOH and 5 g of sodium sulfite. The methylene chloride layer was washed with brine, dried over MgSO_4 , and concentrated in vacuo. The residue was purified by HPLC (20% EtOAc-hexane) to give a third fraction which was further purified by HPLC (10% EtOAc-hexane) to give 0.12 g of syrup in the first fraction.

Recrystallization from hexane gave 0.08 g (17%) of 19a, mp 89.5-105.5 °C. The mother liquor from the first fraction was combined with the second fraction and was further purified by HPLC to give additional 19a in the first fraction and 60 mg of 19b in the second fraction.


171

Crystallization from hexane gave 56 mg of a white solid.

Example 16

3-Butyl-3-ethyl-4,5-dihydroxy-5-phenyl-2,3,4,5-tetrahydro-benzothiepine-1,1-dioxide (20)

This product was isolated along with 6b from hydrogenation of a mixture of 8a and 8b.

**Example 17**

3-Butyl-3-ethyl-4-hydroxy-5-phenylthio-2,3,4,5-tetrahydro-benzothiepine-1,1-dioxide (21)


A mixture of 25 mg (0.085 mmole) of 19b, 0.27 g (2.7 mmole) of thiophenol, 0.37 g (2.7 mmole) of potassium carbonate, and 4 mL of DMF was stirred at room temperature under N_2 for 19 h. The reaction mixture was poured into water and extracted with methylene chloride. The methylene chloride layer was washed successively with 10% NaOH and brine, dried over MgSO_4 , and concentrated in vacuo to give 0.19 g of semisolid which contain substantial amounts of diphenyl disulfide. This material was purified by HPLC (5% EtOAc-hexane) to remove diphenyl disulfide in the first fraction. The column was then eluted with 20% EtOAc-hexane to give 17 mg of a first fraction, 4 mg of a second fraction and 11 mg of a third fraction which were three different isomers of 21, i.e. 21a, 21b, and 21c, respectively, by ^1H NMR and mass spectra.

Example 18

Alternative Synthesis of 6c and 6d

A. Preparation from 2-((2-benzoylphenylthio)methyl)-2-ethylthoxanone (2)

Step 1. 2-((2-benzoylphenylthio)methyl)-2-ethylthoxanone (44)




172

To a solution of 9.0 g (0.025 mole) of compound 2 in 100 ml of methylene chloride was added 14.6 g (0.025 mol) of 50-60% MCPBA portionwise. The reaction mixture was stirred at room temperature for 64 h then was stirred with 200 ml of 1 M potassium carbonate and filtered through Celite. The methylene chloride layer was washed twice with 300 ml of 1 M potassium carbonate, once with 10% sodium hydroxide and once with brine. The insoluble solid formed during washing was removed by filtration through Celite. The methylene chloride solution was dried and concentrated in vacuo to give 9.2 g (95%) of semisolid. A portion (2.6 g) of this solid was purified by HPLC (10% ethyl acetate-hexane) to give 1.9 g of crystals, mp 135-136 °C

15

Step 2. 2-((2-Benzylphenylsulfonyl)methyl)-2-ethylhexanal (45)



A solution of 50 g (0.13 mole) of crude 44 in 250 ml of methylene chloride was divided in two portions and charged to two Fisher-Porter bottles. To each bottle was charged 125 ml of methanol and 5 g of 10% Pd/C. The bottles were pressurized with 70 psi of hydrogen and the reaction mixture was stirred at room temperature for 7 h before being charged with an additional 5 g of 10% Pd/C. The reaction mixture was again hydrogenated with 70 psi of hydrogen for 7 h. This procedure was repeated one more time but only 1 g of Pd/C was charged to the reaction mixture. The combined reaction mixture was filtered and concentrated in vacuo to give 46.8 g of 45 as brown oil.

25


Step 3. (3a,4a,5a) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (6c), and (3a,4b,5b) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (6d)

35

To a solution of 27.3 g (73.4 mmole) of 45 in 300 ml of anhydrous THF cooled to 2 °C with an ice bath was added 9.7 g (73.4 mmole) of 95% potassium t-butoxide. The reaction mixture was stirred for 20 min, quenched with 300 ml of 10% HCl and extracted with methylene chloride. The methylene chloride layer was dried over magnesium sulfate and concentrated in vacuo to give 24.7 g of yellow oil. Purification by HPLC (ethyl acetate-hexane) yielded 9.4 g of recovered 45 in the first fraction, 5.5 g (20%) of 6c in the second fraction and 6.5 g (24%) of 6d in the third fraction.

10

B. Preparation from 2-hydroxydiphenylmethane
Step 1. 2-mercaptodiphenylmethane (46)



To a 500 ml flask was charged 16 g (0.33 mol) of 60% sodium hydride oil dispersion. The sodium hydride was washed twice with 50 ml of hexane. To the reaction flask was charged 100 ml of DMF. To this mixture was added a solution of 55.2 g (0.3 mol) of 2-hydroxydiphenylmethane in 200 ml of DMF in 1 h while temperature was maintained below 30 °C by an ice-water bath. After complete addition of the reagent, the mixture was stirred at room temperature for 30 min then cooled with an ice bath. To the reaction mixture was added 49.4 g (0.4 mole) of dimethyl thiocarbamoyl chloride at once. The ice bath was removed and the reaction mixture was stirred at room temperature for 18 h before being poured into 300 ml of water. The organic was extracted into 500 ml of toluene. The toluene layer was washed successively with 10% sodium hydroxide and brine and was concentrated in vacuo to give 78.6 g of a yellow oil which was 95% pure dimethyl O-2-benzylphenyl thiocarbamate. This oil was heated at 280-300 °C in a kugelrohr pot under house vacuum for 30 min. The residue was kugelrohr distilled at 1 torr (180-280 °C). The distillate (56.3 g) was crystallized from methanol to give 37.3 g (46%) of the rearranged product dimethyl

20


25

30

35


5-2-benzylphenyl thiocarbamate as a yellow solid. A mixture of 57 g (0.21 mole) of this yellow solid, 30 g of potassium hydroxide and 150 ml of methanol was stirred overnight then was concentrated in vacuo. The residue was diluted with 200 ml of water and extracted with ether. The aqueous layer was made acidic with concentrated HCl. The oily suspension was extracted into ether. The ether extract was dried over magnesium sulfate and concentrated in vacuo. The residue was crystallized from hexane to give 37.1 g (88%) of 2-mercaptopdiphenylmethane as a yellow solid.

Step 2. 2-((2-Benzylphenylthio)methyl)-2-ethylhexanal (47)



A mixture of 60 g (0.3 mole) of yellow solid from step 1, 70 g (0.3 mole) of compound 1 from preparation 1, 32.4 g (0.32 mole) of triethylamine, 120 ml of 2-methoxyethyl ether was held at reflux for 6 hr and concentrated in vacuo. The residue was triturated with 500 ml of water and 30 ml of concentrate HCl. The organic was extracted into 400 ml of ether. The ether layer was washed successively with brine, 10% sodium hydroxide and brine and was dried over magnesium sulfate and concentrated in vacuo. The residue (98.3 g) was purified by HPLC with 2-5% ethyl acetate-hexane as eluent to give 2-((2-benzylphenylthio)methyl)-2-ethylhexanal 47 as a yellow syrup.

Step 3. 2-((2-Benzylphenylsulfonyl)methyl)-2-ethylhexanal (45)




To a solution of 72.8 g (0.21 mole) of yellow syrup from step 2 in 1 liter of methylene chloride cooled to 10 °C was added 132 g of 50-60% MCPBA in 40 min. The reaction mixture was stirred for 2 h. An additional 13 g of 50-60% MCPBA was added to the reaction mixture. The reaction mixture was stirred for 2 h and filtered

175

through Celite. The methylene chloride solution was washed twice with 1 liter of 1 M potassium carbonate then with 1 liter of brine. The methylene chloride layer was dried over magnesium sulfate and concentrated to 76 g of 2-((2-benzylphenylsulfonyl)methyl)-2-ethylhexanal 45 as a syrup.

Step 4. (3a, 4a, 5a) 3-Butyl-3-ethyl-6-hydroxy-5-phenyl-2,3,6,5-tetrahydrobenzothiepine-1,1-dioxide (6c), and (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,6,5-tetrahydrobenzothiepine-1,1-dioxide (6d)

Reaction of 45 with potassium t-butoxide according to the procedure in step 3 of procedure A gave pure 6c and 6d after HPLC.



Example 19
(3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-8-methoxy-5-phenyl-2,3,6,5-tetrahydrobenzothiepine-1,1-dioxide (25) and (3a, 4a, 5a) 3-Butyl-3-ethyl-6-hydroxy-8-methoxy-5-phenyl-2,3,6,5-tetrahydrobenzothiepine-1,1-dioxide (26)
Step 1. Preparation of 2-((2-benzoyl-6-methoxyphenylthio)methyl)-2-ethylhexanal (22)

2-Hydroxy-4-methoxybenzophenone was converted to the dimethyl O-2-benzoylphenyl thiocarbamate by methods previously described in example 18. The product can be isolated by recrystallization from ethanol. Using this improved isolation procedure no chromatography was needed. The thermal rearrangement was performed by reacting the thiocarbamate (5 g) in diphenyl ether at 260 °C as previously described. The improved isolation procedure which avoided a chromatography step was described below.

The crude pyrolysis product was then heated at 65 °C in 100 ml of methanol and 100 ml of THF in the presence of 3.5 g of KOH for 4 h. After removing THF and methanol

176


by rotary evaporation the solution was extracted with 5 % NaOH and ether. The base layer was acidified and extracted with ether to obtain a 2.9 g of crude thiophenol product. The product was further purified by titrating the desired mercaptan into base with limited KOH. After acidification and extraction with ether pure 2-mercapto-4-methoxybenzophenone (2.3 g) was isolated.

2-mercapto-4-methoxybenzophenone can readily be converted to the 2-((2-benzoyl-4-methoxyphenylthio)methyl)-2-ethylhexanal (22) by reaction with 2-ethyl-2-(mesyloxymethyl)hexanal (1) as previously described.

Step 2. 2-((2-Benzoyl-5-methoxyphenylsulfonyl)methyl)-2-ethylhexanal (23)




Substrate 22 was readily oxidized to 2-((2-benzoyl-5-methoxyphenyl-sulfonyl)methyl)-2-ethylhexanal (23) as described in example 18.



Step 3. 2-((2-Benzoyl-5-methoxyphenylsulfonyl)methyl)-2-ethylhexanal (24)

Sulfone 23 was then reduced to 2-((2-benzoyl-5-methoxyphenyl-sulfonyl)methyl)-2-ethylhexanal (24) as described in example 18.



Step 4. (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-8-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (25) and (3a, 4a, 5a) 3-Butyl-3-ethyl-4-hydroxy-8-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (26)



A 3-neck flask equipped with a powder addition funnel, thermocouple and nitrogen bubbler was charged with 19.8 g (0.05 mole) of sulfone 24 in 100 ml dry THF. The reaction was cooled to -1.6 °C internal

177

temperature by means of ice/salt bath. Slowly add 5.61 g (0.05 mole) of potassium t-butoxide by means of the powder addition funnel. The resulting light yellow solution was maintained at -1.6 °C. After 30 min reaction 400 ml of cold ether was added and this solution was extracted with cold 10 % HCl. The acid layer was extracted with 300 ml of methylene chloride. The organic layers were combined and dried over magnesium sulfate and after filtration stripped to dryness to obtain 19.9 g of product. ¹H nmr and glpc indicated a 96% conversion to a 50/50 mixture of 25 and 26. The only other observable compound was 4% starting sulfone 24.

The product was then dissolved in 250 ml of 90/10 hexane/ethyl acetate by warming to 50 °C. The solution was allowed to cool to room temperature and in this way pure 26 can be isolated. The crystallization can be enhanced by addition of a seed crystal of 26. After 2 crystallizations the mother liquor which was now 85.4% 25 and has a dry weight of 8.7 g. This material was dissolved in 100 ml of 90/10 hexane/ethyl acetate and 10 ml of pure ethyl acetate at 40 °C. Pure 25 can be isolated by seeding this solution with a seed crystal of 25 after storing it overnight at 0 °C.



Example 20

(3a, 4a, 5a) 3-Butyl-3-ethyl-4,8-dihydroxy-8-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (27)

In a 25 ml round bottomed flask, 1 g of 26 (2.5 mmoles) and 10 ml methylene chloride were cooled to -78 °C with stirring. Next 0.7 ml of boron tribromide (7.5 mmole) was added via syringe. The reaction was allowed to slowly warm to room temperature and stirred for 6 h. The reaction was then diluted with 50 ml methylene chloride and washed with saturated NaCl and then water. The organic layer was dried over magnesium

178

sulfate. The product (0.88g) 27 was characterized by NMR and mass spectra.

Example 21

General Alkylation of phenol 27

A 25 ml flask was charged with 0.15 g of 27 (0.38 mmole), 5 ml anhydrous DMF, 54 mg of potassium carbonate (0.38 mmole) and 140 mg ethyl iodide (0.9 mmole). The reaction was stirred at room temperature overnight. The reaction was diluted with 50 ml ethyl ether and washed with water (25 ml) then 5% NaOH (20 ml) and then sat. NaCl. After stripping off the solvent the ethoxylated product 28 was obtained in high yield. The product was characterized by NMR and mass spectra. This same procedure was used to prepare products listed in table 1 from the corresponding iodides or bromides. For higher boiling alkyl iodides and bromides only one equivalent of the alkyl halide was used.




Table 1

Compound No.	R
27	H
26	Me
28	Et
29	hexyl
30	Ac
31	(CH ₂) ₆ -N-phtalimide

Example 22

(3a, 4a, 5a) 3-Butyl-3-ethyl-6-hydroxy-7-hydroxyamino-5-phenyl-2,3,4,5-tetrahydrobenzothiazepine-1,1-dioxide (37) and (3a, 4b, 5b) 3-Butyl-3-ethyl-6-hydroxy-7-hydroxyamino-5-phenyl-2,3,4,5-tetrahydrobenzothiazepine-1,1-dioxide (38)

Step 1. Preparation of 2-chloro-5-nitrophenylmethane (32)




Procedure adapted from reference: Synthesis -Stuttgart 9 770-772 (1986) Olah G. Et al

Under nitrogen, a 3 neck flask was charged with 45 g (0.172 mole) of 2-chloro-5-nitrobenzophenone in 345 ml methylene chloride and the solution was cooled to ice/water temperature. By means of an additional funnel, 150 g (0.172 mole) of trifluoromethane sulfonic acid in 345 ml methylene chloride was added slowly. Next 30 g of triethylsilane (0.172 mole) in 345 ml methylene chloride was added dropwise to the chilled solution. Both addition steps (trifluoromethane sulfonic acid and triethylsilane) were repeated. After the additions were completed the reaction was allowed to slowly warm up to room temperature and stirred for 12 h under nitrogen. The reaction mixture was then poured into a chilled stirred solution of 1600 ml of saturated sodium bicarbonate. Gas evolution occurred. Poured into a 4 liter separatory funnel and separated layers. The methylene chloride layer was isolated and

combined with two 500 ml methylene chloride extractions of the aqueous layer. The methylene chloride solution was dried over magnesium sulfate and concentrated in vacuo. The residue was recrystallized from hexane to give 39 g product. Structure 32 was confirmed by mass spectra and proton and carbon NMR.

Step 2. Preparation of 2-((2-benzyl-4-ethyl-1-nitrophenoxy)methyl)-2-ethylhexanal (33)



The 2-chloro-5-nitrodiphenylmethane product 32 (40 g, 0.156 mole) from above was placed in a 2 liter 2 neck flask with water condenser. Next 150 ml DMSO and 7.18 g (0.156 mole) of lithium sulfide was added and the


solution was stirred at 75 °C for 12 h. The reaction

was cooled to room temperature and then 51.7 g of mesylate IV was added in 90 ml DMSO. The reaction mixture was heated to 80 °C under nitrogen. After 12 h monitored by TLC and added more mesylate if necessary. Continued the reaction until the reaction was

completed. Next the reaction mixture was slowly poured into a 1900 ml of 5% acetic aqueous solution with stirring, extracted with 4 X 700 ml of ether, and dried over MgSO₄. After removal of ether, 82.7 g of product was isolated. The material can be further purified by silica gel chromatography using 95% hexane and 5% ethyl acetate. If pure mesylate was used in this step there was no need for further purification. The product 33 was characterized by mass spectra and NMR.

Step 3. Oxidation of the nitro product 33 to the sulfone 2-((2-benzyl-4-nitrophenoxy)methyl)-2-ethylhexanal (34)


The procedure used to oxidize the sulfide 33 to the sulfone 34 has been previously described.



Step 4. Reduction of 34 to 2-((2-benzyl-4-hydroxyaminophenoxy)methyl)-2-ethylhexanal (35)


A 15 g sample of 34 was dissolved in 230 ml of ethanol and placed in a 500 ml rb flask under nitrogen. Next 1.5 g of 10 wt.% Pd/C was added and hydrogen gas was bubbled through the solution at room temperature until the nitro substrate 34 was consumed. The reaction could be readily monitored by silica gel TLC using 80/20 hexane/EtOAc. Product 35 was isolated by filtering off the Pd/C and then stripping off the EtOH solvent. The product was characterized by NMR and mass spectra.

Step 5. Preparation of the 2-((2-benzyl-4-N,O-di-(t-butoxy-carbonyl)hydroxyaminophenoxy)methyl)-2-ethylhexanal (36).




A 13.35 g sample of 35 (0.0344 mole) in 40 ml of dry THF was stirred in a 250 ml round bottomed flask. Next added 7.52 g (0.0344 mole) of di-t-butyl dicarbonate in 7 ml THF. Heated at 60 °C overnight. Stripped off THF and redissolved in methylene chloride. Extracted with 1 % HCl; and then 5% sodium bicarbonate.

The product was further purified by column chromatography using 90/10 hexane/ethyl acetate and then 70/30 hexane/ethyl acetate. The product 36 was obtained (4.12 g) which appeared to be mainly the di-(t-butoxycarbonyl) derivatives by proton NMR.




Step 6. (3a,4a,5a) 3-Butyl-3-ethyl-4-hydroxy-7-hydroxyamino-5-phenyl-2,3,4,5-tetrahydrobenzothioepine-1,1-dioxide (37) and (3a,4b,5b) 3-Butyl-3-ethyl-4-hydroxy-7-hydroxyamino-5-phenyl-2,3,4,5-tetrahydrobenzothioepine-1,1-dioxide (38)



A 250ml 3-neck round bottomed flask was charged with 4 g of 36 (6.8 mmoles), and 100 ml of anhydrous THF and cooled to -78 °C under a nitrogen atmosphere. Slowly add 2.29 g potassium tert-butoxide (20.4 mmoles) with

stirring and maintaining a -78 °C reaction temperature. After 1 h at -78 °C the addition of base was completed and the temperature was brought to -10 °C by means of a ice/salt bath. After 3 h at -10 °C, only trace 36 remained by TLC. Next add 35 ml of deionized water to the reaction mixture at -10 °C and stirred for 5 min. Striped off most of the THF and added to separatory funnel and extracted with ether until all of the organic was removed from the water phase. The combined ether phases were washed with saturated NaCl and then dried over sodium sulfate. The only products by TLC and NMR were the two BOC protected isomers of 37 and 38. The isomers were separated by silica gel chromatography using 85% hexane and 15 % ethyl acetate; BOC-37 (0.71 g) and BOC- 38 (0.78 g).

Next the BOC protecting group was removed by reacting 0.87 g of BOC-38 (1.78 mmol) with 8.7 ml of 4 M HCl (34.8 mmol) in dioxane for 30 min. Next added 4.74 g of sodium acetate (34.8 mmol) to the reaction mixture and 16.5 ml ether and stirred until clear. After transferring to a separatory funnel extracted with ether and water and then dried the ether layer with sodium sulfate. After removing the ether, 0.665 g of 38 was isolated. Isomer 37 could be obtained in a similar procedure.



Example 23
(3a,4a,5a) 3-Butyl-3-ethyl-7-(n-hexylamino)-4-hydroxy-5-phenoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (40) and (3a,4b,5b) 3-Butyl-3-ethyl-7-(n-hexylamino)-4-hydroxy-5-phenoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (41)

Step 1. 2-((2-Benzyl-6-(n-hexylamino)phoxysulfonyl)methyl)-2-ethylhexanol (39)

In a Fischer porter bottle weighed out 0.5 g of 36 (1.2 mmol) and dissolved in 3.8 ml of ethanol under

183

nitrogen. Next added 0.1 g of Pd/C and 3.8 ml of hexanal. Seal and pressure to 50 psi of hydrogen gas. Stirred for 48 h. After filtering off the catalyst and removing the solvent by rotary evaporation 39 was isolated by column chromatography (0.16 g) using 90/10 hexane ethyl acetate and gradually increasing the mobile phase to 70/30 hexane/ethyl acetate. The product was characterized by NMR and mass spectra.

Step 2. (3a,4a,5a) 3-Butyl-3-ethyl-7-(n-hexylamino)-4-hydroxy-5-phenoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (40) and (3a,4b,5b) 3-Butyl-3-ethyl-7-(n-hexylamino)-4-hydroxy-5-phenoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (41)



A 2-neck, 25 ml round bottomed flask with stir bar was charged with 0.158 g 39 (0.335 mmol) and 5 ml anhydrous THF under nitrogen. Cool to -10 °C by means of a salt/water bath. Slowly add 0.113 g of potassium tert butoxide (0.335 mmol). After 15 min at -10 °C all of the starting material was consumed by TLC and only the two isomers 40 and 41 were observed. Next added 5 ml of chilled 10% HCl and stirred at -10 °C for 5 min. Transferred to a separatory funnel and extract with ether. Dried over sodium sulfate. Proton NMR of the dried product (0.143 g) indicated only the presence of the two isomers 40 and 41. The two isomers were separated by silica gel chromatography using 90/10 hexane ethyl acetate and gradually increasing the mobile phase to 70/30 hexane/ethyl acetate. 40 (53.2 mg); 41(58.9 mg).

Example 24


Quaternization of amine substrates 40 and 41



Amine products such as 40 and 41 can be readily alkylated to quaternary salts by reaction with alkyl halides. For example 40 in DMF with 5 equivalents of


184

methyl iodide in the presence of 2,6 dimethyl lutidine produces the dimethylhexylamino quaternary salt.

Example 25

(3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-5-(4-iodophenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (42)

In a 25 ml round bottomed flask 0.5 g (1.3 mmole) of **6d**, 0.67 g of mercuric triflate were dissolved in 20 ml of dry methylene chloride with stirring. Next 0.34 g of Iodine was added and the solution was stirred at room temperature for 30 h. The reaction was then diluted with 50 ml methylene chloride and washed with 10 ml of 1 M sodium thiosulfate; 10 ml of saturated KI; and dried over sodium sulfate. See Tetrahedron, Vol.50, No. 17, pp 5139-5146 (1994) Bachki, F. Et al. Mass spectrum indicated a mixture of **6d**, mono iodide **42** and a diiodide adduct. The mixture was separated by column chromatography and **42** was characterized by NMR and mass spectra.

Example 26

(3a, 4b, 5b) 3-Butyl-5-(4-carbomethoxyphenyl)-3-ethyl-4-hydroxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (43)

A 0.1 g sample of **42** (0.212 mmole), 2.5 ml dry methanol, 38 μ l triethylamine (0.275 mmole), 0.3 ml toluene and 37 mg of palladium chloride (0.21 mmole) was charged to a glass lined mini reactor at 300 psi carbon monoxide. The reaction was heated at 100 °C overnight. The catalyst was filtered and a high yield of product was isolated.


The product was characterized by NMR and mass spectra.

Note the ester functionalized product **43** can be converted to the free acid by hydrolysis.

185

Example 27

(3a, 4a, 5a) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (48), and (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (49)



Step 1. 2-Mercapto-5-methoxybenzophenone (50)


Reaction of 66.2 g of 4-methoxythiophenol with 360 ml of 2.5 N n-butyllithium, 105 g of tetramethylethylenediamine and 66.7 g of benzonitrile in 600 ml cyclohexane according to the procedure in WO 93/16055 gave 73.2 g of brown oil which was kugelrohr distilled to remove 4-methoxythiophenol and gave 43.86 g of crude **50** in the pot residue.

Step 2. 2-((2-Benzoyl-4-methoxyphenylthio)methyl)-2-ethylhexanal (51)



Reaction of 10 g (0.04 mole) of crude **50** with 4.8 g (0.02 mole) of mesylate **1** and 3.2 ml (0.23 mole) of triethylamine in 50 ml of diglyme according to the procedure for the preparation of **2** gave 10.5 g of crude product which was purified by HPLC (5% ethyl acetate-hexane) to give 1.7 g (22%) of **51**.

Step 3. 2-((2-Benzoyl-4-methoxyphenylsulfonyl)methyl)-2-ethyl-hexanal (52)



A solution of 1.2 g (3.1 mmole) of **51** in 25 ml of methylene chloride was reacted with 2.0 g (6.2 mmole) of 50-60% MCPBA according to the procedure of step 2 of procedure A in example 18 gave 1.16 g (90%) of **52** as a yellow oil.


Step 4. 2-((2-Benzyl-4-methoxyphenylsulfonyl)methyl)-2-ethylhexanal (53)




186

Hydrogenation of 1.1 g of 52 according to the procedure of step 3 of procedure A of example 18 gave 53 as a yellow oil (1.1 g).

Step 5. (3a,4a,5a) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (48), and (3a,4b,5b) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (49)



A solution of 1.1 g of 53, 0.36 g of potassium t-butoxide and 25 ml of anhydrous THF was held at reflux for 2 h and worked up as in step 4 of procedure A of example 18 to give 1.07 g of a crude product which was purified by HPLC to give 40 mg (48) of 48 as crystals, mp 153-154 °C and 90 mg (88) of 49 as solid, mp 136-140 °C.




Example 28
5-Phenyl-2,3-dihydrospirobenzothiepine-3,1'-cyclohexane (57)


Step 1. 1-(Hydroxymethyl)-cyclohexanecarboxaldehyde (54)

To a cold (0 °C) mixture of 100 g (0.891 mole) of cyclohexanecarboxaldehyde, 76.5 g of 37% of formaldehyde in 225 ml of methanol was added dropwise 90 ml of 1 N Sodium hydroxide in 1 h. The reaction mixture was stirred at room temperature over 48 then was evaporated to remove methanol. The reaction mixture was diluted with water and extracted with methylene chloride. The organic layer was washed with water, brine, and dried over sodium sulfate and concentrated under vacuum to give 75 g (59.7%) of thick oil. Proton NMR and mass spectra were consistent with the product.

Step 2. 1-(mesyloxymethyl)-cyclohexanecarboxaldehyde (55)




To a cold (0 °C) mixture of alcohol 54 (75 g, 0.54 mole) and 65.29 g (0.57 mole) of methanesulfonyl chloride in 80 ml of methylene chloride was added a solution of pyridine (47.96 g, 0.57 mole) in 40 ml of methylene chloride. The reaction mixture was stirred at room temperature for 18 h then quenched with water, acidified with conc. HCl and extracted with methylene chloride. The organic layer was washed with water, brine, and dried over sodium sulfate and concentrated under vacuum to give 91.63 g (77.8%) of thick oil. Proton NMR and mass spectra were consistent with the product.



Step 3. 1-(2-benzoylphenylthio)methyl)-cyclohexanecarboxaldehyde (56)

A mixture of 69 g (0.303 mole) of 2-mercaptobenzophenone, 82 g (0.303 mole) of mesylate 55, 32 g of triethylamine, and 150 ml of diglyme was stirred and held at reflux for 24 h. The mixture was cooled, poured into dil. HCl and extracted with methylene chloride. The organic layer was washed with 10% NaOH, water, brine, and dried over sodium sulfate and concentrated under vacuum to remove excess diglyme. This was purified by silica gel flush column (5% EtOAc: Hexane) and gave 18.6 g (75.9%) of yellow oil. Proton NMR and mass spectra were consistent with the product.

Step 4. 5-Phenyl-2,3-dihydrospirobenzothiepine-3,1'-cyclohexane (57)



To a mixture of 6.19 g of zinc dust and 100 ml of dry DME was added TiCl₄ (16.8 g, 0.108 mole). The reaction mixture was heated to reflux for 2 h. A solution of compound 56 (8.3 g, 0.023 mole) in 50 ml of DME was added dropwise to the reaction mixture in 1 h and the mixture was held at reflux for 18 h. The mixture was

cooled, poured into water and extracted with ether. The organic layer was washed with water, brine, and dried over sodium sulfate, filtered through celite and concentrated under vacuum. The residue was purified by HPLC (10% EtOAc: Hexane) to give 4.6 g (64%) of white solid, mp 90-91 °C. Proton and carbon NMR and mass spectra were consistent with the product.

Example 29

8b-Phenyl-1a,2,3,8b-tetrahydrospiro(benzothiepine)(4,5-b)oxirane-2,1'-cyclohexane)-4,4-dioxido (58)




(58)

To a solution of 57 (4.6 g, 15 mmole) in 50 ml chloroform under nitrogen was added 55% MCPBA (16.5 g, 52.6 mmole) portionwise with spatula. The reaction was held at reflux for 18 h and washed with 10% NaOH(3X), water, brine, and dried over sodium sulfate and concentrated under vacuum to give 5 g of crude product. This was recrystallized from Hexane/EtOAc to give 4.31 g (81%) of yellow solid, mp 154-155 °C. Proton and carbon NMR and mass spectra were consistent with the product.

Example 30

trans-4-Hydroxy-5-phenyl-2,3,4,5-tetrahydro spiro(benzothiepine-3,1'-cyclohexane)-1,1-dioxide (59)




(59)

A mixture of 0.5 g (1.4 mmole) of 58, 20 ml of ethanol, 10 ml of methylene chloride and 0.4 g of 10% Pd/C catalyst was hydrogenated with 70 psi hydrogen for 3 h at room temperature. The crude reaction slurry was filtered through Celite and evaporated to dryness. The residue was purified by HPLC (10% EtOAc:Hexane, 25% EtOAc:Hexane). The first fraction was 300 mg (60%) as a white solid, mp 99-100 °C. Proton NMR showed this was a trans isomer. The second fraction gave 200 mg of solid which was impure cis isomer.

Example 31

cis-4-Hydroxy-5-phenyl-2,3,4,5-tetrahydro spiro(benzothiepine-3,1'-cyclohexane)-1,1-dioxide (60)



(60)

To a solution of 0.2 g (0.56 mmole) of 59 in 20 ml of CH₂Cl₂, was added 8 g of 50% NaOH and one drop of Aliquat-336 (methyltriprilylammonium chloride) phase transfer catalyst. The reaction mixture was stirred for 10 h at room temperature. Twenty g of ice was added to the mixture and the mixture was extracted with CH₂Cl₂, (3x10 ml) washed with water, brine and dried over MgSO₄, and concentrated in vacuo to recover 0.15 g of crude product. This was recrystallized from Hexane/EtOAc to give 125 mg of white crystal, mp 209-210 °C. Proton and carbon NMR and mass spectra were consistent with the product.

Example 32


(3a,4a,5a) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine (61), and (3a,4b,5b) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine (62)



(61)

(62)

To a solution of 0.5 g (1.47 mmole) of compound 47 in 5 ml of anhydrous THF was added 0.17 g (1.47 mmole) of 95% potassium t-butoxide. The reaction mixture was stirred at room temperature for 18 h and quenched with 10 ml of 10% HCl. The organic was extracted into methylene chloride. The methylene chloride extract was dried over magnesium sulfate and concentrated in vacuo. The residue was purified by HPLC (2% EtOAc:hexane) to give 47 mg of 61 in the second fraction and 38 mg of 62 in the third fraction. Proton NMR and mass spectra were consistent with the assigned structures.

Example 33

(63)

(64)

(3a, 4a, 5a) 3-Butyl-3-ethyl-4-hydroxy-7-amino-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (63) and (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-7-amino-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (64)

An autoclave was charged with 200 mg of 37 in 40 cc ethanol and .02 g 10 & Pd/C. After purging with nitrogen the clave was charged with 100 psi hydrogen and heated to 55 C. The reaction was monitored by TLC and mass spec and allowed to proceed until all of 37 was consumed. After the reaction was complete the catalyst was filtered and the solvent was removed in vacuo and the only observable product was amine 63. This same procedure was used to produce 64 from 38.


Example 34

(3a, 4a, 5a) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-(3'-methoxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (65), and (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-(3'-methoxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (66).

Alkylation of e-methoxyphenol with 3-methoxybenzyl chloride according to the procedure described in J. Chem. Soc. 2431 (1958) gave 4-methoxy-2-(3'-methoxybenzyl)phenol in 35% yield. This material was converted to compound 65, mp 138.5-141.5 °C, and compound 66, mp 115.5-117.5 °C, by the procedure similar to that in Example 18 method B.

Example 35

(3a, 4a, 5a) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-(3'-(trifluoromethyl)phenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (67), and (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-(3'-(trifluoromethyl)phenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (68).



Alkylation of 4-methoxyphenol with 3-(trifluoromethyl)benzyl chloride according to the procedure described in J. Chem. Soc. 2431 (1958) gave 4-methoxy-2-(3'-(trifluoromethyl)benzyl)phenol. This material was converted to compound 67, mp 226.5-228 °C, and compound 68, mp 188-190 °C, by the procedure similar to that in Example 18 method B.

Example 36

(3a, 4a, 5a) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (69), and (3a, 4b, 5b) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (70).

Alkylation of 4-methoxyphenol with 4-fluorobenzyl chloride according to the procedure described in J. Chem. Soc. 2431 (1958) gave 4-methoxy-2-(4'-fluorobenzyl)phenol. This material was converted to compound 69 and compound 70 by the procedure similar to that in Example 18 method B.


Example 37

(3a, 4a, 5a) 3-Butyl-3-ethyl-5-(3'-fluorophenyl)-4-hydroxy-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (71), and (3a, 4b, 5b) 3-Butyl-3-ethyl-5-(3'-fluorophenyl)-4-hydroxy-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (72).

Alkylation of 4-methoxyphenol with 3-fluorobenzyl chloride according to the procedure described in J. Chem. Soc. 2431 (1958) gave 4-methoxy-2-(3'-fluorobenzyl)phenol. This material was converted to compound 71 and compound 72 by the procedure similar to that in Example 18 method B.

Example 38

(3a, 4a, 5a) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-(3'-(trifluoromethyl)phenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (73), and (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-(3'-(trifluoromethyl)phenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxido (74).



(3a, 4a, 5a) 3-Butyl-3-ethyl-5-(2'-fluorophenyl)-4-hydroxy-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (73), and (3a, 4b, 5b) 3-Butyl-3-ethyl-5-(2'-fluorophenyl)-4-hydroxy-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (74).

Alkylation of 4-methoxyphenol with 2-fluorobenzyl chloride according to the procedure described in J. Chem. Soc. 2431 (1958) gave 4-methoxy-2-(2'-fluorobenzyl)phenol. This material was converted to compound 73 and compound 74 by the procedure similar to that in Example 18 method B.

Example 39

(3a, 4a, 5a) 3-Butyl-7-bromo-3-ethyl-4-hydroxy-5-(3'-methoxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (75), and (3a, 4b, 5b) 3-Butyl-7-bromo-3-ethyl-4-hydroxy-5-(3'-methoxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (76).

Alkylation of 4-bromophenol with 3-methoxybenzyl chloride according to the procedure described in J. Chem. Soc. 2431 (1958) gave 4-bromo-2-(3'-methoxybenzyl)phenol. This material was converted to compound 75, mp 97-101.5 °C, and compound 76, mp 102-106 °C, by the procedure similar to that in Example 18 method B.

Example 40

(3a, 4a, 5a) 3-Butyl-3-ethyl-7-fluoro-5-(4'-fluorophenyl)-4-hydroxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (77), and (3a, 4b, 5b) 3-Butyl-3-ethyl-7-fluoro-5-(4'-fluorophenyl)-4-hydroxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (78).

Alkylation of 4-fluorophenol with 4-fluorobenzyl chloride according to the procedure described in J.

193


Chem. Soc. 2431 (1958) gave 4-fluoro-2-(4'-fluorobenzyl)phenol. This material was converted to compound 77, mp 228-230 °C, and compound 78, mp 134.5-139 °C, by the procedure similar to that in Example 18 method B.

Example 41

(3a, 4a, 5a) 3-Butyl-3-ethyl-7-fluoro-4-hydroxy-5-(3'-methoxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (79), and (3a, 4b, 5b) 3-Butyl-3-ethyl-7-fluoro-4-hydroxy-5-(3'-methoxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (80).

Alkylation of 4-fluorophenol with 3-methoxybenzyl chloride according to the procedure described in J.

Chem. Soc. 2431 (1958) gave 4-fluoro-2-(3'-methoxybenzyl)phenol. This material was converted to compound 79, as a solid and compound 80, mp 153-155 °C, by the procedure similar to that in Example 18 method B.




Example 42

(3a, 4b, 5b) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-methylthio-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (81).


A mixture of 0.68 (1.66 mmol) of compound 77, 0.2 g (5 mmol) of sodium methanethiolate and 15 ml of anhydrous DMF was stirred at room temperature for 16 days. The reaction mixture was dilute with ether and washed with water and brine and dried over $MgSO_4$. The ether solution was concentrated in vacuo. The residue was purified by HPLC (20% ethyl acetate in hexanes). The first fraction was impure (3a, 4a, 5a) 3-butyl-3-ethyl-4-hydroxy-7-methylthio-5-(4'-fluorophenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide. The second fraction was compound 81, mp 185-186.5 °C.

194



Example 43
(3a, 4b, 5b) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-(1-pyrrolidinyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (82).

A mixture of 0.53 g (1.30 mmol) of compound 78 and 5 ml of pyrrolidine was held at reflux for 1 h. The reaction mixture was diluted with ether and washed with water and brine and dried over $MgSO_4$. The ether solution was concentrated in vacuo. The residue was crystallized from ether-hexanes to give compound 82, mp 174.5-177 °C.



Example 44
(3a, 4b, 5b) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-(1-morpholinyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (83).

A mixture of 0.4 g (0.98 mmol) of compound 78 and 5.0 g (56 mmol) of morpholine was held at reflux for 2 h and concentrated in vacuo. The residue was diluted with ether (30 ml) and washed with water and brine and dried over $MgSO_4$. The ether solution was concentrated in vacuo. The residue was recrystallized from ether-hexanes to give compound 83, mp 176.5-187.5 °C.




Example 45

(3a, 4a, 5a) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-methyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (84), and (3a, 4b, 5b) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-methyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (85).

Alkylation of 4-methylphenol with 4-fluorobenzyl chloride according to the procedure described in J. Chem. Soc. 2431 (1958) gave 4-methyl-2-(4'-fluorobenzyl)phenol. This material was converted to

195

compound 84 and compound 85 by the procedure similar to that in Example 18, method B.



Example 46
(3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-5-(4'-fluorophenyl)-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (86), and (3a, 4b, 5b) 3-Butyl-3-ethyl-4,7-dihydroxy-5-(4'-fluorophenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (87).

To a solution of 0.52 (1.2 mmol) of compound 66 in 20 ml of methylene chloride was added 1.7 g (6.78 mmol) of boron tribromide. The reaction mixture was cooled to -78 °C and was stirred for 4 min. An additional 0.3 ml of boron tribromide was added to the reaction mixture and the reaction mixture was stirred at -78 °C for 1 h and quenched with 2 N HCl. The organic was extracted into ether. The ether layer was washed with brine, dried over $MgSO_4$, and concentrated in vacuo. The residue (0.48 g) was purified by HPLC (30% ethyl acetate in hexanes). The first fraction was 0.11 g of compound 86 as a white solid, mp 171.5-173 °C. The second fraction was crystallized from chloroform to give 0.04 g of compound 87 as a white solid, mp 264 °C (dec).




Example 47

(3a, 4b, 5b) 3-Butyl-3-ethyl-4,7-dihydroxy-5-(4'-fluorophenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (88).

Reaction of compound 70 with excess boron tribromide at room temperature and worked up as in Example 46 gave compound 88 after an HPLC purification.


Example 48



196

(3a, 4b, 5b) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-(1-azetidinyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (89).

A mixture of 0.20 g (0.49 mmol) of compound 78, and 2.0 g (35 mmol) of aztidine was held at reflux for 3 h and concentrated in vacuo. The residue was diluted with ether (30 ml) and washed with water and brine and dried over MgSO₄. The ether solution was concentrated on a steam bath. The separated crystals were filtered to give 0.136 g of 89 as prisms, mp 196.5-199.5 °C.

Example 49  (91)
(3a, 4a, 5a) 3-Butyl-3-ethyl-5-(3'-methoxyphenyl)-4-hydroxy-7-methylthio-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (90). (3a, 4b, 5b) 3-Butyl-3-ethyl-5-(3'-methoxyphenyl)-4-hydroxy-7-methylthio-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (91).

A mixture of 0.4 g (0.95 mmol) of compound 79, 0.08 g (1.14 mmol) of sodium methanethiolate and 15 ml of anhydrous DMF was stirred at 60 °C for 2 h. An additional 1.4 mmol of sodium methanethiolate was added to the reaction mixture and the mixture was stirred at 60 °C for an additional 2 h. The reaction mixture was triturated with 100 ml of water and extracted with methylene chloride. The methylene chloride water mixture was filtered through Celite and the methylene chloride layer was dried over MgSO₄ and concentrated in vacuo. The first fraction (0.1 g) was compound 90, mp 117-121 °C. The second fraction (0.16 g) was compound 91, mp 68-76 °C.

A mixture of 2.6 g (0.04 mole) of zinc dust, 7.2 g (0.047 mole) of TiCl₄, and 80 mL of anhydrous ethylene glycol dimethyl ether (DME) was held at reflux for 2 h. The reaction mixture was cooled to 5 °C. To the reaction mixture was added dropwise a solution of 3.54 g (0.01 mole) of 2 in 30 mL of DME in 40 min. The reaction mixture was stirred at room temperature for 16 h and then was held at reflux for 2 h and cooled before being poured into brine. The organic was extract into methylene chloride. The methylene chloride extract was dried over MgSO₄ and concentrated in vacuo. The residue was purified by HPLC (hexane) to give 1.7 g (43%) of 3 as an oil in the first fraction. The second fraction was discarded and the third fraction was further purified by HPLC (hexane) to give 0.07 g (2%) of 4a in the earlier fraction and 0.1 g (3%) of 4b in the later fraction.

Example 2

cis-3-Butyl-3-ethyl-5-phenyl-2,3-dihydrobenzothiepine-5H)-4-one-1,1-dioxide (5a) and *trans*-3-Butyl-3-ethyl-5-phenyl-2,3-dihydro-benzothiepin-(5H)-4-one-1,1-dioxide (5b)

To a solution of 1.2 g (3.5 mmole) of 50-60% MCPBA in 20 mL of methylene chloride was added 0.59 g (1.75 mmole) of a mixture of 4a and 4b in 10 mL of methylene chloride. The reaction mixture was stirred for 20 h. An additional 1.2 g (1.75 mmole) of 50-60% MAPBA was added and the reaction mixture was stirred for an additional 3 h then was triturated with 50 mL of 10% NaOH. The insoluble solid was filtered. The methylene chloride layer of the filtrate was washed with brine, dried over MgSO₄, and concentrated in vacuo. The residual syrup was purified by HPLC (5% EtOAc-hexane) to give 0.2 g (30%) of 5a as an oil in the first fraction and 0.17 g (26%) of 5b as an oil in the second fraction.

Example 3

(3a, 4a, 5b) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (6a), (3a, 4b, 5a) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (6b), (3a, 4a, 5a) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (6c), and (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (6d)

A. Reduction of 5a and 5b with Sodium Borohydride

To a solution of 0.22 g (0.59 mmole) of 5b in 10 mL of ethanol was added 0.24 g (6.4 mmole) of sodium borohydride. The reaction mixture was stirred at room temperature for 18 h and concentrated in vacuo to remove ethanol. The residue was triturated with water and extracted with methylene chloride. The methylene chloride extract was dried over MgSO_4 and concentrated in vacuo to give 0.2 g of syrup. In a separate experiment, 0.45 g of 5a was treated with 0.44 g of sodium borohydride in 10 mL of ethanol and was worked up as described above to give 0.5 g of syrup which was identical to the 0.2 g of syrup obtained above. These two materials were combined and purified by HPLC using 10% EtOAc-hexane as eluant. The first fraction was 0.18 g (27%) of 6a as a syrup. The second fraction was 0.2 g (30%) of 6b also as a syrup. The column was then eluted with 20% EtOAc-hexane to give 0.077 g (11%) of 6c in the third fraction as a solid. Recrystallization from hexane gave a solid, mp 179-181 °C. Finally, the column was eluted with 30% EtOAc-hexane to give 0.08 g (12%) of 6d in the fourth fraction as a solid. Recrystallization from hexane gave a solid, mp 160-161 °C.

B. Conversion of 6a to 6c and 6d with NaOH and PTC

To a solution of 0.29 g (0.78 mmole) of 6a in 10 mL CH_2Cl_2 , was added 9 g of 40% NaOH. The reaction mixture was stirred for 0.5 h at room temperature and was added one drop of Aliquat-336 (methyltriisopropylammonium chloride) phase transfer catalyst (PTC). The mixture was stirred for 0.5 h at room temperature before being treated with 25 mL of ice-crystals then was extracted with CH_2Cl_2 (3x10 mL), dried over MgSO_4 and concentrated in vacuo to recover 0.17 g of a colorless film. The components of this mixture were separated using an HPLC and eluted with EtOAc-hexane to give 12.8 mg (4%) of 2-(2-benzylphenylsulfonylethyl)-2-ethylhexenal in the first fraction, 30.9 mg (11%) of 6c in the second fraction and 90.0 mg (31%) of 6d in the third fraction.

Oxidation of 6a to 5b

To a solution of 0.20 g (0.52 mmole) of 6a in 5 mL of CH_2Cl_2 , was added 0.23 g (1.0 mmole) of pyridinium chlorochromate. The reaction mixture was stirred for 2 h then was treated with additional 0.23 g of pyridinium chlorochromate and stirred overnight. The dark reaction mixture was poured into a ceramic filterfrit containing silica gel and was eluted with CH_2Cl_2 . The filtrate was concentrated in vacuo to recover 167 mg (87%) of 5b as a colorless oil.

Example 4

3-Butyl-3-ethyl-5-phenyl-2,3-dihydrobenzothiepine-1,1-dioxide (7)

To a solution of 5.13 g (15.9 mmole) of 3 in 50 mL of CH₂Cl₂ was added 10 g (31.9 mmole) of 50-60% MCPBA (m-chloroperoxybenzoic acid) portionwise causing a mild reflux and formation of a white solid. The reaction mixture was allowed to stir overnight under N₂ and was triturated with 25 mL of water followed by 50 mL of 10% NaOH solution. The organic was extracted into CH₂Cl₂ (4x20 mL). The CH₂Cl₂ extract was dried over MgSO₄ and evaporated to dryness to recover 4.9 g (87%) of an opaque viscous oil.

Example 5

(1a, 2b, 8ba) 2-Butyl-2-ethyl-8b-phenyl-1a, 2, 3, 8b-tetrahydro-benzothiepine[4, 5-b]oxirene-4, 4-dioxide (8a)
(1a, 2a, 8ba) 2-Butyl-2-ethyl-8b-phenyl-1a, 2, 3, 8b-tetrahydro-benzothiepine [4, 5-b]oxirene-4, 4-dioxide (8b)

To 1.3 g (4.03 mole) of 3 in 25 mL of CHCl₃ was added portionwise 5 g (14.1 mmole) of 50-60 % MCPBA causing a mild exotherm. The reaction mixture was stirred under N₂ overnight and was then held at reflux for 3 h. The insoluble white slurry was filtered. The filtrate was extracted with 10% potassium carbonate (3x50 mL), once with brine, dried over MgSO₄, and concentrated in vacuo to give 1.37 g of a light yellow oil. Purification by HPLC gave 0.65 g of crystalline product. This product is a mixture of two isomers. Trituration of this crystalline product in hexane recovered 141.7 mg (10%) of a white crystalline product. This isomer was characterized by NMR and mass spectra to be the (1a, 2b, 8ba) isomer 8a. The hexane filtrate was concentrated in vacuo to give 206 mg of white film which is a mixture of 30% 8a and 70% 8b by ¹H NMR.

201

Example 6

cis-3-Butyl-3-ethyl-5-phenyl-2,3,4,5-tetrahydro-benzothiepine-1,1-dioxide (9a), trans-3-Butyl-3-ethyl-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (9b), and 3-Butyl-3-ethyl-4-hydroxy-5-cyclohexylidene-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (10)

A mixture of 0.15 g (0.4 mmole) of a 3:7 mixture of 8a and 8b was dissolved in 15 mL MeOH in a 3 oz. Fisher/Porter vessel, then was added 0.1 g of 10% Pd/C catalyst. This mixture was hydrogenated at 70 psi H₂ for 5 h and filtered. The filtrate was evaporated to dryness in vacuo to recover 0.117 g of a colorless oil. This material was purified by HPLC eluting with EtOAc-hexane. The first fraction was 4.2 mg (3%) of 9b. The second fraction, 5.0 mg (4%), was a 50/50 mixture of 9a and 9b. The third fraction was 8.8 mg (6%) of 6a. The fourth fraction was 25.5 mg (18%) of 6b. The fifth fraction was 9.6 mg (7%) of a mixture of 6b and a product believed to be 3-butyl-3-ethyl-4,5-dihydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide based on mass spectrum. The sixth fraction was 7.5 mg (5%) of a mixture of 6a and one of the isomers of 10, 10a.

Example 7

In another experiment, a product (3.7 g) from epoxidation of 3 with excess MCPBA in refluxing CHCl₃, under air was hydrogenated in 100 mL of methanol using 1 g of 10% Pd/C catalyst and 70 psi hydrogen. The product was purified by HPLC to give 0.9 g (25%) of 9b, 0.45 g (13%) of 9a, 0.27 g (7%) of 6a, 0.51 g (14%) of 6b, 0.02 g (1%) of 6c, 0.06 g (2%) of one isomer of 10, 10a and 0.03 g (1%) of another isomer of 10, 10b.

202

Example 82-(2-Benzoylphenylthio)methyl)butyraldehyde (11)

To an ice bath cooled solution of 9.76 g (0.116 mole) of 2-ethylacrolein in 40 mL of dry THF was added 24.6 g (0.116 mole) of 2-mercaptobenzophenone in 40 mL of THF followed by 13 g (0.128 mole) of triethylamine. The reaction mixture was stirred at room temperature for 3 days, diluted with ether, and was washed successively with dilute HCl, brine, and 1 M potassium carbonate. The ether layer was dried over MgSO₄ and concentrated in vacuo. The residue was purified by HPLC (10% EtOAc-hexane) to give 22 g (64%) of 11 in the second fraction. An attempt to further purify this material by Kugelrohr distillation at 0.5 torr (160-190 °C) gave a fraction (12.2 g) which contained starting material indicating a reversed reaction during distillation. This material was dissolved in ether (100 mL) and was washed with 50 mL of 1 M potassium carbonate three times to give 6.0 g of a syrup which was purified by HPLC (10% EtOAc-hexane) to give 5.6 g of pure 11.

Example 93-Ethyl-5-phenyl-2,3-dihydrobenzothiepine (12)

To a mixture of 2.61 g (0.04 mole) of zinc dust and 60 mL of DME was added 7.5 g (0.048 mole) of TiCl₄. The reaction mixture was held at reflux for 2 h. A solution of 2.98 g (0.01 mole) of 11 was added dropwise in 1 h. The reaction mixture was held at reflux for 18 h, cooled and poured into water. The organic was extracted into ether. The ether layer was washed with brine and filtered through Celite. The filtrate was dried over MgSO₄ and concentrated. The residual oil (2.5 g) was purified by HPLC to give 2.06 g (77%) of 12 as an oil in the second fraction.

203

Example 101a,2a,8a)-2-Ethyl-8b-phenyl-1a,2,3,8b-tetrahydro-benzothiepine-[4,5-b]oxirano-4,4-dioxide (13)

To a solution of 1.5 g (5.64 mmole) of 12 in 25 mL of CHCl₃ was added 6.8 g (19.4 mmole) of 50-60% MCPB portionwise causing an exotherm and formation of a white solid. The mixture was stirred at room temperature overnight diluted with 100 mL methylene chloride and washed successively with 10% K₂CO₃ (4x50 mL), water (twice with 25 mL) and brine. The organic layer was then dried over MgSO₄ and evaporated to dryness to recover 1.47 g of an off white solid. ¹H NMR indicated that only one isomer is present. This solid was slurried in 200 mL of warm Et₂O and filtered to give 0.82 g (46%) of 13 as a white solid, mp 185-186.5 °C.

Example 11

(3a,4b,5a)-3-Ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydro-benzothiepine-1,1-dioxide (14a), (3a,4b,5b)-3-Ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (14b), and cis-3-Ethyl-5-phenyl-2,3,4,5-tetrahydro-benzothiepine-1,1-dioxide (15)

A mixture of 0.5 g (1.6 mole) of 13, 50 mL of acetic acid and 0.5 g of 10% Pd/C catalyst was hydrogenated with 70 psi hydrogen for 4 h. The crude reaction slurry was filtered and the filtrate was stirred with 150 mL of a saturated NaHCO₃ solution followed by 89 g of NaHCO₃ powder portionwise to neutralize the rest of acetic acid. The mixture was extracted with methylene chloride (4x25 mL), then the organic layer was dried over MgSO₄ and concentrated in vacuo to give 0.44 g (87%) of a voluminous white solid which was purified by HPLC (EtOAc-Hexane) to give 26.8 mg (6%) of 15 in the first fraction, 272 mg (54%) of 14a as a solid, mp 142-143.5 °C, in the second fraction, and 35 mg (7%) of impure 14b in the third fraction.

204

Example 12**2-Ethyl-2-((2-Hydroxymethylphenyl)chloromethyl)hexenal (16)**

A mixture of 5.0 g (0.036 mole) of 2-mercaptobenzyl alcohol, 6.4 g (0.032 mole) of 1, 3, 6 g (0.036 mole) of triethylamine and 25 mL of 2-methoxyethyl ether was held at reflux for 7 h. Additional 1.1 g of mercaptobenzyl alcohol and 0.72 g of triethylamine was added to the reaction mixture and the mixture was held at reflux for additional 16 h. The reaction mixture was cooled and poured into 6N HCl and extracted with methylene chloride. The methylene chloride extract was washed twice with 10% NaOH, dried over MgSO₄ and concentrated in vacuo to give 9.6 g of residue. Purification by HPLC (20% EtOAc-hexane) gave 3.7 g (41%) of 16 as an oil.

Example 13**2-Ethyl-2-((2-formylphenyl)chloromethyl)hexenal (17)**

A mixture of 3.7 g of 16, 5.6 g (0.026 mole) of pyridinium chlorochromate, 2 g of Celite and 30 mL of methylene chloride was stirred for 18 h and filtered through a bed of silica gel. The silica gel was eluted with methylene chloride. The combined methylene chloride eluant was purified by HPLC (20% EtOAc-hexane) to give 2.4 g (66%) of an oil.

Example 14**3-Butyl-3-ethyl-2,3-dihydrobenzothiepine (18)**

A mixture of 2.6 g (0.04 mole) of zinc dust, 7.2 g (0.047 mole) of TiCl₄, and 50 mL of DME was held at reflux for 2 h and cooled to room temperature. To this mixture was added 2.4 g (8.6 mmole) of 17 in 20 mL of DME in 10 min. The reaction mixture was stirred at room

265

temperature for 2 h and held at reflux for 1 h then was let standing at room temperature over weekend. The reaction mixture was poured into dilute HCl and was stirred with methylene chloride. The methylene chloride-water mixture was filtered through Celite. The methylene chloride layer was washed with brine, dried over MgSO₄, and concentrated in vacuo to give 3.0 g of a residue. Purification by HPLC gave 0.41 g (20%) of 18 as an oil in the early fraction.

Example 15**(1aa,2a,8ba) 2-Butyl-2-ethyl-1a,2,3,8b-tetrahydro-benzothiepine(4,5-b)oxirene-4,4-dioxide (19a) and (1aa,2b,8ba) 2-Butyl-2-ethyl-1a,2,3,8b-tetrahydro-benzothiepine(4,5-b)oxirene-4,4-dioxide (19b)**

To a solution of 0.4 g of 0.4 g (1.6 mmole) of 18 in 30 mL of methylene chloride was added 2.2 g (3.2 mmole) of 50-60% MCPBA. The reaction mixture was stirred for 2 h and concentrated in vacuo. The residue was dissolved in 30 mL of CHCl₃ and was held at reflux for 18 h under N₂. The reaction mixture was stirred with 100 mL of 10% NaOH and 5 g of sodium sulfite. The methylene chloride layer was washed with brine, dried over MgSO₄ and concentrated in vacuo. The residue was purified by HPLC (20% EtOAc-hexane) to give a third fraction which was further purified by HPLC (10% EtOAc-hexane) to give 0.12 g of syrup in the first fraction.

Recrystallization from hexane gave 0.08 g (17%) of 19a, mp 89.5-105.5 °C. The mother liquor from the first fraction was combined with the second fraction and was further purified by HPLC to give additional 19a in the first fraction and 60 mg of 19b in the second fraction. Crystallization from hexane gave 56 mg of a white solid.

Example 16

266

3-Butyl-3-ethyl-6,5-dihydroxy-5-phenyl-2,3,6,5-tetrahydro-benzothiepine-1,1-dioxide (20)

This product was isolated along with 8a from hydrogenation of a mixture of 8a and 8b.

Example 17

3-Butyl-3-ethyl-4-hydroxy-5-phenylthio-2,3,6,5-tetrahydro-benzothiepine-1,1-dioxide (21)

A mixture of 25 mg (0.085 mmole) of 19b, 0.27 g (2.7 mmole) of thiophenol, 0.37 g (2.7 mmole) of potassium carbonate, and 4 mL of DMF was stirred at room temperature under N₂ for 19 h. The reaction mixture was poured into water and extracted with methylene chloride. The methylene chloride layer was washed successively with 10% NaOH and brine, dried over MgSO₄, and concentrated in vacuo to give 0.19 g of semisolid disulfide. This material was purified by HPLC (5% EtOAc-hexane) to remove diphenyl disulfide in the first fraction. The column was then eluted with 20% EtOAc-hexane to give 17 mg of a first fraction, 4 mg of a second fraction and 11 mg of a third fraction which were three different isomers of 21, i.e. 21a, 21b, and 21c, respectively, by ¹H NMR and mass spectra.

Example 18

Alternative Synthesis of 6c and 6d

A. Preparation from 2-((2-Benzoylphenylthio)methyl)-2-ethylhexanal (2)

Step 1. 2-((2-Benzoylphenylsulfonyl)methyl)-2-ethylhexanal (44)

To a solution of 9.0 g (0.025 mole) of compound 2 in 100 mL of methylene chloride was added 14.6 g (0.025 mol) of 50-60% MCPBA portionwise. The reaction mixture was stirred at room temperature for 64 h then was stirred with 200 mL of 1 M potassium carbonate and

203

filtered through Celite. The methylene chloride layer was washed twice with 300 mL of 1 M potassium carbonate, once with 10% sodium hydroxide and once with brine. The insoluble solid formed during washing was removed by filtration through Celite. The methylene chloride solution was dried and concentrated in vacuo to give 9.2 g (95%) of semisolid. A portion (2.6 g) of this solid was purified by HPLC (10% ethyl acetate-hexane) to give 1.9 g of crystals, mp 135-136 °C

Step 2. 2-((2-Benzoylphenylsulfonyl)methyl)-2-ethylhexanal (45)

A solution of 50 g (0.13 mole) of crude 44 in 250 mL of methylene chloride was divided in two portions and charged to two Fisher-Porter bottles. To each bottle was charged 125 mL of methanol and 5 g of 10% Pd/C. The bottles were pressurized with 70 psi of hydrogen and the reaction mixture was stirred at room temperature for 7 h before being charged with an additional 5 g of 10% Pd/C. The reaction mixture was again hydrogenated with 70 psi of hydrogen for 7 h. This procedure was repeated one more time but only 1 g of Pd/C was charged to the reaction mixture. The combined reaction mixture was filtered and concentrated in vacuo to give 46.8 g of 45 as brown oil.

Step 3. (3a, 4a, 5a) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,6,5-tetrahydrobenzothiepine-1,1-dioxide (6c), and (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,6,5-tetrahydrobenzothiepine-1,1-dioxide (6d)

To a solution of 27.3 g (73.4 mmole) of 45 in 300 mL of anhydrous THF cooled to 2 °C with an ice bath was added 9.7 g (73.4 mmole) of 95% potassium t-butoxide. The reaction mixture was stirred for 20 min, quenched with 300 mL of 10% HCl and extracted with methylene chloride. The methylene chloride layer was dried over magnesium sulfate and concentrated in vacuo to give

208

24.7 g of yellow oil. Purification by HPLC (ethyl acetate-hexane) yielded 9.4 g of recovered 45 in the first fraction, 5.5 g (20%) of 6c in the second fraction and 6.5 g (24%) of 6d in the third fraction.

B. Preparation from 2-hydroxydiphenylmethane

Step 1. 2-mercaptodiphenylmethane (46)

To a 500 ml flask was charged 16 g (0.33 mol) of 60% sodium hydride oil dispersion. The sodium hydride was washed twice with 50 ml of hexane. To the reaction flask was charged 100 ml of DMF. To this mixture was added a solution of 55.2 g (0.3 mol) of 2-hydroxydiphenylmethane in 200 ml of DMF in 1 h while temperature was maintained below 30 °C by an ice-water bath. After complete addition of the reagent, the mixture was stirred at room temperature for 30 min then cooled with an ice bath. To the reaction mixture was added 49.4 g (0.4 mole) of dimethyl thiocarbamoyl chloride at once. The ice bath was removed and the reaction mixture was stirred at room temperature for 18 h before being poured into 300 ml of water. The organic was extracted into 500 ml of toluene. The toluene layer was washed successively with 10% sodium hydroxide and brine and was concentrated in vacuo to give 78.6 g of a yellow oil which was 95% pure dimethyl O-2-benzylphenyl thiocarbamate. This oil was heated at 280-300 °C in a kugelrohr pot under house vacuum for 30 min. The residue was kugelrohr distilled at 1 torr (180-280 °C). The distillate (56.3 g) was crystallized from methanol to give 37.3 g (46%) of the rearranged product dimethyl S-2-benzylphenyl thiocarbamate as a yellow solid. A mixture of 57 g (0.21 mole) of this yellow solid, 30 g of potassium hydroxide and 150 ml of methanol was stirred overnight then was concentrated in vacuo. The residue was diluted with 200 ml of water and extracted with ether. The aqueous layer was made acidic with concentrate HCl. The oily suspension was extracted into ether. The ether extract was dried over magnesium

sulfate and concentrated in vacuo. The residue was crystallized from hexane to give 37.1 g (88%) of 2-mercaptodiphenylmethane as a yellow solid.

Step 2. 2-((2-Benzylphenylthio)methyl)-2-ethylhexanal (47)

A mixture of 60 g (0.3 mole) of yellow solid from step 1, 70 g (0.3 mole) of compound 1 from preparation 1, 32.4 g (0.32 mole) of triethylamine, 120 ml of 2-methoxyethyl ether was held at reflux for 6 hr and concentrated in vacuo. The residue was triturated with 500 ml of water and 30 ml of concentrate HCl. The organic was extracted into 400 ml of ether. The ether layer was washed successively with brine, 10% sodium hydroxide and brine and was dried over magnesium sulfate and concentrated in vacuo. The residue (98.3 g) was purified by HPLC with 2-5% ethyl acetate-hexane as eluent to give 2-((2-benzylphenylthio)methyl)-2-ethylhexanal 47 as a yellow syrup.

Step 3. 2-((2-Benzylphenylsulfonyl)methyl)-2-ethylhexanal (45)

To a solution of 72.8 g (0.21 mole) of yellow syrup from step 2 in 1 liter of methylene chloride cooled to 10 °C was added 132 g of 50-60% MCPBA in 40 min. The reaction mixture was stirred for 2 h. An additional 13 g of 50-60% MCPBA was added to the reaction mixture. The reaction mixture was stirred for 2 h and filtered through Celite. The methylene chloride solution was washed twice with 1 liter of 1 M potassium carbonate then with 1 liter of brine. The methylene chloride layer was dried over magnesium sulfate and concentrated to 76 g of 2-((2-benzylphenylsulfonyl)methyl)-2-ethylhexanal 45 as a syrup.

Step 4. (3a,4a,5a) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (6c), and

(3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (6d)

Reaction of 45 with potassium t-butoxide according to the procedure in step 3 of procedure A gave pure 6c and 6d after HPLC.

Example 19

(3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-8-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (25) and (3a, 4a, 5a) 3-Butyl-3-ethyl-4-hydroxy-8-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (26)
Step 1. Preparation of 2-((2-benzoyl-4-methoxyphenylthio)methyl)-2-ethylhexanal (22)

2-Hydroxy-4-methoxybenzophenone was converted to the dimethyl O-2-benzoylphenyl thiocarbamate by methods previously described in example 18. The product can be isolated by recrystallization from ethanol. Using this improved isolation procedure no chromatography was needed. The thermal rearrangement was performed by reacting the thiocarbamate (5 g) in diphenyl ether at 260 °C as previously described. The improved isolation procedure which avoided a chromatography step was described below.

The crude pyrolysis product was then heated at 65 °C in 100 ml of methanol and 100 ml of THF in the presence of 3.5 g of KOH for 4 h. After removing THF and methanol by rotary evaporation the solution was extracted with 5 % NaOH and ether. The base layer was acidified and extracted with ether to obtain a 2.9 g of crude thiophenol product. The product was further purified by titrating the desired mercaptan into base with limited KOH. After acidification and extraction with ether pure 2-mercapto-4-methoxybenzophenone (2.3 g) was isolated.

2-mercapto-4-methoxybenzophenone can readily be converted to the 2-((2-benzoyl-4-

methoxyphenylthio)methyl)-2-ethylhexanal (22) by reaction with 2-ethyl-2-(mesyloxymethyl)hexanal (1) as previously described.

Step 2. 2-((2-Benzoyl-5-methoxyphenylsulfonyl)methyl)-2-ethylhexanal (23)

Substrate 22 was readily oxidized to 2-((2-benzoyl-5-methoxyphenyl-sulfonyl)methyl)-2-ethylhexanal (23) as described in example 18.

Step 3. 2-((2-benzyl-5-methoxyphenylsulfonyl)methyl)-2-ethylhexanal (24)

Sulfone 23 was then reduced to 2-((2-benzyl-5-methoxyphenyl-sulfonyl)methyl)-2-ethylhexanal (24) as described in example 18.

Step 4. (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-8-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (25) and (3a, 4a, 5a) 3-Butyl-3-ethyl-4-hydroxy-8-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (26)

A 3-neck flask equipped with a powder addition funnel, thermocouple and nitrogen bubbler was charged with 19.8 g (0.05 mole) of sulfone 24 in 100 ml dry THF. The reaction was cooled to -1.6 °C internal temperature by means of ice/salt bath. Slowly add 5.61 g (0.05 mole) of potassium t-butoxide by means of the powder addition funnel. The resulting light yellow solution was maintained at -1.6 °C. After 30 min reaction 400 ml of cold ether was added and this solution was extracted with cold 10 % HCl. The acid layer was extracted with 300 ml of methylene chloride. The organic layers were combined and dried over magnesium sulfate and after filtration stripped to dryness to obtain 19.9 g of product. ¹H nmr and glpc indicated a 96% conversion to a 50/50 mixture of 25 and

26. The only other observable compound was 4% starting sulfone 24.

The product was then dissolved in 250 ml of 90/10 hexane/ethyl acetate by warming to 50 °C. The solution was allowed to cool to room temperature and in this way pure 26 can be isolated. The crystallization can be enhanced by addition of a seed crystal of 26. After 2 crystallizations the mother liquor which was now 85.4% 25 and has a dry weight of 8.7 g. This material was dissolved in 100 ml of 90/10 hexane/ethyl acetate and 10 ml of pure ethyl acetate at 40 °C. Pure 25 can be isolated by seeding this solution with a seed crystal of 25 after storing it overnight at 0 °C.

Example 20

(3a,4a,5a) 3-Butyl-3-ethyl-4,8-dihydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (27)

In a 25 ml round bottomed flask, 1 g of 26 (2.5 mmoles) and 10 ml methylene chloride were cooled to - 78 °C with stirring. Next 0.7 ml of boron tribromide (7.5 mmole) was added via syringe. The reaction was allowed to slowly warm to room temperature and stirred for 6 h. The reaction was then diluted with 50 ml methylene chloride and washed with saturated NaCl and then water. The organic layer was dried over magnesium sulfate. The product (0.88g) 27 was characterized by NMR and mass spectra.

Example 21

General Alkylation of phenol 27

A 25 ml flask was charged with 0.15 g of 27 (0.38 mmole), 5 ml anhydrous DMF, 54 mg of potassium carbonate (0.38 mmole) and 140 mg ethyl iodide (0.9 mmole). The reaction was stirred at room temperature overnight. The reaction was diluted with 50 ml ethyl ether and washed with water (25 ml) then 5% NaOH (20 ml) and then sat. NaCl. After stripping off the solvent

the ethoxylated product 28 was obtained in high yield. The product was characterized by NMR and mass spectra. This same procedure was used to prepare products listed in table 1 from the corresponding iodides or bromides. For higher boiling alkyl iodides and bromides only one equivalent of the alkyl halide was used.




Table 1

Compound No.	R
27	H
26	Me
28	Et
29	hexyl
30	Ac
31	(CH ₂) ₆ -N-pthalimide

Example 22

(3a, 4a, 5a) 3-Butyl-3-ethyl-4-hydroxy-7-hydroxyamino-5-phenyl-2,3,4,5-tetrahydrobenzothiazepine-1,1-dioxide (37) and (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-7-hydroxyamino-5-phenyl-2,3,4,5-tetrahydrobenzothiazepine-1,1-dioxide (38)

Step 1. Preparation of 2-chloro-5-nitrodiphenylmethane (32)

Procedure adapted from reference: Synthesis - Stuttgart 9 770-772 (1986) Olah G. Et al

Under nitrogen, a 3 neck flask was charged with 45 g (0.172 mole) of 2-chloro-5-nitrobenzophenone in 345 ml methylene chloride and the solution was cooled to ice/water temperature. By means of an additional funnel, 150 g (0.172 mole) of trifluoromethane sulfonic acid in 345 ml methylene chloride was added slowly. Next 30 g of triethylsilane (0.172 mole) in 345 ml methylene chloride was added dropwise to the chilled solution. Both addition steps (trifluoromethane sulfonic acid and triethylsilane) were repeated. After the additions were completed the reaction was allowed to slowly warm up to room temperature and stirred for 12 h under nitrogen. The reaction mixture was then poured into a chilled stirred solution of 1600 ml of saturated sodium bicarbonate. Gas evolution occurred. Poured into a 4 liter separatory funnel and separated layers. The methylene chloride layer was isolated and combined with two 500 ml methylene chloride extractions

of the aqueous layer. The methylene chloride solution was dried over magnesium sulfate and concentrated in vacuo. The residue was recrystallized from hexane to give 39 g product. Structure 32 was confirmed by mass spectra and proton and carbon NMR.

Step 2. Preparation of 2-((2-benzyl-4-nitrophenylthio)methyl)-2-ethylhexenal (33)

The 2-chloro-5-nitrodiphenylmethane product 32 (40 g, 0.156 mole) from above was placed in a 2 liter 2 neck flask with water condenser. Next 150 ml DMSO and 7.18 g (0.156 mole) of lithium sulfide was added and the solution was stirred at 75 °C for 12 h. The reaction was cooled to room temperature and then 51.7 g of mesylate IV was added in 90 ml DMSO. The reaction mixture was heated to 80 °C under nitrogen. After 12 h monitored by TLC and added more mesylate if necessary. Continued the reaction until the reaction was completed. Next the reaction mixture was slowly poured into a 1900 ml of 5% acetic aqueous solution with stirring, extracted with 4 X 700 ml of ether, and dried over MgSO₄. After removal of ether, 82.7 g of product was isolated. The material can be further purified by silica gel chromatography using 95% hexane and 5 % ethyl acetate. If pure mesylate was used in this step there was no need for further purification. The product 33 was characterized by mass spectra and NMR.

Step 3. Oxidation of the nitro product 33 to the sulfone 2-((2-benzyl-4-nitrophenylsulfonyl)methyl)-2-ethylhexenal (34)

The procedure used to oxidize the sulfide 33 to the sulfone 34 has been previously described.

Step 4. Reduction of 34 to 2-((2-benzyl-4-hydroxyaminophenylsulfonyl)methyl)-2-ethylhexenal (35)

A 15 g sample of 34 was dissolved in 230 ml of ethanol and placed in a 500 ml rx flask under nitrogen. Next 1.5 g of 10 wt. % Pd/C was added and hydrogen gas was bubbled through the solution at room temperature until the nitro substrate 34 was consumed. The reaction could be readily monitored by silica gel TLC using 80/20 hexane/EtOAc. Product 35 was isolated by filtering off the Pd/C and then stripping off the EtOH solvent. The product was characterized by NMR and mass spectra.

Step 5. Preparation of the 2-((2-benzyl-6-*N*,*O*-di-(*t*-butoxy-carbonyl)hydroxyaminophenylsulfonyl)methyl)-2-ethylhexanal (36).

A 13.35 g sample of 35 (0.0344 mole) in 40 ml of dry THF was stirred in a 250 ml round bottomed flask. Next added 7.52 g (0.0344 mole) of di-*t*-butyl dicarbonate in 7 ml THF. Heated at 60 °C overnight. Stripped off THF and redissolved in methylene chloride. Extracted with 1 % HCl; and then 5% sodium bicarbonate.

The product was further purified by column chromatography using 90/10 hexane/ethyl acetate and then 70/30 hexane/ethyl acetate. The product 36 was obtained (4.12 g) which appeared to be mainly the di-(*t*-butoxycarbonyl) derivatives by proton NMR.

Step 6. (3a,4a,5a) 3-Butyl-3-ethyl-4-hydroxy-7-hydroxyamino-5-phenyl-2,3,4,5-tetrahydrobenzothioipino-1,1-dioxido (37) and (3a,4b,5b) 3-Butyl-3-ethyl-4-hydroxy-7-hydroxyamino-5-phenyl-2,3,4,5-tetrahydrobenzothioipino-1,1-dioxido (38)

A 250ml 3-neck round bottomed flask was charged with 4 g of 36 (6.8 mmoles), and 100 ml of anhydrous THF and cooled to -78 °C under a nitrogen atmosphere. Slowly add 2.29 g potassium tert-butoxide (20.4 mmoles) with stirring and maintaining a -78 °C reaction temperature. After 1 h at -78 °C the addition of base was completed and the temperature was brought to -10 °C by means of a

ice/salt bath. After 3 h at -10 °C, only trace 36 remained by TLC. Next add 35 ml of deionized water to the reaction mixture at -10 °C and stirred for 5 min. Stripped off most of the THF and added to separatory funnel and extracted with ether until all of the organic was removed from the water phase. The combined ether phases were washed with saturated NaCl and then dried over sodium sulfate. The only products by TLC and NMR were the two BOC protected isomers of 37 and 38. The isomers were separated by silica gel chromatography using 85% hexane and 15 % ethyl acetate; BOC-37 (0.71 g) and BOC- 38 (0.78 g).

Next the BOC protecting group was removed by reacting 0.87 g of BOC-38 (1.78 mmoles) with 8.7 ml of 4 M HCl (34.8 mmoles) in dioxane for 30 min. Next added 4.74 g of sodium acetate (34.8 mmoles) to the reaction mixture and 16.5 ml ether and stirred until clear. After transferring to a separatory funnel extracted with ether and water and then dried the ether layer with sodium sulfate. After removing the ether, 0.665 g of 38 was isolated. Isomer 37 could be obtained in a similar procedure.

Example 23

(3a,4a,5a) 3-Butyl-3-ethyl-7-(*n*-hexylamino)-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothioipino-1,1-dioxido (40) and (3a,4b,5b) 3-Butyl-3-ethyl-7-(*n*-hexylamino)-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothioipino-1,1-dioxido (41)

Step 1. 2-((2-Benzyl-6-(*n*-

hexylamino)phenylsulfonyl)methyl)-2-ethylhexanal (39)

In a Fischer porter bottle weighed out 0.5 g of 34 (1.2 mmoles) and dissolved in 3.8 ml of ethanol under nitrogen. Next added 0.1 g of Pd/C and 3.8 ml of hexanal. Seal and pressure to 50 psi of hydrogen gas. Stirred for 48 h. After filtering off the catalyst and removing the solvent by rotary evaporation 39 was

isolated by column chromatography (0.16 g) using 90/10 hexane ethyl acetate and gradually increasing the mobile phase to 70/30 hexane/ethyl acetate. The product was characterized by NMR and mass spectra.

Step 2. (3a, 4a, 5a) 3-Butyl-3-ethyl-7-(n-hexylamino)-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (40) and (3a, 4b, 5b) 3-Butyl-3-ethyl-7-(n-hexylamino)-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (41)

A 2-neck, 25 ml round bottomed flask with stir bar was charged with 0.158 g 39 (0.335 mmole) and 5 ml anhydrous THF under nitrogen. Cool to -10 °C by means of a salt/water bath. Slowly add 0.113 g of potassium tert butoxide (0.335 mmole). After 15 min at -10 °C all of the starting material was consumed by TLC and only the two isomers 40 and 41 were observed. Next added 5 ml of chilled 10% HCl and stirred at -10 °C for 5 min. Transferred to a separatory funnel and extract with ether. Dried over sodium sulfate. Proton NMR of the dried product (0.143 g) indicated only the presence of the two isomers 40 and 41. The two isomers were separated by silica gel chromatography using 90/10 hexane ethyl acetate and gradually increasing the mobile phase to 70/30 hexane/ethyl acetate. 40 (53.2 mg); 41 (58.9 mg).

Example 24

Quaternization of amine substrates 40 and 41

Amine products such as 40 and 41 can be readily alkylated to quaternary salts by reaction with alkyl halides. For example 40 in DMF with 5 equivalents of methyl iodide in the presence of 2,6 dimethyl lutidine produces the dimethylhexylamino quaternary salt.

Example 25

(3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-5-(4-iodophenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (42)

In a 25 ml round bottomed flask 0.5 g (1.3 mmole) of 64, 0.67 g of mercuric triflate were dissolved in 20 ml of dry methylene chloride with stirring. Next 0.34 g of Iodine was added and the solution was stirred at room temperature for 30 h. The reaction was then diluted with 50 ml methylene chloride and washed with 10 ml of 1 M sodium thiosulfate; 10 ml of saturated KI ; and dried over sodium sulfate. See Tetrahedron, Vol.50, No. 17, pp 5139-5146 (1994) Bachki, F. Et al. Mass spectrum indicated a mixture of 6d , mono iodide 42 and a diiodide adduct. The mixture was separated by column chromatography and 42 was characterized by NMR and mass spectra.

Example 26

(3a, 4b, 5b) 3-Butyl-5-(4-carbomethoxyphenyl)-3-ethyl-4-hydroxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (43)

A 0.1 g sample of 42 (0.212 mmole), 2.5 ml dry methanol, 38 µl triethylamine (0.275 mmole) , 0.3 ml toluene and 37 mg of palladium chloride (0.21 mmole) was charged to a glass lined mini reactor at 300 psi carbon monoxide. The reaction was heated at 100 °C overnight. The catalyst was filtered and a high yield of product was isolated.

The product was characterized by NMR and mass spectra.

Note the ester functionalized product 43 can be converted to the free acid by hydrolysis.

Example 27

(3a, 4a, 5a) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (48), and (3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-7-

methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (49)

Step 1. 2-Mercapto-5-methoxybenzophenone (50)

Reaction of 66.2 g of 4-methoxythiophenol with 360 ml of 2.5 N n-butyllithium, 105 g of tetramethylethylenediamine and 66.7 g of benzonitrile in 600 ml cyclohexane according to the procedure in WO 93/16055 gave 73.2 g of brown oil which was kugelrohr distilled to remove 4-methoxythiophenol and gave 43.86 g of crude 50 in the pot residue.

Step 2. 2-((2-Benzoyl-4-methoxyphenylthio)methyl)-2-ethylhexanal (51)

Reaction of 10 g (0.04 mole) of crude 50 with 4.8 g (0.02 mole) of mesylate 1 and 3.2 ml (0.23 mole) of triethylamine in 50 ml of diglyme according to the procedure for the preparation of 2 gave 10.5 g of crude product which was purified by HPLC (5% ethyl acetate-hexane) to give 1.7 g (22%) of 51.

Step 3. 2-((2-Benzoyl-4-methoxyphenylsulfonyl)methyl)-2-ethylhexanal (52)

A solution of 1.2 g (3.1 mmoles) of 51 in 25 ml of methylene chloride was reacted with 2.0 g (6.2 mmoles) of 50-60% MCPBA according to the procedure of step 2 of procedure A in example 18 gave 1.16 g (90%) of 52 as a yellow oil.

Step 4. 2-((2-Benzoyl-4-methoxyphenylsulfonyl)methyl)-2-ethylhexanal (53)

Hydrogenation of 1.1 g of 52 according to the procedure of step 3 of procedure A of example 18 gave 53 as a yellow oil (1.1 g).

221

Step 5. (3a,4a,5a) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (48), and (3a,4b,5b) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (49)

A solution of 1.1 g of 53, 0.36 g of potassium t-butoxide and 25 ml of anhydrous THF was held at reflux for 2 h and worked up as in step 4 of procedure A of example 18 to give 1.07 g of a crude product which was purified by HPLC to give 40 mg (4%) of 48 as crystals, mp 153-154 °C and 90 mg (8%) of 49 as solid, mp 136-140 °C.

Example 28

5-Phenyl-2,3-dihydrospirobenzothiepine-3,1'-cyclohexane (57)

Step 1. 1-(Hydroxymethyl)-cyclohexanecarboxaldehyde (54)

To a cold (0 °C) mixture of 100 g (0.891 mole) of cyclohexanecarboxaldehyde, 76.5 g of 37% of formaldehyde in 225 ml of methanol was added dropwise 90 ml of 1 N Sodium hydroxide in 1 h. The reaction mixture was stirred at room temperature over 48 h then was evaporated to remove methanol. The reaction mixture was diluted with water and extracted with methylene chloride. The organic layer was washed with water, brine, and dried over sodium sulfate and concentrated under vacuum to give 75 g (59.7%) of thick oil. Proton NMR and mass spectra were consistent with the product.

Step 2. 1-(mesyloxymethyl)cyclohexanecarboxaldehyde (55)

To a cold (0 °C) mixture of alcohol 54 (75 g, 0.54 mole) and 65.29 g (0.57 mole) of methanesulfonyl chloride in 80 ml of methylene chloride was added a solution of pyridine (47.96 g, 0.57 mole) in 40 ml of methylene

222

chloride. The reaction mixture was stirred at room temperature for 18 h then quenched with water, acidified with conc. HCl and extracted with methylene chloride. The organic layer was washed with water, brine, and dried over sodium sulfate and concentrated under vacuum to give 91.63 g (77.8%) of thick oil. Proton NMR and mass spectra were consistent with the product.

Step 3. 1-((2-Benzoylphenylthio)methyl)cyclohexanecarboxaldehyde (56)

A mixture of 69 g (0.303 mole) of 2-mercaptobenzophenone, 82 g (0.303 mole) of mesylate 55, 32 g of triethylamine, and 150 ml of diglyme was stirred and held at reflux for 24 h. The mixture was cooled, poured into dil. HCl and extracted with methylene chloride. The organic layer was washed with 10% NaOH, water, brine, and dried over sodium sulfate and concentrated under vacuum to remove excess diglyme. This was purified by silica gel flush column (5% EtOAc: Hexane) and gave 18.6 g (75.98%) of yellow oil. Proton NMR and mass spectra were consistent with the product.

Step 4. 5-Phenyl-2,3-dihydrospirobenzochloephine-3,1'-cyclohexane (57)

To a mixture of 6.19 g of zinc dust and 100 ml of dry DME was added TiCl₄ (16.8 g, 0.108 mole). The reaction mixture was heated to reflux for 2 h. A solution of compound 56 (8.3 g, 0.023 mole) in 50 ml of DME was added dropwise to the reaction mixture in 1 h and the mixture was held at reflux for 18 h. The mixture was cooled, poured into water and extracted with ether. The organic layer was washed with water, brine, and dried over sodium sulfate, filtered through celite and concentrated under vacuum. The residue was purified by HPLC (10% EtOAc: Hexane) to give 4.6 g (64%) of white

223

solid, mp 90-91 °C. Proton and carbon NMR and mass spectra were consistent with the product.

Example 22

8b-Phenyl-1a,2,3,8b-tetrahydrospiro(benzochloephino[4,5-b]oxirene-2,1'-cyclohexane)-4,4-dioxide (58)

To a solution of 57 (4.6 g, 15 mmole) in 50 ml chloroform under nitrogen was added 55% MCPBA (16.5 g, 52.6 mmole) portionwise with spatula. The reaction was held at reflux for 18 h and washed with 10% NaOH (3X), water, brine, and dried over sodium sulfate and concentrated under vacuum to give 5 g of crude product. This was recrystallized from Hexane/EtOAc to give 4.31 g (81%) of yellow solid, mp 154-155 °C. Proton and carbon NMR and mass spectra were consistent with the product.

Example 30

trans-4-Hydroxy-5-phenyl-2,3,4,5-tetrahydro spiro(benzochloephine-3,1'-cyclohexane)-1,1-dioxide (59)

A mixture of 0.5 g (1.4 mmole) of 58, 20 ml of ethanol, 10 ml of methylene chloride and 0.4 g of 10% Pd/C catalyst was hydrogenated with 70 psi hydrogen for 3 h at room temperature. The crude reaction slurry was filtered through Celite and evaporated to dryness. The residue was purified by HPLC (10% EtOAc:Hexane, 25% EtOAc:Hexane). The first fraction was 300 mg (60%) as a white solid, mp 99-100 °C. Proton NMR showed this was a trans isomer. The second fraction gave 200 mg of solid which was impure cis isomer.

Example 31

cis-4-Hydroxy-5-phenyl-2,3,4,5-tetrahydro spiro(benzochloephine-3,1'-cyclohexane)-1,1-dioxide (60)

224

To a solution of 0.2 g (0.56 mmole) of **59** in 20 ml of CH_2Cl_2 , was added 8 g of 50% NaOH and one drop of Aliquat-336 (methyltricaprylammonium chloride) phase transfer catalyst. The reaction mixture was stirred for 10 h at room temperature. Twenty g of ice was added to the mixture and the mixture was extracted with CH_2Cl_2 (3x10 ml) washed with water, brine and dried over MgSO_4 and concentrated in vacuo to recover 0.15 g of crude product. This was recrystallized from Hexane/EtOAc to give 125 mg of white crystal, mp 209-210 °C. Proton and carbon NMR and mass spectra were consistent with the product.

Example 32

(**3a, 4a, 5a**) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine (**61**), and (**3a, 4b, 5b**) 3-Butyl-3-ethyl-4-hydroxy-5-phenyl-2,3,4,5-tetrahydrobenzothiepine (**62**)

To a solution of 0.5 g (1.47 mmole) of compound **47** in 5 ml of anhydrous THF was added 0.17 g (1.47 mmole) of 95% potassium t-butoxide. The reaction mixture was stirred at room temperature for 18 h and quenched with 10 ml of 10% HCl. The organic was extracted into methylene chloride. The methylene chloride extract was dried over magnesium sulfate and concentrated in vacuo. The residue was purified by HPLC (2% EtOAc-hexane) to give 47 mg of **61** in the second fraction and 38 mg of **62** in the third fraction. Proton NMR and mass spectra were consistent with the assigned structures.

Example 33

(**3a, 4a, 5a**) 3-Butyl-3-ethyl-4-hydroxy-7-amino-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (**63**) and (**3a, 4b, 5b**) 3-Butyl-3-ethyl-4-hydroxy-7-amino-5-phenyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (**64**)

An autoclave was charged with 200 mg of **37** in 40 cc ethanol and .02 g 10 % Pd/C. After purging with

nitrogen theclave was charged with 100 psi hydrogen and heated to 55 °C. The reaction was monitored by TLC and mass spec and allowed to proceed until all of **37** was consumed. After the reaction was complete the catalyst was filtered and the solvent was removed in vacuo and the only observable product was amine **63**. This same procedure was used to produce **64** from **38**.

Example 34

(**3a, 4a, 5a**) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-(3'-methoxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (**65**), and (**3a, 4b, 5b**) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-(3'-methoxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (**66**). Alkylation of e-methoxyphenol with 3-methoxybenzyl chloride according to the procedure described in J. Chem. Soc. 2431 (1958) gave 4-methoxy-2-(3'-methoxybenzyl)phenol in 35% yield. This material was converted to compound **65**, mp 138.5-141.5 °C, and compound **66**, mp 115.5-117.5 °C, by the procedure similar to that in Example 18 method B.

Example 35

(**3a, 4a, 5a**) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-(3'-trifluoromethyl)phenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (**67**), and (**3a, 4b, 5b**) 3-Butyl-3-ethyl-4-hydroxy-7-methoxy-5-(3'-trifluoromethyl)phenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (**68**). Alkylation of 4-methoxyphenol with 3-(trifluoromethyl)benzyl chloride according to the procedure described in J. Chem. Soc. 2431 (1958) gave 4-methoxy-2-(3'-(trifluoromethyl)benzyl)phenol. This material was converted to compound **67**, mp 226.5-228 °C, and compound **68**, mp 188-190 °C, by the procedure similar to that in Example 18 method B.

Example 36

(3a, 4a, 5a) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (69), and (3a, 4b, 5b) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (70).

Alkylation of 4-methoxyphenol with 4-fluorobenzyl chloride according to the procedure described in J. Chem. Soc, 2431 (1958) gave 4-methoxy-2-(4'-fluorobenzyl)phenol. This material was converted to compound 69 and compound 70 by the procedure similar to that in Example 18 method B.

Example 37

(3a, 4a, 5a) 3-Butyl-3-ethyl-5-(3'-fluorophenyl)-4-hydroxy-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (71), and (3a, 4b, 5b) 3-Butyl-3-ethyl-5-(3'-fluorophenyl)-4-hydroxy-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (72).

Alkylation of 4-methoxyphenol with 3-fluorobenzyl chloride according to the procedure described in J. Chem. Soc, 2431 (1958) gave 4-methoxy-2-(3'-fluorobenzyl)phenol. This material was converted to compound 71 and compound 72 by the procedure similar to that in Example 18 method B.

Example 38

(3a, 4a, 5a) 3-Butyl-3-ethyl-5-(2'-fluorophenyl)-4-hydroxy-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (73), and (3a, 4b, 5b) 3-Butyl-3-ethyl-5-(2'-fluorophenyl)-4-hydroxy-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (74).

Alkylation of 4-methoxyphenol with 2-fluorobenzyl chloride according to the procedure described in J. Chem. Soc, 2431 (1958) gave 4-methoxy-2-(2'-fluorobenzyl)phenol. This material was converted to

227

compound 73 and compound 74 by the procedure similar to that in Example 18 method B.

Example 39

(3a, 4a, 5a) 3-Butyl-7-bromo-3-ethyl-4-hydroxy-5-(3'-methoxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (75), and (3a, 4b, 5b) 3-Butyl-7-bromo-3-ethyl-4-hydroxy-5-(3'-methoxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (76).

Alkylation of 4-bromophenol with 3-methoxybenzyl chloride according to the procedure described in J. Chem. Soc, 2431 (1958) gave 4-bromo-2-(3'-methoxybenzyl)phenol. This material was converted to compound 75, mp 97-101.5 °C, and compound 76, mp 102-106 °C, by the procedure similar to that in Example 18 method B.

Example 40

(3a, 4a, 5a) 3-Butyl-3-ethyl-7-fluoro-5-(4'-fluorophenyl)-4-hydroxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (77), and (3a, 4b, 5b) 3-Butyl-3-ethyl-7-fluoro-5-(4'-fluorophenyl)-4-hydroxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (78).

Alkylation of 4-fluorophenol with 4-fluorobenzyl chloride according to the procedure described in J. Chem. Soc, 2431 (1958) gave 4-fluoro-2-(4'-fluorobenzyl)phenol. This material was converted to compound 77, mp 228-230 °C, and compound 78, mp 134.5-139 °C, by the procedure similar to that in Example 18 method B.

Example 41

(3a, 4a, 5a) 3-Butyl-3-ethyl-7-fluoro-4-hydroxy-5-(3'-methoxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (79), and (3a, 4b, 5b) 3-Butyl-3-ethyl-7-fluoro-

228

40hydroxy-5-(3'-methoxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (80).

Alkylation of 4-fluorophenol with 3-methoxybenzyl chloride according to the procedure described in J. Chem. Soc. 2431 (1958) gave 4-fluoro-2-(3'-methoxybenzyl)phenol. This material was converted to compound 79, as a solid and compound 80, mp 153-155 °C, by the procedure similar to that in Example 18 method B.

Example 42

(3a, 4b, 5b) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-methylthio-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (81).

A mixture of 0.68 (1.66 mmol) of compound 77, 0.2 g (5 mmol) of sodium methanethiolate and 15 ml of anhydrous DMF was stirred at room temperature for 16 days. The reaction mixture was dilute with ether and washed with water and brine and dried over M_2SO_4 . The ether solution was concentrated in vacuo. The residue was purified by HPLC (20% ethyl acetate in hexanes). The first fraction was impure (3a, 4a, 5a) 3-butyl-3-ethyl-4-hydroxy-7-methylthio-5-(4'-fluorophenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide. The second fraction was compound 81, mp 185-186.5 °C.

Example 43

(3a, 4b, 5b) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-(1-pyrrolidinyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (82).

A mixture of 0.53 g (1.30 mmol) of compound 78 and 5 ml of pyrrolidine was held at reflux for 1 h. The reaction mixture was diluted with ether and washed with water and brine and dried over M_2SO_4 . The ether solution was concentrated in vacuo. The residue was

229

crystallized from ether-hexanes to give compound 82, mp 174.5-177 °C.

Example 44

(3a, 4b, 5b) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-(1-morpholinyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (83).

A mixture of 0.4 g (0.98 mmol) of compound 78 and 5.0 g (56 mmol) of morpholine was held at reflux for 2 h and concentrated in vacuo. The residue was diluted with ether (30 ml) and washed with water and brine and dried over M_2SO_4 . The ether solution was concentrated in vacuo. The residue was recrystallized from ether-hexanes to give compound 83, mp 176.5-187.5 °C.

Example 45

(3a, 4a, 5a) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-methyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (84), and (3a, 4b, 5b) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-methyl-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (85).

Alkylation of 4-methylphenol with 4-fluorobenzyl chloride according to the procedure described in J. Chem. Soc. 2431 (1958) gave 4-methyl-2-(4'-fluorobenzyl)phenol. This material was converted to compound 84 and compound 85 by the procedure similar to that in Example 18 method B.

Example 46

(3a, 4b, 5b) 3-Butyl-3-ethyl-4-hydroxy-5-(4'-hydroxyphenyl)-7-methoxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (86), and (3a, 4b, 5b) 3-Butyl-3-ethyl-4,7-dihydroxy-5-(4'-hydroxyphenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (87).

230

To a solution of 0.52 (1.2 mmol) of compound 66 in 20 ml of methylene chloride was added 1.7 g (6.78 mmol) of boron tribromide. The reaction mixture was cooled to -78 °C and was stirred for 4 min. An additional 0.3 ml of boron tribromide was added to the reaction mixture and the reaction mixture was stirred at -78 °C for 1 h and quenched with 2 N HCl. The organic was extracted into ether. The ether layer was washed with brine, dried over $MgSO_4$, and concentrated in vacuo. The residue (0.48 g) was purified by HPLC (30% ethyl acetate in hexanes). The first fraction was 0.11 g of compound 86 as a white solid, mp 171.5-173 °C. The second fraction was crystallized from chloroform to give 0.04 g of compound 87 as a white solid, mp 264 °C (dec).

Example 47

(3a,4b,5b) 3-Butyl-3-ethyl-4,7-dihydroxy-5-(4'-fluorophenyl)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (88).

Reaction of compound 70 with excess boron tribromide at room temperature and worked up as in Example 46 gave compound 88 after an HPLC purification.

Example 48

(3a,4b,5b) 3-Butyl-3-ethyl-5-(4'-fluorophenyl)-4-hydroxy-7-(1-azetidiny)-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (89).

A mixture of 0.20 g (0.49 mmol) of compound 78, and 2.0 g (35 mmol) of azetidine was held at reflux for 3 h and concentrated in vacuo. The residue was diluted with ether (30 ml) and washed with water and brine and dried over $MgSO_4$. The ether solution was concentrated on a steam bath. The separated crystals were filtered to give 0.136 g of 89 as prisms, mp 196.5-199.5 °C.

Example 49


(3a,4a,5a) 3-Butyl-3-ethyl-5-(3'-methoxyphenyl)-4-hydroxy-7-methylthio-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (90). (3a,4b,5b) 3-Butyl-3-ethyl-5-(3'-methoxyphenyl)-4-hydroxy-7-methylthio-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide (91).

A mixture of 0.4 g (0.95 mmol) of compound 79, 0.08 g (1.14 mmol) of sodium methanethiolate and 15 ml of anhydrous DMF was stirred at 60 °C for 2 h. An additional 1.4 mmol of sodium methanethiolate was added to the reaction mixture and the mixture was stirred at 60 °C for an additional 2 h. The reaction mixture was triturated with 100 ml of water and extracted methylene chloride. The methylene chloride water mixture was filtered through Celite and the methylene chloride layer was dried over $MgSO_4$ and concentrated in vacuo. The first fraction (0.1 g) was compound 90, mp 117-121 °C. The second fraction (0.16 g) was compound 91, mp 68-76 °C.

Example 50

Preparation of polyethyleneglycol functionalized benzothiepine **A**.

5



No. 141

No. 136

A 50 ml rb flash under a nitrogen atmosphere was charged with 0.54 g of M-Tres-5000 (Polyethyleneglycol Tresylate [methoxy-PEG-Tres, MW 5000] purchased from Shearwater Polymers Inc., 2130 Memorial Parkway, SW, Huntsville, Alabama 35801), 0.055 g Compound No. 136, 0.326 C₆CO, and 2cc anhydrous acetonitrile. The reaction was stirred at 30 C for 5 days and then the solution was filtered to remove salts. Next, the acetonitrile was removed under vacuum and the product was dissolved in THF and then precipitated by addition of hexane. The polymer precipitate was isolate by filtration from the solvent mixture (THF/hexane). This precipitation procedure was continued until no Compound No. 136 was detected in the precipitated product (by TLC SiO₂). Next, the polymer precipitate was dissolved in water and filtered and the water soluble polymer was dialyzed for 48 hours through a cellulose dialysis tube (Spectrum® 7, 45 mm x 0.5 ft, cutoff 1,000 MW). The polymer solution was then removed from the dialysis tube and lyophilized until dried. The NMR was consistent with the desired product **A** and gel permeation


233

chromatography indicated the presence of a 4500 MW polymer and also verified that no free Compound No. 136 was present. This material was active in the IBAT in vitro cell assay.

5


Example 51

Preparation of Compound 140



10

No. 140



No. 111


15

A 2-necked 50 ml round bottom Flask was charged with 0.42g of Tres-3400 (Polyethyleneglycol Tresylate [Tres-PEG-Tres, MW 3400] purchased from Shearwater Polymers Inc., 2130 Memorial Parkway, SW, Huntsville, Alabama 35801), 0.1 potassium carbonate, 0.100g of Compound No. 111 and 5 ml anhydrous DMF. Stir for 6 days at 27 °C. TLC indicated the disappearance of the starting

20


234

Compound No. 111. The solution was transferred to a separatory funnel and diluted with 50 cc methylene chloride and then extracted with water. The organic layer was evaporated to dryness by means of a rotary evaporator. Dry wt. 0.4875 g. Next, the polymer was dissolved in water and then dialyzed for 48 hours at 40 °C through a cellulose dialysis tube (spectrum® 7, 45mm x 0.5 ft, cutoff 1,000 MW). The polymer solution was then removed from the dialysis tube and lyophilized until dried 0.341 g). NMR was consistent with the desired product B.


Example 52

No. 134

A 10 cc vial was charged with 0.21 g of Compound No. 136 (0.5mmoles), 0.17g (1.3 mmoles)potassium carbonate, 0.6g (1.5 mmoles) of 1,2-bis-(2-iodoethoxy)-ethane and 10 cc DMF. The reaction was stirred for 4 days at room temperature and then worked up by washing with ether/water. The ether layer was stripped to dryness and the desired product Compound No. 134 was isolated on a silica gel column using 80/20 hexane ethyl acetate.

Example 53

No. 112

Example 54


No. 113

A two necked 25 ml round bottom flask was charged with 0.5g (1.24mmoles) of 69462, 13 mls of anhydrous DMF, 0.055g of 60% NaH dispersion and 0.230g (0.62 mmoles) of 1,2-Bis (2-iodoethoxy)ethane) at 10 °C under nitrogen. Next, the reaction was slowly heated to 40 °C. After 14 hours all of the Compound No. 113 was consumed and the reaction was cooled to room temperature and extracted with ether/water. The ether layer was evaporated to dryness and then chromatographed on Silicage (80/20 ethyl


acetate/hexane). Isolated Compound No. 112 (0.28 g) was characterized by NMR and mass spec.

Example 55

5



No. 135



No. 136

In a 50 ml round bottom Flask, add 0.7g (1.8 mmoles) of Compound No. 136, 0.621g of potassium carbonate, 6 ml DMF, and 0.33g of 1,2-Bis (2-iodoethoxyethane). Stir at 40 °C under nitrogen for 12 hours. The workup and isolation was the same procedure for Compound No. 112.

10

Examples 56 and 57 (Compound Nos. 131 and 137)

The compositions of these compounds are shown in Table

3.

The same procedure as for Example 55 except appropriate benzoethiopia was used.

20

237


Example 58 (Compound No. 139)

The composition of this compound is shown in Table 3. Same procedure as for Example 55 with appropriate benzoethiopia 1,6 diiodohexane was used instead of 1,2-Bis (2-iodoethoxyethane).

5

Example 59 (Compound No. 101)

10




No. 101

This compound is prepared by condensing the 7-NH, benzoethiopia with the 1,12-dodecane dicarboxylic acid or acid halide.

15

Example 60 (Compound No. 104)


238




No. 104

2-Chloro-4-nitrobenzophenone is reduced with triethylsilane and trifluoromethane sulfonic acid to 2-chloro-4-nitrodiphenylmethane 32. Reaction of 32 with lithium sulfide followed by reacting the resulting sulfide with mesylate IV gives sulfide-aldehyde XXIII. Oxidation of XXIII with 2 equivalents of MCPBA yields sulfone-aldehyde XXV (see Scheme 5). Reduction of the sulfone-aldehyde XXV formaldehyde and 100 psi hydrogen and 55 C for 12 hours catalyzed by palladium on carbon in the same reaction vessel yields the substituted dimethylamine derivative XXVII. Cyclization of XXVII with potassium t-butoxide yields a mixture of substituted amino derivatives of this invention Compound No. 104.


Scheme 6



239




Example 61



No. 102

A 1 oz. Fisher-porter bottle was charged with 0.14 g (0.34 mmoles) of 70112, 0.97 gms (6.8 mmoles) of methyl iodide, and 7 ml of anhydrous acetonitrile. Heat to 50 °C for 4 days. The quat. Salt Compound No. 192 was isolated by concentrating to 1 cc acetonitrile and then precipitating with diethyl ether.


Example 62



No. 125


A 0.1 g (0.159 mmoles) sample of Compound No. 134 was dissolved in 15 ml of anhydrous acetonitrile in a Fischer-porter bottle and then trimethylamine was bubbled through the solution for 5 minutes at 0 °C and then capped and warmed to room temperature. The reaction was stirred overnight and the desired product was isolated by removing solvent by rotary evaporation.

10

Example 63 (Compound No. 295)

No. 295

15




No. 113

241

Sodium Hydride 60% (11 mg, 0.27 mmoles) in 1 cc of acetonitrile at 0 °C was reacted with 0.248 mmoles (.10 g) of Compound No. 54 in 2.5cc of acetonitrile at 0 °C. Next, 0.980g 2.48 mmoles) of 1,2-Bis [2-iodoethoxyethane]. After warming to room temperature, stir for 14 hours. The product was isolated by column chromatography.

5

Example 64 (Compound No. 286)

No. 286


Following a procedure similar to the one described in Example 86, infra (see Compound No. 118), the title compound was prepared and purified as a colorless solid; mp 180-181 °C; ¹H NMR (CHCl₃) δ 0.85 (t, J = 6 Hz, 3H, 0.92 (t, J = 6 Hz, 3H), 1.24-1.42 (m, 2H), 1.46-1.56 (m, 1H), 1.64-1.80 (m, 1H), 2.24-2.38 (m, 1H), 3.15 (AB, J_{AB} = 15 Hz, Dv = 42 Hz, 2H), 4.20 (d, J = 8 Hz, 1H), 5.13 (s, 2H), 5.53 (s, 1H), 6.46 (s, 1H), 6.68 (s, 1H), 7.29-7.51 (m, 10H), 7.74 (d, J = 8 Hz, 1H), 8.06 (d, J = 8 Hz, 1H). FAIMS m/z 494 (M⁺H), HRMS calcd for (M⁺H) 494.2001, found 494.1993. Anal. Calcd. for C₂₄H₂₁NO₃S: C, 68.13; H, 6.33; N, 2.84. Found: C, 68.19; H, 6.56; N, 2.74.

15

20

25


242

Example 65 (Compound No. 287)

No. 287

Following a procedure similar to the one described in Example 89, *infra* (see Compound No. 121), the title compound was prepared and purified as a colorless


solid: mp 245-246 °C, ¹H NMR (CDCl₃) δ 0.84 (t, J = 6 Hz, 3H), 0.92 (t, J = 6 Hz, 3H), 1.28, (d, J = 8 Hz, 1H), 1.32-1.42 (m, 1H), 1.48-1.60 (m, 1H), 1.64-1.80 (m, 1H), 2.20-2.36 (m, 1H), 3.09 (AB, J_M = 15 Hz, Dv = 42 Hz, 2H), 3.97 (bs, 2H), 4.15 (d, J = 8 Hz, 1H), 5.49 (s, 1H), 5.95 (s, 1H), 6.54 (d, J = 7 Hz, 1H), 7.29-7.53 (m, 5H), 7.88 (d, J = 8 Hz, 1H); ESMS 366 (M+1).
Anal. Calcd. for C₂₄H₂₃NO₂S: C, 66.82; H, 7.01; N, 3.90.
 Found: C, 66.54; H, 7.20; N, 3.69.

Example 66 (Compound No. 288)

No. 288

243

Following a procedure similar to the one described in Example 89, *infra* (see Compound No. 121), the title compound was prepared and purified by silica gel chromatography to give the desired product as a colorless solid: mp 185-186°C; ¹H NMR (CDCl₃) δ 1.12 (s, 3H), 1.49 (s, 3H), 3.00 (d, J = 15 Hz, 1H), 3.28 (d, J = 15 Hz, 1H), 4.00 (s, 1H), 5.30 (s, 1H), 5.51 (s, 1H), 5.97 (s, 1H), 6.56 (dd, J = 2.1, 8.4 Hz, 1H), 7.31-7.52 (m, 5H), 7.89 (d, J = 8.4 Hz, 1H). MS (FAB+) (M+H) m/z 332.

Example 67 (Compound No. 289)


No. 289

Following a procedure similar to the one described in Example 89 (see Compound No. 121), the title

compound was prepared and purified by silica gel chromatography to give the desired product as a white solid: mp 205-206 °C; ¹H NMR (CDCl₃) δ 0.80-0.95 (m, 6H), 1.10-1.70 (m, 7H), 2.15 (m, 1H), 3.02 (d, J = 15.3 Hz, 2H), 3.15 (d, J = 15.1 Hz, 2H), 3.96 (s, br, 2H), 4.14 (d, J = 7.8 Hz, 1H), 5.51 (s, 1H), 5.94 (d, J = 2.2, 1H), 6.54 (dd, J = 8.5, 2.2 Hz, 1H), 7.28-7.50 (m, 6H), 7.87 (d, J = 8.5 Hz, 1H). MS (FAB): m/z 388 (M+H).

Example 68 (Compound No. 290)

244



No. 290

Following a procedure similar to the one described in Example 89, *infra* (see Compound No. 121), the title compound was prepared and purified as a colorless

solid: mp = 96-98 °C, ¹H NMR (CDCl₃) δ 0.92 (t, J = 7

Hz, 6H), 1.03-1.70 (m, 11H), 2.21 (t, J = 8 Hz, 1H),


3.09 (AB, J_{AB} = 18 Hz, DV = 38 Hz, 2H), 3.96 (bs, 2H),

4.14 (d, J = 7 Hz, 1H), 5.51 (s, 1H), 5.94 (s, 1H),

6.56 (d, J = 9 Hz, 1H), 7.41-7.53 (m, 6H), 7.87 (d, J =

8 Hz, 1H); FAB/MS m/z 416 (M+H).

Example 69




No. 291

Following a procedure similar to the one described in Example 86, *infra* (see Compound No. 118), the title compound was prepared and purified as a colorless

solid: ¹H NMR (CDCl₃) δ 0.91 (t, J = 7 Hz, 6H), 1.02-


1.52 (m, 11H), 1.60-1.70 (m, 1H), 2.23 (t, J = 8 Hz,

Example 70 (Compound No. 292)

No. 292

Following a procedure similar to the one described in Example 89, *infra* (see Compound No. 121), the title compound was prepared and purified as a colorless

solid: mp = 111-112.5°C, ¹H NMR (CDCl₃) δ 0.90 (t, J = 8 Hz, 6H), 1.03-1.50 (m, 10H), 1.55-1.70 (m, 2H), 2.18 (t, J = 12 Hz, 2H), 3.07 (AB, J_A = 15 Hz, Dv = 45 Hz, 2H), 4.09 (bs, 2H), 5.49 (s, 1H), 5.91 (s, 1H), 6.55 (d, J = 9 Hz, 1H), 7.10 (t, J = 7 Hz, 2H), 7.46 (t, J = 6 Hz, 2H), 7.87 (d, J = 9 Hz, 1H).

Example 71 (Compound No. 293)

No. 293

During the preparation of Compound No. 290 from Compound No. 291 using BBr₃, the title compound was

isolated: ¹H NMR (CDCl₃) δ 0.85 (t, J = 6 Hz, 6H), 0.98-1.60 (m, 10H), 1.50-1.66 (m, 2H), 2.16 (t, J = 8 Hz, 1H), 3.04 (AB, J_A = 15 Hz, Dv = 41 Hz, 2H), 4.08 (s, 1H), 4.12 (s, 1H), 5.44 (s, 1H), 5.84 (s, 1H), 6.42 (d, J = 9 Hz, 1H), 7.12 (d, J = 8 Hz, 2H), 7.16-7.26 (m, 10H), 7.83 (d, J = 8 Hz, 1H); ESMS m/z 512 (M+Li).

Example 72 (Compound No. 294)

Following a procedure similar to the one described in Example 60 (Compound No. 104), the title compound was prepared and purified as a colorless solid: ¹H NMR (CDCl₃) δ 0.90 (t, J = 6 Hz, 6H), 1.05-1.54 (m, 9H), 1.60-1.70 (m, 1H), 2.24 (t, J = 8 Hz, 1H), 2.80 (s, 6H), 3.05 (AB, J_A = 15 Hz, Dv = 42 Hz, 2H), 4.05-4.18 (m, 2H), 5.53 (s, 1H), 5.93 (s, 1H), 6.94 (d, J = 9 Hz, 1H), 7.27-7.42 (m, 4H), 7.45 (d, J = 8 Hz, 2H), 7.87 (d, J = 9 Hz, 1H); ESMS m/z 444 (M+H).

Structures of the compounds of Examples 33 to 72 are shown in Tables 3 and 3A.

Examples 73-79, 87, 88, and 91-102

Using in each instance a method generally described in those of Examples 1 to 72 appropriate to the substituents to be introduced, compounds were prepared having the structures set forth in Table 3. The starting materials illustrated in the reaction schemes shown above were varied in accordance with principles of organic synthesis well known to the art to introduce the indicated substituents in the 4- and 5- positions (R', R', R') and in the indicated position on the benzo ring (R').

Structures of the the compounds produced in Examples 73-102 are set forth in Tables 3 and 3A.

Examples 80-84

- 5 Preparation of 115, 116, 111, 113
Preparation of 4-chloro-3-[4-methoxy-phenylmethyl]-nitrobenzene.
In a 500 ml 2-necked rb flask weigh out 68.3 gms phosphorus pentachloride (0.328 mole 1.1 eq). Add 50 mls chlorobenzene. Slowly add 60 gms 2-chloro-5-nitrobenzoic acid (0.298 mole). Stir at room temp overnight under N₂ then heat 1 hr at 50C.
Remove chlorobenzene by high vacuum. Wash residue with hexane. Dry wt=55.5 gms.
- 15 In the same rb flask, dissolve acid chloride (55.5 g 0.25 mole) from above with 100 mls anisole (about 3.4 eq). Chill solution with ice bath while purging with N₂. Slowly add 40.3g aluminum chloride (1.2 eq 0.3 mole). Stir under N₂ for 24 hrs.
- 20 After 24 hrs, the solution was poured into 300 mls 1N HCl soln. (cold). Stir this for 15 min. Extract several times with diethyl ether. Extract organic layer once with 2% aqueous NaOH then twice with water. Dry organic layer with MgSO₄, dry on vac line. Solid is washed well with ether and then ethanol before drying. Wt=34.57g (mixture of meta, ortho and para).

Elemental	theory	found
C	57.65	57.45
H	3.46	5.51
N	4.8	4.8
Cl	12.15	12.16

249

With the next step of the reduction of the ketone with trifluoromethane sulfonic acid and triethyl silane, crystallization with ethyl acetate/hexane affords pure 4-chloro-3-[4-methoxy-phenylmethyl]-nitrobenzene.

- 5 4-Chloro-3-[4-methoxy-phenylmethyl]-nitrobenzene was then reacted as specified in the synthesis of 117 and 118 from 2-chloro-4-nitrophenylmethane. From these procedures 115 and 116 can be synthesized. Compounds 111 and 113 can be synthesized from the procedure used to prepare compound 121.
- 10

Compound 114 can be prepared by reaction of 116 with ethyl mercaptan and aluminum trichloride.

Examples 85 and 86

- 15 Preparation of 117 and 118

2-Chloro-4-nitrobenzophenone is reduced with triethylsilane and trifluoromethane sulfonic acid to 2-chloro-4-nitrodiphenylmethane 32. Reaction of 32 with lithium sulfide followed by reacting the resulting sulfide with mesylate IV gives sulfide-aldehyde XXIII. Oxidation of XXIII with 2 equivalents of MCPBA yields sulfone-aldehyde XXIII. Oxidation of XXIII with 2 equivalents of MCPBA yields sulfone-aldehyde XXIV (see Scheme 5).

- 25 The sulfone-aldehyde (31.8 g) was dissolved in ethanol/toluene and placed in a parr reactor with 100 ml toluene and 100 ml of ethanol and 3.2 g of 10% Pd/C and heated to 55 C and 100 psi of hydrogen gas for 14 hours. The reaction was then filtered to remove the catalyst. The amine product (.076 moles, 29.5 g) from this reaction was then reacted with benzyl chloroformate (27.4g) in toluene in the presence of 35 g of potassium carbonate and stirred at room
- 30

250

temperature overnight. After work up by extraction with water, the CBZ protected amine product was further purified by precipitation from toluene/hexane.

The CBZ protected amine product was then reacted with 3 equivalents of potassium t-butoxide in THF at 0 C to yield compounds 117 and 118 which were separated by silica gel column chromatography.

Examples 89 and 90

10 Preparation of 121 or 122

Compound 118 (.013 moles, 6.79g) is dissolved in 135 ml of dry chloroform and cooled to -78 C, next 1.85 ml of boron tribromide (4.9 g) was added and the reaction is allowed to warm to room temperature.

15 Reaction is complete after 1.5 hours. The reaction is quenched by addition of 10% potassium carbonate at 0 C and extract with ether. Removal of ether yields compound 121. A similar procedure can be used to produce 122 from 117.

20

Examples 93-96

Compounds 126, 127, 128 and 129 as set forth in Table 3 were prepared substantially in the manner described above for compounds 115, 116, 111 and 113, respectively, except that fluorobenzene was used as a starting material in place of anisole.

25

TABLE 3
Specific compounds (#102-111, 113-130, 132-134, 136, 138, 142-144, 262-296)






TABLE 3A
Bridged Benzothiephenes (#101, 112, 131, 135, 137, 139-141)


38	74	Et-	n-Bu-	HO-	H-	O-F-Ph-	H-	7-CH ₃ O-
42	81	Et-	n-Bu-	HO-	H-	P-F-Ph-	H-	7-CH ₃ S-
45	85	Et-	n-Bu-	HO-	H-	P-F-Ph-	H-	7-CH ₃ -
45	84	Et-	n-Bu-	H-	HO-	H-	P-F-Ph-	7-CH ₃ -
44	83	Et-	n-Bu-	HO-	H-	P-F-Ph-	H-	7-(N)- morpholine.
43	82	Et-	n-Bu-	HO-	H-	P-F-Ph-	H-	7-(N)- pyrrolid- dine
64	286	Et-	Et-	HO-	H-	Ph-	H-	7-NH-CBZ
65	287	Et-	Et-	HO-	F-	Ph-	H-	7-NH ₂ -
66	288	CH ₃ -	CH ₃ -	HO-	F-	Ph-	H-	7-NH ₂ -
67	289	n- C ₃ H ₇ -	n- C ₃ H ₇ -	HO-	H-	Ph-	H-	7-NH ₂ -
68	290	n-Bu-	n-Bu-	HO-	H-	Ph-	H-	7-NH ₂ -
69	291	n-Bu-	n-Bu-	HO-	H-	Ph-	H-	7-NH-CBZ
70	292	n-Bu-	n-Bu-	HO-	H-	P-F-Ph-	H-	7-NH ₂ -
71	293	n-Bu-	n-Bu-	HO-	H-	Ph-	H-	7-PnCH ₂ N-
72	294	n-Bu-	n-Bu-	HO-	H-	Ph-	H-	7-(CH ₃) ₂ N-
63	295	Et-	n-Bu-	HO-	H-	P-I- (C ₂ H ₅ O) ₂ - Ph-	H-	7-NH ₂ -
102	296	Et-	n-Bu-	HO-	H-	I ⁻ , P- (CH ₃) ₃ N ⁺ (C ₂ H ₅ O) ₂ Ph-	H-	7-NH ₂ -




CPD #101 (Ex. 59)




CPD #112 (Ex. 53)




CPD #131 (Ex. 56)




CPD #135 (Ex. 53)




CPD #137 (Ex. 57)



CPD #139 (Ex. 58)



CPD #140 (Ex. 51)




(Ex. 50)

Examples 104-231

Using in each instance a method generally described in those of Examples 1 to 72 appropriate to the substituents to be introduced, including where necessary other common synthesis expedients well known to the art, compounds are prepared having the structures set forth in Table 4. The starting materials illustrated in the reaction schemes shown above are varied in accordance with principles of organic synthesis well known to the art in order to introduce the indicated substituents in the 4- and 5-positions (R^1 , R^1 , R^1 , R^1) and in the indicated position on the benzo ring (R^1).

TABLE 4
Alternative compounds #1 (302-312, 314-430)



Comp#	R^1	$(R^2)_q$
302	p-F-Ph-	7-(1-azetidine)
303	p-F-Ph-	7-EtS-
304	p-F-Ph-	7-CH ₃ (O)-
305	p-F-Ph-	7-CH ₃ (O) ₂ -
306	p-F-Ph-	7-PhS-
307	p-F-Ph-	7-CH ₃ S- 9-CH ₃ S-
308	p-F-Ph-	7-CH ₃ O- 9-CH ₃ O-
309	p-F-Ph-	7-Et-
310	p-F-Ph-	7-IPt-
311	p-F-Ph-	7-t-Bu-
312	p-F-Ph-	7-(1-pyrazole)-
314	m-CH ₃ O-Ph	7-(1-azetidine)
315	m-CH ₃ O-Ph-	7-(1-azetidine)
316	m-CH ₃ O-Ph-	7-EtS-
317	m-CH ₃ O-Ph-	7-CH ₃ (O)-
318	m-CH ₃ O-Ph-	7-CH ₃ (O) ₂ -
319	m-CH ₃ O-Ph-	7-PhS-


320	m-CH ₃ O-Ph	7-CH ₃ S- 9-CH ₃ S-	337	p-F-Ph-	Ph-
321	m-CH ₃ O-Ph	7-CH ₃ O- 9-CH ₃ O-	338	p-F-Ph-	7-CH ₃ C(=CH ₂)-
322	m-CH ₃ O-Ph	7-Et-	339	p-F-Ph-	7-cyclopropyl
323	m-CH ₃ O-Ph	7-IPr-	340	p-F-Ph-	7-(CH ₃) ₂ NH-
324	m-CH ₃ O-Ph	7-t-Bu-	341	p-F-Ph-	7-(N)-azetidide 9-CH ₃ S-
325	p-F-Ph-	6-CH ₃ O- 7-CH ₃ O- 8-CH ₃ O-	342	p-F-Ph-	7-(N-pyrrolidine) 9-CH ₃ S-
326	p-F-Ph-	7-(1-azetidide) 9-CH ₃ -	343	p-F-Ph-	7-(CH ₃) ₂ N- 9-CH ₃ S-
327	p-F-Ph-	7-EtS- 9-CH ₃ -	344	m-CH ₃ O-Ph-	7-(1-pyrazole)
328	p-F-Ph-	7-CH ₃ S(O)- 9-CH ₃ -	345	m-CH ₃ O-Ph-	7-(N,N')-methylpiperazine
329	p-F-Ph-	7-CH ₃ S(O) ₂ - 9-CH ₃ -	346	m-CH ₃ O-Ph-	Ph-
330	p-F-Ph-	7-PhS- 9-CH ₃ -	347	m-CH ₃ O-Ph-	7-CH ₃ C(=CH ₂)-
331	p-F-Ph-	7-CH ₃ S- 9-CH ₃ -	348	m-CH ₃ O-Ph-	7-cyclopropyl
332	p-F-Ph-	7-CH ₃ O- 9-CH ₃ -	349	m-CH ₃ O-Ph-	7-(CH ₃) ₂ NH-
333	p-F-Ph-	7-CH ₃ - 9-CH ₃ -	350	m-CH ₃ O-Ph-	7-(N)-azetidide 9-CH ₃ S-
334	p-F-Ph-	7-CH ₃ O- 9-CH ₃ O-	351	m-CH ₃ O-Ph-	7-(N-pyrrolidine)- 9-CH ₃ S-
335	p-F-Ph-	7-(1-pyrazole)	352	m-CH ₃ O-Ph-	7-(CH ₃) ₂ N- 9-CH ₃ S-
336	p-F-Ph-	7-(N,N')-methylpiperazine	353	m-CH ₃ O-Ph-	6-CH ₃ O- 7-CH ₃ O- 8-CH ₃ O-
			354	m-CH ₃ O-Ph-	7-(1-azetidide) 9-CH ₃ -

355	m-CH ₃ O-Ph-	7-Ets- 9-CH ₃ -	375	thien-2-yl	7-CH ₃ S-
356	m-CH ₃ O-Ph-	7-CH ₃ S(O)- 9-CH ₃ -	376	thien-2-yl	7-(1-azetidine)
357	m-CH ₃ O-Ph-	7-CH ₃ S(O)2- 9-CH ₃ -	377	thien-2-yl	7-Me-
358	m-CH ₃ O-Ph-	7-PhS- 9-CH ₃ -	378	5-Cl-thien-2-yl	7-(1-azetidine)
359	m-CH ₃ O-Ph-	7-CH ₃ S- 9-CH ₃ -	379	5-Cl-thien-2-yl	7-(1-azetidine)
360	m-CH ₃ O-Ph-	7-CH ₃ O- 9-CH ₃ -	380	5-Cl-thien-2-yl	7-Ets-
361	m-CH ₃ O-Ph-	7-CH ₃ - 9-CH ₃ -	381	5-Cl-thien-2-yl	7-CH ₃ S(O)-
362	m-CH ₃ O-Ph-	7-CH ₃ O- 9-CH ₃ O-	382	5-Cl-thien-2-yl	7-CH ₃ S(O)2-
			383	5-Cl-thien-2-yl	7-PhS-
			384	5-Cl-thien-2-yl	7-CH ₃ S- 9-CH ₃ S-
			385	5-Cl-thien-2-yl	7-CH ₃ O- 9-CH ₃ O-
363	thien-2-yl	7-(1-azetidine)	386	5-Cl-thien-2-yl	7-Et-
364	thien-2-yl	7-Ets-	387	5-Cl-thien-2-yl	7-IPr-
365	thien-2-yl	7-CH ₃ S(O)-	388	5-Cl-thien-2-yl	7-t-Bu-
366	thien-2-yl	7-CH ₃ S(O)2-	389	5-Cl-thien-2-yl	7-CH ₃ O-
367	thien-2-yl	7-PhS-	390	5-Cl-thien-2-yl	7-CH ₃ S-
368	thien-2-yl	7-CH ₃ S- 9-CH ₃ S-	391	5-Cl-thien-2-yl	7-Me
369	thien-2-yl	7-CH ₃ O- 9-CH ₃ O-	392	thien-2-yl	7-(1-azetidine) 9-CH ₃ -
370	thien-2-yl	7-Et-	393	thien-2-yl	7-Ets- 9-CH ₃ -
371	thien-2-yl	7-IPr-	394	thien-2-yl	7-CH ₃ S(O)- 9-CH ₃ -
372	thien-2-yl	7-t-Bu-	395	thien-2-yl	7-CH ₃ S(O)2- 9-CH ₃ -
373	thien-2-yl	7-(1-pyrrole)-			
374	thien-2-yl	7-CH ₃ O-			

396	thien-2-yl	7-PhS- 9-CH ₃ -	417	5-Cl-thien-2-yl	7-(N)-azetidine 9-CH ₃ S-
397	thien-2-yl	7-CH ₃ S- 9-CH ₃ -	418	5-Cl-thien-2-yl	7-(N-pyrrolidine)- 9-CH ₃ S-
398	thien-2-yl	7-CH ₃ O- 9-CH ₃ -	419	5-Cl-thien-2-yl	7-(CH ₃) ₂ N- 9-CH ₃ S-
399	thien-2-yl	7-CH ₃ - 9-CH ₃ -	420	5-Cl-thien-2-yl	7-(1-azetidine) 9-CH ₃ -
400	thien-2-yl	7-CH ₃ O- 9-CH ₃ O-	421	5-Cl-thien-2-yl	7-EtS- 9-CH ₃ -
401	thien-2-yl	7-(1-pyrazole)	422	5-Cl-thien-2-yl	7-CH ₃ S(O)- 9-CH ₃ -
402	thien-2-yl	7-(N,N')-methylpiperazine	423	5-Cl-thien-2-yl	7-CH ₃ S(O) ₂ - 9-CH ₃ -
403	thien-2-yl	Ph-	424	5-Cl-thien-2-yl	7-PhS- 9-CH ₃ -
404	thien-2-yl	7-CH ₃ C(=CH ₂)-	425	5-Cl-thien-2-yl	7-CH ₃ S- 9-CH ₃ -
405	thien-2-yl	7-cyclopropyl	426	5-Cl-thien-2-yl	7-CH ₃ O- 9-CH ₃ -
406	thien-2-yl	7-(CH ₃) ₂ NH -	427	5-Cl-thien-2-yl	7-CH ₃ - 9-CH ₃ -
407	thien-2-yl	7-(N)-azetidine 9-CH ₃ S-	428	5-Cl-thien-2-yl	7-CH ₃ O- 9-CH ₃ O-
408	thien-2-yl	7-(N-pyrrolidine) 9-CH ₃ S-	429	thien-2-yl	6-CH ₃ O- 7-CH ₃ O- 8-CH ₃ O-
409	thien-2-yl	7-(CH ₃) ₂ N- 9-CH ₃ S-	430	5-Cl-thien-2-yl	6-CH ₃ O- 7-CH ₃ O- 8-CH ₃ O-
411	5-Cl-thien-2-yl	7-(1-pyrazole)			
412	5-Cl-thien-2-yl	7-(N,N')-methylpiperazine			
413	5-Cl-thien-2-yl	Ph-			
414	5-Cl-thien-2-yl	7-CH ₃ C(=CH ₂)-			
415	5-Cl-thien-2-yl	7-cyclopropyl			
416	5-Cl-thien-2-yl	7-(CH ₃) ₂ NH -			

Example 232-1394

Using in each instance a method generally described in those of Examples 1 to 72 appropriate to the substituents to be introduced, including where necessary other common synthesis expedients well known to the art, compounds are prepared having the structures set forth in Table 1. The starting materials illustrated in the reaction schemes shown above are varied in accordance with principles of organic synthesis well known to the art in order to introduce the indicated substituents in the 4- and 5-positions (R^1 , R^1 , R^1) and in the indicated position on the benzo ring (R^1).

Example 1395**Dibutyl 4-fluorobenzene dialdehyde****Step 1: Preparation of dibutyl 4-fluoro benzene dialdehyde**

To a stirred solution of 17.5 g (123 mmol) of 2,5-difluorobenzaldehyde (Aldrich) in 615 mL of DMSO at ambient temperature was added 6.2 g (135 mmol) of lithium sulfide (Aldrich). The dark red solution was stirred at 75 C for 1.5 hours, or until the starting material was completely consumed, and then 34 g (135 mmol) of dibutyl mesylate aldehyde was added at about 50 C. The reaction mixture was stirred at 75 C for three hours or until the reaction was completed. The cooled solution was poured into water and extracted

with ethyl acetate. The combined extracts were washed with water several times, dried (MgSO₄) and concentrated in vacuo. Silica gel chromatographic purification of the crude product gave 23.6 g (59%) of fluorobenzene dialdehyde as a yellow oil: ¹H NMR (CDCl₃) δ 0.87 (t, J = 7.05 Hz, 6H, 1.0-1.4 (m, 8H), 1.5-1.78 (m, 4H), 3.09 (s, 2H), 7.2-7.35 (m, 1H), 7.5-7.6 (m, 2H), 9.43 (s, 1H), 10.50 (d, J = 2.62 Hz, 1H).

Step 2: Preparation of dibutyl 4-fluorobenzyl alcohol
To a solution of 22.6 g (69.8 mmol) of the dialdehyde obtained from Step 1 in 650 mL of THF at -60 C was added 69.8 mL (69.8 mmol) of DIBAL (1M in THF) via a syringe. The reaction mixture was stirred at -40 C for 20 hours. To the cooled solution at -40 C was added sufficient amount of ethyl acetate to quench the excess of DIBAL, followed by 3 N HCl. The mixture was extracted with ethyl acetate, washed with water, dried (MgSO₄), and concentrated in vacuo. Silica gel chromatographic purification of the crude product gave 13.5 g (58%) of recovered starting material, and 8.1 g (36%) of the desired fluorobenzyl alcohol as a colorless oil: ¹H NMR (CDCl₃) δ 0.88 (t, J = 7.05 Hz, 6H), 1.0-1.4 (m, 8H), 1.5-1.72 (m, 4H), 1.94 (br s, 1H), 3.03 (s, 2H), 4.79 (s, 2H), 6.96 (dt, J = 8.46, 3.02 Hz, 1H), 7.20 (dd, J = 9.47, 2.82 Hz, 1H), 7.42 (dd, J = 8.67, 5.64, 1H), 9.40 (s, 1H).

Step 3: Preparation of dibutyl 4-fluorobenzyl bromide
To a solution of 8.1 g (25 mmol) of benzyl alcohol obtained from Step 2 in 100 mL of DMF at -40 C was added 47 g (50 mmol) of bromotriphenylphosphonium bromide (Aldrich). The resulting solution was stirred cold for 30 min, then was allowed to warm to 0 C. To the mixture was added 10% solution of sodium sulfite and ethyl acetate. The extract was washed a few times with water, dried (MgSO₄), and concentrated in vacuo.


The mixture was stirred in small amount of ethyl acetate/hexane mixture (1:4 ratio) and filtered through a pad of silica gel, eluting with same solvent mixture. The combined filtrate was concentrated in vacuo to give 9.5 g (98%) of the desired product as a colorless oil:

¹H NMR (CDCl₃) δ 0.88 (t, J = 7.05 Hz, 6H), 1.0-1.4 (m, 8H), 1.55-1.78 (m, 4H), 3.11 (s, 2H), 4.67 (s, 2H), 7.02 (dt, J = 8.46, 3.02 Hz, 1H), 7.15 (dd, J = 9.47, 2.82 Hz, 1H), 7.46 (dd, J = 8.67, 5.64, 1H), 9.45 (s, 1H).


Step 4: Preparation of sulfonyl 4-fluorobenzyl bromide

To a solution of 8.5 g (25 mmol) of sulfide obtained from Step 3 in 200 mL of CH₂Cl₂ at 0 °C was added 15.9 g (60 mmol) of MCPBA (64% peracid). The resulting solution was stirred cold for 10 min, then was allowed to stirred ambient temperature for 5 hours. To the mixture was added 10% solution of sodium sulfite and ethyl acetate. The extract was washed several times with saturated Na₂CO₃, dried (MgSO₄), and concentrated in vacuo to give 10.2 g (98%) of the desired product as a colorless oil: ¹H NMR (CDCl₃) δ 0.91 (t, J = 7.05 Hz, 6H), 1.03-1.4 (m, 8H), 1.65-1.82 (m, 2H), 1.90-2.05 (m, 2H), 3.54 (s, 2H), 5.01 (s, 2H), 7.04-7.23 (m, 1H), 7.30 (dd, J = 8.87, 2.42 Hz, 1H), 8.03 (dd, J = 8.86, 5.64, 1H), 9.49 (s, 1H).

Example 1396



Generic Scheme X



Generic Scheme X: The nucleophilic substitution of an appropriately substituted 2-fluorobenzaldehyde with lithium sulfide or other nucleophilic sulfide anion in polar solvent (such as DMF, DMA, DMSO, etc), followed by the addition of dialkyl mesylate aldehyde (X), provided a dialkyl benzene dialdehyde Y. DIBAL reduction of the dialdehyde at low temperature yielded benzyl alcohol monaldehyde Z. Conversion of benzyl alcohol to benzyl bromide, followed by oxidation of sulfide to sulfone yielded the key intermediate W.

Preparation of N-propylsulfonic acid


To a solution of 51 mg (111 μm) Compound X in ethanol (400 μl) was added 1,3 propane sultone (19.5 μl , 222 μm). The reaction was stirred in a sealed vial at 55 °C for 25 hr. Sample was concentrated under a nitrogen stream and purified by reversed phase chromatography using acetonitrile/water as eluent (30-45%) and afforded the desired material as an off-white solid (28.4 mg, 44%): $^1\text{H NMR}$ (CDCl_3) δ 0.82-0.96 (m, 6H), 1.11-1.52 (m of m, 10H), 1.58-1.72 (m, 1H), 2.08-2.21 (m, 1H), 2.35-2.50 (m, 2H), 2.93 (s, 6H), 3.02-3.22 (m of m, 5H), 3.58-3.76 (m, 2H), 4.15 (s, 1H), 5.51 (s, 1H), 6.45-6.58 (m, 1H), 6.92-7.02 (m, 1H), 7.35-7.41 (m, 1H), 7.41-7.51 (m, 2H), 8.08 (d, J = 8.1 Hz, 1H), 8.12-8.25 (m, 1H); MS ES- M-H m/z 579.

Example 1397

The 7-fluoro, 9-fluoro and 7,9-difluoro analogs of benzotheipine compounds of this invention can be reacted with sulfur and nitrogen nucleophiles to give the corresponding sulfur and nitrogen substituted analogs. The following example demonstrates the

synthesis of these analogs.

3,3-Dibutyl-5a-(4'-fluorophenyl)-4a-hydroxy-7-methylthio-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide.



5

A mixture of 0.4 g of 3,3-dibutyl-7-fluoro-5a-(4'-fluorophenyl)-4a-hydroxy-2,3,4,5-

tetrahydrobenzothiepine-1,1-dioxide, prepared by

previously described method, 0.12 g of sodium

methanethiolate and 20 ml of DMF was stirred at 50 C

for 3 days. An additional 0.1 g of sodium

methanethiolate was added to the reaction mixture and

the mixture was stirred for additional 20 h at 50 C

then was concentrated in vacuo. The residue was

trituated with water and extracte wiith ether. The

ether extract was dried over MgSO₄ and concentrated in

vacuo to 0.44 g of an oil. Purification by HPLC (10%

EtOAc in hexane) gave 0.26 g of needles, mp 164-165.5

°C.

20

3,3-Dibutyl-9-dimethylamino-7-fluoro-5a-(4'-


fluorophenyl)-4a-hydroxy-2,3,4,5-

tetrahydrobenzothiepine-1,1-dioxide and 7,9-

Bis(dimethylamino)-3,3-dibutyl-5a-(4'-fluorophenyl)-4a-





hydroxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide.

25




A solution of 0.105 g of 3,3-dibutyl-7,9-difluoro-5a-(4'-fluorophenyl)-4a-hydroxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide, prepared by the method described previously, in 20 ml of 2 N dimethylamine in THF was heated at 160 °C in a sealed Parr reactor overnight. The reaction mixture was cooled and concentrated in vacuo. The residue was triturated with 25 ml of water and extracted with ether. The ether extract was dried over MgSO_4 and concentrated in vacuo. The residue was purified by HPLC (10% EtOAc in hexane) to give 35 mg of an earlier fraction which was identified as 3,3-dibutyl-9-dimethylamino-7-fluoro-5a-(4'-fluorophenyl)-4a-hydroxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide, MS (CI) m/e 480 ($M^+ + 1$), and 29 mg of a later fraction which was identified as 7,9-bis(dimethylamino)-3,3-dibutyl-5a-(4'-fluorophenyl)-4a-hydroxy-2,3,4,5-tetrahydrobenzothiepine-1,1-dioxide, MS (CI) m/e 505 ($M^+ + 1$).

The compounds of this invention can also be synthesized using cyclic sulfate (A, below) as the reagent as shown in the following scheme. The following example describes a procedure for using the cyclic sulfate as the reagent.



Dibutyl cyclic sulfite:




A solution of 2,2-dibutyl-1,3-propanediol (103g, 0.548 mol) and triethylamine (221g, 2.19 mol) in anhydrous methylene chloride (500 ml) and was stirred at 0 degrees C under nitrogen. To the mixture, thionyl chloride (97.8 g, 0.82 mol) was added dropwise and within 5 min the solution turned yellow and then turned black when the addition was completed within half an hour. The reaction mixture was stirred for 3 hrs. GC showed that there was no starting material left. The mixture was washed with ice water twice then with brine twice. The organic phase was dried over magnesium sulfate and concentrated under vacuum to give the cyclic sulfite 128 g (100%) as a black oil. Mass spectrum (MS) was consistent with the product.

To a solution of the above compound (127.5g, 0.54 mol) in 600 ml acetonitrile and 500 ml of water cooled in an ice bath under nitrogen was added ruthenium(III) chloride (1 g) and sodium periodate (233 g, 1.08 mol). The reaction was stirred overnight and the color of the solution turned black. GC showed that there was no starting material left. The mixture was extracted with 300 ml of ether and the ether extract was washed three times with brine. The organic phase was dried over magnesium sulfate and passed through celite. The filtrate was concentrated under vacuum and gave the cyclic sulfate 133 g (97.8%) as an oil. Proton, carbon NMR and MS were consistent with the product.

277


2-[(2-(4'-Fluorobenzyl)-4-methylphenylthio)methyl]-2-butylhexanol:



Sodium hydride (60% oil dispersion), 0.27 g (6.68 mmole), was washed with hexane and the hexane wash was decanted. To the washed sodium hydride was added 20 ml of 2-methoxyethyl ether (diglyme) and the mixture was cooled in an ice bath. A solution of 1.55 g (6.68 mmole) of 2-(4'-fluorobenzyl)-4-methylbenzenethiol in 10 ml of 2-methoxyethyl ether was added dropwise to the reaction mixture in 15 min. A mixture of 2.17 g (8.68 mmole) of the dibutyl cyclic sulfate in 10 ml of 2-methoxyethyl ether was added once and stirred for 30 min at 0 C then at room temperature for 1 hr under nitrogen. GC showed that there was no thiol left. The solvent was evaporated and triturated with water then was extracted with ether twice. The water layer was separated, treated with 20 ml of 10% NaOH then was boiled for 30 min and cooled, acidified with 6N HCl and boiled for 10 min. The reaction mixture was washed and extracted with ether. The organic layer was washed successively with water and brine, dried over magnesium sulfate and concentrated under vacuum to give 2.47 g (92.5%) of an oil. Proton NMR, ¹³C NMR and MS were consistent with the product.


278

2-((2-(4'-Fluorobenzyl)-6-methylphenyl)ethio)methyl]-2-butylhexanol




To a solution of the above product (2 g, 4.9 mmol) in 40 ml methylene chloride cooled in an ice bath under nitrogen was added pyridinium chlorochromate (2.18 g, 9.9 mmol) at once. The reaction was stirred with 3 hrs and filtered through a bed of silica gel. The filtrate was concentrated under vacuum to give 1.39 g (70%) of an oil. Proton, carbon NMR and MS were consistent with the product.

2-((2-(4'-Fluorobenzyl)-4-methylphenyl)sulfonyl)methyl]-2-butylhexanol



To a solution of the above product (0.44 g, 1.1 mmole) in 20 ml methylene chloride solution cooled in an ice bath under nitrogen was added 70% m-chloroperbenzoic acid (0.54 g, 2.2 mmol) at once. The reaction mixture was stirred for 18 hrs and filtered. The filtrate was washed successively with 10% NaOH (3x), water and brine, dried over magnesium sulfate and concentrated under vacuum to give 0.42 g (90%) of an oil. Proton, carbon NMR and MS were consistent with the product.

3,3-Dibutyl-7-methyl-5a-(4'-fluorophenyl)-6a-hydroxy-2,3,4,5-tetrahydrobenzothioflavino-1,1-dioxide




A mixture of 0.37 g (0.85 mmol) of the above product in 30 ml of anhydrous THF was stirred at 0 °C. Then potassium t-butoxide (102 mg, 0.85 mmol) was added. After 3 hrs, TLC showed that there was a product and some starting material left. The crude reaction mixture was acidified with 10% HCl and extracted with ether. The ether extract was washed successively with water and brine, dried with MgSO₄ and concentrated under vacuum. The residue was purified by HPLC (10% EtOAc-Hexane). The first fraction was 0.1 g of starting material as an oil and the second fraction was a white solid, 0.27 g (75%). Proton NMR and carbon NMR were consistent with the desired product. Mass spectrum (CI)

also confirmed the product, m/e 433 (M⁺ 1).

Example 1398

Step 1



$C_{14}H_9ClNO_4$, fw=291.69

In an inert atmosphere, weigh out 68.3 gms phosphorus pentachloride (0.328mole Aldrich 15,777-5) into a 2-necked 500ml round bottom flask. Fit flask with a N₂ inlet adapter and suba seal. Remove from inert atmosphere and begin N₂ purge. Add 50mls anhydrous chlorobenzene (Aldrich 28,451-3) to the PCl₅ via syringe and begin stirring with magnetic stir bar.


- 5
 - 10
 - 15
 - 20
 - 25
- Weigh out 60 gms 2-chloro-5-nitrobenzoic acid (0.298 mole Aldrich 12,511-3). Slowly add to the chlorobenzene solution while under N₂ purge. Stir at room temperature overnight. After stirring at room temperature for ~20hrs, place in oil bath and heat at 50C for 1hr. Remove chlorobenzene by high vacuum. Wash residue with anhydrous hexane. Dry acid chloride wt=61.95gms. Store in inert and dry atmosphere.
- In inert atmosphere, dissolve acid chloride with 105mls anhydrous anisole (0.97 mole Aldrich 29,629-5). Place solution in a 2-necked 500ml round bottom flask.

- 30
- Weigh out 45.1gms aluminum chloride (0.34 moles Aldrich 29,471-3) and place in a solid addition funnel. Fit reaction flask with addition funnel and a N₂ inlet adapter. Remove from inert atmosphere. Chill reaction solution with ice bath and begin N₂ purge. Slowly add

AlCl₃ to chilled solution. After addition is complete, allow to warm to room temperature. Stir overnight.

Quench reaction by pouring into a solution of 300 mls 1N HCl and ice. Stir 15 min. Extract twice with ether. Combine organic layers and extract twice with 2% NaOH, then twice with deionized H₂O. Dry with MgSO₄, filter and rotovap to dryness. Remove anisole by high vacuum. Crystallize product from 90% ethanol 10% ethyl acetate. Dry on vacuum line. Wt=35.2gms. Yield 41%. Obtain NMR and mass spec (m/z=292).

Step 2



Dissolve 38.10gms (0.131 moles) of the benzophenone from step 1 in 250mls anhydrous methylene chloride. Place in a 3 liter flask fitted with N₂ inlet, addition funnel and stopper. Stir with magnetic stir bar. Chill solution with ice bath.

Prepare a solution of 39.32 gms trifluoromethane sulfonic acid (0.262 mole Aldrich 15,853-4) and 170 mls anhydrous methylene chloride. Place in addition funnel and add dropwise to chilled solution under N₂. Stir 5 minutes after addition is complete.

Prepare a solution of 22.85 gms triethyl silane (0.197mole Aldrich 23,019-7) and 170mls anhydrous methylene chloride. Place in addition funnel and add dropwise to chilled solution under N₂. Stir 5 minutes after addition is complete.

Prepare a second solution of 39.32 gms trifluoromethane sulfonic acid and 170mls anhydrous


283

methylene chloride. Place in addition funnel and add dropwise to chilled solution under N₂. Stir 5 minutes after addition is complete.

Prepare a second solution of 22.85 gms triethyl silane and 170mls anhydrous methylene chloride. Place in addition funnel and add dropwise to chilled solution under N₂. After all additions are made allow to slowly warm to room temperature overnight. Stir under N₂ overnight.

Prepare 1300 mls saturated NaHCO₃ in a 4 liter beaker. Chill with ice bath. While stirring vigorously, slowly add reaction mixture. Stir at chilled temperature for 30 min. Pour into a separatory funnel and allow separation. Remove organic layer and extract aqueous layer 2 times with methylene chloride. Dry organic layers with MgSO₄. Crystallize from ethanol. Dry on vacuum line. Dry wt=28.8gms. Confirm by NMR and mass spec (m/z=278).

Step 3



Dissolve 10.12 gms (0.035 moles) of product 2 with 200 mls anhydrous DMSO. Place in a 500 ml round bottom flask with magnetic stir bar. Fit flask with water condenser, N₂ inlet, and stopper. Add 1.84 gms Li₂S (0.040 moles Aldrich 21,324-1). Place flask in oil bath and heat at 75°C under N₂ overnight then cool to

284

room temperature.

Weigh out 10.59 gms dibutyl mesylate (0.040 moles). Dissolve with anhydrous DMSO and add to reaction solution. Purge well with N₂, heat overnight at 80°C.


5

Cool to room temperature. Prepare 500 mls of 5% acetic acid in a 2 liter beaker. While stirring, slowly add reaction mixture. Stir 30 min. Extract with ether 3 times. Combine organic layers and extract with water and sat'd NaCl. Dry organic layer with MgSO₄, filter and rotovap to dryness. Dry oil on vacuum line. Obtain pure product by column chromatography using 95% hexane and 5% ethyl acetate as the mobile phase. Dry wt=7.8 gms. Obtain NMR and mass spec (m/z=444).

10

15

Step 4



$C_{21}H_{29}NO_6$ fw=475.61

20

Dissolve 9.33 gms (0.021 moles) of product 3 with 120 mls anhydrous methylene chloride. Place in a 250 ml round bottom flask with magnetic stir bar. Fit flask with N₂ inlet and stopper. Chill solution with ice bath under N₂ purge. Slowly add 11.54 gms 3-chloroperbenzoic acid (0.0435 moles, Fluka 25800, -65%). After addition is complete warm to room temperature and monitor reaction by TLC. Reaction goes

25

30


285

quickly to the sulphoxide intermediate but takes 8 hrs to convert to the sulphone. Chill solution over night in freezer. Filter solid from reaction, extract filtrate with 10% K₂CO₃. Extract aqueous layer twice with methylene chloride. Combine organic layers and dry with MgSO₄. Filter and rotovap to dryness. Obtain pure product by crystallizing from ethanol or isolating by column chromatography. Obtain NMR and mass spec (m/z=476).

5

10

Step 5



$C_{27}H_{37}NO_6$ fw=473.68

15

Reaction is done in a 300 ml stainless steel Parr stirred mini reactor. Place 9.68 gms (0.0204 moles) of product 4 in reactor base. Add 160 mls ethanol. For safety reasons next two compounds are added in a N₂ atmosphere glove bag. In glove bag, add 15.3 mls formaldehyde (0.204 moles, Aldrich 25,254-9, about 37 wt% in water) and 1.45 gms 10% Pd/Carbon (Aldrich 20,569-9). Seal reactor before removing from glove bag. Purge reactor three times with H₂. Heat to 55°C under H₂. Run reaction at 200 psig H₂, 55°C, and a stir rate of 250 rpm. Run overnight under these conditions. Cool reactor and vent H₂. Purge with N₂. Check progress of run by TLC. Reaction is a mixture of desired product and intermediate. Filter reaction

20


25

286

mixture over a bed of celite washing well with ether. Rotovap and redissolve with ether. Extract with water. Dry organic layer with MgSO_4 , filter and rotovap to dryness. Dry on vacuum line.

Charge reactor again with same amounts, seal reactor and run overnight under same conditions. After second run all of the material has been converted to the desired product. Cool and vent H_2 pressure. Purge with N_2 . Filter over a bed of celite, washing well with ether. Rotovap to dryness. Dissolve with ether and extract with water. Dry organic layer with MgSO_4 , filter and rotovap to dryness. Dry on vacuum line. Obtain NMR and mass spec ($m/z=474$).

Step 6



20

Dissolve 8.97 gms (0.0189 mole) of product 5 with 135 ml anhydrous THF. Place in a 250 ml round bottom flask with magnetic stir bar. Fit flask with N_2 inlet and stopper. Chill solution with ice/salt bath under N_2 purge. Slowly add 2.55 gms potassium t-butoxide (0.227 mole Aldrich 15,667-1). After addition is complete, continue to stir at -10°C monitoring by TLC. Once reaction is complete, quench by adding 135 ml 10% HCl stirring 10 min. Extract three times with


25

287

ether. Dry organic layer with MgSO_4 , filter and rotovap to dryness. Crystallize from ether. Obtain NMR and mass spec ($m/z=474$).

5

Step 7



10

Dissolve 4.67 gms (0.01 moles) of product 6 with 100 ml anhydrous chloroform. Place in a 250 ml round bottom flask with magnetic stir bar. Fit flask with N_2 inlet adapter and suba seal. Chill solution with dry ice /acetone bath under a N_2 purge. Slowly add, via syringe, 2.84 ml boron tribromide (0.03 moles Aldrich 20,220-7). Stir at cold temperature for 15 min after addition then allow to warm to room temperature. Monitor reaction progress by TLC. Reaction is usually complete in 3 hrs.

15


20

Chill solution with ice bath. Quench with 100 ml 10% K_2CO_3 while stirring rapidly. Stir 10 min, then transfer to sep funnel and allow separation. Remove aqueous layer. Extract organic layer once with 10% HCl, once H_2O , and once with saturated NaCl solution. Dry organic layer with MgSO_4 , filter and rotovap to dryness. Crystallize product from ether. Obtain NMR and mass spec ($m/z=460$).

25

288

Step 8


 $C_{14}H_{24}NO_4SI$ fw=701.71

5 Weigh 0.38 gms NaH (9.57 mmoles Aldrich 19,923-0 60% disp. in mineral oil) in a 250 ml round bottom flask with magnetic stir bar. Fit flask with N_2 inlet and stopper. Chill NaH with ice bath and begin N_2 purge.


10 Dissolve 4.0 gms (8.7 mmoles) of product 7 with 60 mls anhydrous DMF. Add to the cold NaH. Stir at cold temperature for 30 min. Add 1.33 gms K_2CO_3 (9.57 mmoles Fisher P-208).

15 Dissolve 16.1 gms 1,2-bis-(2-iodoethoxy)ethane (43.5 mmoles Aldrich 33,343-3) with 60 mls anhydrous DMF. Add to cold reaction mixture. Warm to room temperature then heat to 40°C overnight under N_2 .

20 Cleanup by diluting with ether and extracting sequentially with 5% NaOH, H_2O , and saturated NaCl. Dry organic layer with $MgSO_4$, filter and dry. Obtain pure product by column chromatography using 75% hexane 25% ethyl acetate as the mobile phase. Obtain NMR and mass spec ($m/z=702$).

Step 9

289



 $C_{14}H_{24}NO_4SI$ fw=802.90

5 Dissolve 1.0 gms (1.43 mmoles) of product 8 with 10 mls anhydrous acetonitrile. Place in a 3 ounce Fischer-Porter pressure reaction vessel with magnetic stir bar. Add 2.9 gms triethyl amine (28.6 mmoles Aldrich 23,962-3) dissolved in 10 mls anhydrous acetonitrile. Purge well with N_2 then close system. Heat at 45°C. Monitor reaction by TLC. Reaction is usually complete in 48 hrs.


15 Perform cleanup by removing acetonitrile under vacuum. Redissolve with anhydrous chloroform and precipitate quaternary ammonium salt with ether. Repeat several times. Dry to obtain crystalline product. Obtain NMR and mass spec ($m/z=675$).

20

290

Example 1399**Step 1. Preparation of 1**


To a solution of 144 g of KOH (2560 mmol) in 1.1 L of DMSO was added 120 g of 2-bromobenzyl alcohol (641 mmol) slowly via addition funnel. Then was added 182 g of methyl iodide (80 mL, 1282 mmol) via addition funnel. Stirred at ambient temperature for fifteen minutes. Poured reaction contents into 1.0 L of water and extracted three times with ethyl acetate. The organic layer was dried over MgSO_4 and concentrated *in vacuo*. Purified by silica-gel chromatography through a 200 mL plug using hexanes (100%) as eluant yielded 103.2 g (80%) of 1 as a clear colorless liquid. ^1H NMR (CDCl_3) δ 3.39 (s, 3H), 4.42 (s, 2H), 7.18-7.27 (m, 2H), 7.12 (d, J = 7.45, 1H), 7.50 (s, 1H).

Step 2. Preparation of 2

To a cooled (-78°C) solution of 95 g (472 mmol) of 1 in 1.5 L THF was added 240 mL of 2.5 M n-butyl lithium (576 mmol). The mixture was stirred for one hour, and then to it was added 180 g of zinc iodide (566 mmol) dissolved in 500 mL THF. The mixture was stirred thirty minutes, allowed to warm to 5°C , cooled to -10°C and to it was added 6 g of $\text{Pd(PPh}_3)_4$ (5.2 mmol) and 125 g 2,5-difluorobenzoyl chloride (708 mmol). The mixture was stirred at ambient temperature for 18

291


hours and then cooled to 10°C , quenched with water, partitioned between ethyl acetate and water, and washed organic layer with 1N HCl and with 1N NaOH. The organic layer was dried over MgSO_4 and concentrated *in vacuo*. Purification by silica gel chromatography (Waters Prep-500) using 5% ethyl acetate/hexanes as eluant gave 53.6 g (43 %) of 2 as an orange oil. ^1H NMR (CDCl_3) δ 3.40 (s, 3H), 4.51 (s, 2H), 7.12-7.26 (m, 3H), 7.47 (t, J = 7.50, 1H), 7.57 (d, J = 7.45, 1H), 7.73 (d, J = 7.45, 1H), 7.80 (s, 1H).

Step 3. Preparation of 3

A solution of 53 g (202.3 mmol) of 2 and 11.2 g Li_2S (242.8 mmol) in 250 mL DMF was heated to 100°C for 18 hours. The reaction was cooled (0°C) and 60.7 g of X' (the cyclic sulfate compound of example 1397) (242.8 mmol) in 50 mL DMF was added. Stirred at ambient temperature for 18 hours then condensed *in vacuo*. Added 1 L water to organic residue and extracted twice with diethyl ether. Aqueous layer acidified (pH 1) and refluxed 2 days. Cooled to ambient temperature and extracted with methylene chloride, dried organic layer over MgSO_4 and condensed *in vacuo*. Purification by silica gel chromatography (Waters Prep-500) using 10% ethyl acetate / hexanes as eluant gave 42.9 g (48 %) of 3 as a yellow oil. ^1H NMR (CDCl_3) δ 0.86 (t, J = 7.25 Hz, 6H), 1.10 - 1.26 (m, 12H), 2.83 (s, 2H), 3.32 (s, 2H), 3.40 (s, 3H), 4.48 (s, 3H), 7.02 (dd, J = 8.26 Hz and 2.82 Hz, 1H), 7.16 (dt, J = 8.19 Hz and 2.82 Hz, 1H), 7.45 (t, J = 7.65 Hz, 1H), 7.56-7.61 (m, 2H), 7.69 (d, J = 7.85 Hz, 1H), 7.74 (s, 1H).

292

Step 4. Preparation of 4




- 5 To a cooled (-40 °C) solution of 42.9 g (96.2 mmol) of 3 in 200 mL of methylene chloride was added 21.6 g trifluoromethane sulfonic acid (12.8 mL, 144 mmol) followed by the addition of 22.4 g triethyl silane (30.7 mL, 192.4 mmol). Stirred at -20 °C for two hours, quenched with water and warmed to ambient temperature.

10

Partitioned between methylene chloride and water, dried the organic layer over MgSO_4 and condensed in vacuo. Purification by silica gel chromatography (Waters Prep-500) using 10% ethyl acetate/hexanes as elutant gave 24.2 g (60%) of 4 as a oil. $^1\text{H NMR}$ (CDCl_3) δ 0.89 (t, J = 7.05 Hz, 6H), 1.17 - 1.40 (m, 12H), 1.46 (t, J = 5.84 Hz, 1H), 2.81 (s, 2H), 3.38 (s, 3H), 3.43 (d, J = 5.23 Hz, 2H), 4.16 (s, 2H), 4.42 (s, 2H), 6.80 (d, J = 9.67 Hz, 1H), 6.90 (t, J = 8.46 Hz, 1H), 7.09 (d, J = 7.45 Hz, 1H), 7.15 - 7.21 (m, 2H), 7.25 - 7.32 (m, 2H), 7.42 (m, 1H).

15

Step 5. Preparation of 5




- 25 To a cooled (15-18 °C) solution of 24.2 g (55.8 mmol) of 4 in 100 mL DMSO was added 31.2 g sulfur trioxide pyridine complex (195 mmol). Stirred at ambient temperature for thirty minutes. Poured into cold water

293

and extracted three times with ethyl acetate. Washed organics with 5% HCl (300 mL) and then with brine (300 mL), dried organics over MgSO_4 and condensed in vacuo to give 23.1 g (96 %) of 5 as a light brown oil. $^1\text{H NMR}$ (CDCl_3) δ 0.87 (t, J = 7.05 Hz, 6H), 1.01 - 1.32 (m, 8H), 1.53 - 1.65 (m, 4H), 2.98 (s, 2H), 3.38 (s, 3H), 4.15 (s, 2H), 4.43 (s, 2H), 6.81 (dd, J = 9.66 Hz and 2.82 Hz, 1H), 6.91 (t, J = 8.62 Hz, 1H), 7.07 (d, J = 7.46 Hz, 1H), 7.14 (s, 1H), 7.19 (d, J = 7.65 Hz, 1H), 7.26 - 7.32 (m, 1H), 7.42 (dd, J = 8.66 Hz and 5.64 Hz, 1H), 9.40 (s, 1H).

10

Step 6. Preparation of 6




15

To a cooled (0 °C) solution of 23.1 g (53.6 mmol) of 5 in 200 mL methylene chloride was added 28.6 g meta chloroperoxy-benzoic acid (112.6 mmol). Stirred at ambient temperature for 24 hours. Quenched with 100 mL 10% Na_2SO_3 , partitioned between water and methylene chloride. Dried organic layer over MgSO_4 and condensed in vacuo to give 24.5 g (98%) of 6 as a light yellow oil. $^1\text{H NMR}$ (CDCl_3) δ 0.86 - 1.29 (m, 14H), 1.58 - 1.63 (m, 2H), 1.82 - 1.91 (m, 2H), 3.13 (s, 2H), 3.39 (s, 3H), 4.44 (s, 2H), 4.50 (s, 2H), 6.93 (d, J = 9.07 Hz, 1H), 7.10 - 7.33 (m, 5H), 8.05 (s, 1H), 9.38 (s, 1H).

20


Step 7. Preparation of 7

294



To a solution of 24.5 g (52.9 mmol) of **6** in 20 mL of THF contained in a stainless steel reaction vessel was added 100 mL of a 2.0 M solution of dimethyl amine and 20 mL of neat dimethyl amine. The vessel was sealed and heated to 110 °C for 16 hours. The reaction vessel was cooled to ambient temperature and the contents concentrated *in vacuo*. Purification by silica gel chromatography (Waters Prep-500) using 15 % ethyl acetate/hexanes gave 21.8 g (84 %) of **7** as a clear colorless oil. ¹H NMR (CDCl₃) δ 0.85 (t, J = 7.25 Hz, 6H), 0.93 - 1.29 (m, 8H), 1.49 - 1.59 (m, 2H), 1.70 - 1.80 (m, 2H), 2.98 (s, 8H), 3.37 (s, 3H), 4.41 (s, 2H), 4.44 (s, 2H), 6.42 (s, 1H), 6.58 (dd, J = 9.0 Hz and 2.61 Hz, 1H), 7.13 (d, J = 7.45 Hz, 1H), 7.21 (s, 1H), 7.28 (t, J = 7.85 Hz, 1H), 7.82 (d, J = 9.06 Hz, 1H), 9.36 (s, 1H).

Step 8. Preparation of **8**




A solution of 21.8 g (44.8 mmol) of **7** in 600 mL of THF was cooled to 0 °C. 58.2 mL of a 1 M solution of potassium t-butoxide was added slowly, maintaining the temperature at <5 °C. Stirred for 30 minutes, then

295

quenched with 50 mL of saturated ammonium chloride. The organic layer was partitioned between ethyl acetate and water, dried over MgSO₄ and concentrated *in vacuo*. Purification by recrystallization from ~10% ethyl acetate/hexanes gave 15.1 g of **8** as a white solid. The mother liquor was purified by silica gel chromatography (Waters Prep-500) using 30% ethyl acetate/hexanes as the eluant to give 3.0 g of **8** as a white solid. MS (FAB/LS) m/e 494.6. HRMS (EI) calculated for M+H 487.2756. Found 487.2746.


Step 9. Preparation of **9**



A solution of 2.0 g (4.1 mmol) of **8** in 20 mL of methylene chloride was cooled to -60 °C. 4.1 mL of a 1M solution of boron tribromide was added. Stirred at ambient temperature for thirty minutes. Cooled reaction to -10 °C and quenched with 50 mL of water. The organic layer was partitioned between methylene chloride and water, dried over MgSO₄ and concentrated *in vacuo*. Purification by recrystallization from 50% ethyl acetate/methylene chloride gave 1.95 g (89%) of **9** as a white solid. MS (FAB/LS) m/e 537. HRMS (FAB) calculated for M 536.1834. Found 536.1822.

Step 10. Preparation of **10**

296



A solution of 1.09 g (2.0 mmol) of 9 and 4.9 g (62 mmol) of pyridine in 30 mL of acetonitrile was stirred at ambient temperature for 18 hours. The reaction was concentrated in vacuo. Purification by recrystallization from methanol/ diethyl ether gave 1.19 g (96%) of 10 as an off white solid. MS (FAB) m/e 535.5.


5

10

297

Example 1398

Step 1. Preparation of 2



To a solution of 6.0 g of dibutyl 4-fluorobenzene dialdehyde of Example 1395 (14.3 mmol) in 72 mL of toluene and 54 mL of ethanol was added 4.7 g 3-nitrobenzenesboronic acid (28.6 mmol), 0.8 g of tetrakis (triphenylphosphine) palladium(0) (0.7 mmol) and 45 mL of a 2 M solution of sodium carbonate in water. This heterogeneous mixture was refluxed for three hours, then cooled to ambient temperature and partitioned between ethyl acetate and water. The organic layer was dried over MgSO₄ and concentrated in vacuo.

10


15

Purification by silica gel chromatography (Waters Prep-2000) using ethyl acetate/hexanes (25/75) gave 4.8 g (73%) of the title compound as a yellow solid. ¹H NMR (CDCl₃) δ 0.88 (t, J = 7.45 Hz, 6H), 0.99-1.38 (m, 8H), 1.62-1.75 (m, 2H), 1.85-2.00 (m, 2H), 3.20 (s, 2H), 4.59 (s, 2H), 6.93 (dd, J = 10.5 and 2.4 Hz, 1H), 7.15 (dt, J = 8.4 and 2.85 Hz, 1H), 7.46-7.59 (m, 2H), 8.05-8.16 (m, 3H), 9.40 (s, 1H).

20

298

Step 3. Preparation of 3




5 A solution of 4.8 g (10.4 mmol) of 2 in 500 mL THF was cooled to 0 °C in an ice bath. 20 mL of a 1 M solution of potassium t-butoxide was added slowly, maintaining the temperature at <5 °C. Stirring was continued for 30 minutes, then the reaction was quenched with 100 mL of saturated ammonium chloride. The mixture was partitioned between ethyl acetate and water; the organic layer was washed with brine, then dried (MgSO₄) and concentrated in vacuo. Purification by silica gel chromatography through a 100 mL plug using CH₂Cl₂ as eluent yielded 4.3 g (90%) of 3 as a pale yellow foam.

15 ¹H NMR (CDCl₃) δ 0.93 (t, J = 7.25 Hz, 6H), 1.00-1.55 (m, 8H), 1.59-1.74 (m, 3H), 2.15-2.95 (m, 1H), 3.16 (q_{AB}, J_{AB} = 15.0 Hz, Δν = 33.2 Hz, 2H), 4.17 (d, J = 6.0 Hz, 1H), 5.67 (s, 1H), 6.34 (dd, J = 9.6 and 3.0 Hz, 1H), 7.08 (dt, J = 8.5 and 2.9 Hz, 1H), 7.64 (t, J = 8.1 Hz, 1H), 7.81 (d, J = 8.7 Hz, 1H), 8.13 (dd, J = 9.9 and 3.6 Hz, 1H), 8.23-8.30 (m, 1H), 8.44 (s, 1H). MS (FABH⁺) m/e (relative intensity) 464.5 (100), 446.6 (65). HRMS calculated for M+H 464.1907. Found 464.1905.

299

Step 4. Preparation of 4




5 To a cooled (0 °C) solution of 4.3 g (9.3 mmol) of 3 in 30 mL THF contained in a stainless steel reaction vessel was added 8.2 g dimethyl amine (182 mmol). The vessel was sealed and heated to 110 °C for 16 hours. The reaction vessel was cooled to ambient temperature and the contents concentrated in vacuo. Purification by silica gel chromatography (Waters Prep-2000) using an ethyl acetate/hexanes gradient (10-40% ethyl acetate) gave 4.0 g (88%) of 4 as a yellow solid.

15 NMR (CDCl₃) δ 0.80-0.95 (m, 6H), 0.96-1.53 (m, 8H), 1.60-1.69 (m, 3H), 2.11-2.28 (m, 1H), 2.79 (s, 6H), 3.09 (q_{AB}, J_{AB} = 15.0 Hz, Δν = 45.6 Hz, 2H), 4.90 (d, J = 9.0 Hz, 1H), 5.65 (s, 1H), 5.75 (d, J = 2.1 Hz, 1H), 6.52 (dd, J = 9.6 and 2.7 Hz, 1H), 7.59 (t, J = 8.4 Hz, 1H), 7.85 (d, J = 7.80 Hz, 1H), 7.89 (d, J = 9.0 Hz, 1H), 8.20 (dd, J = 8.4 and 1.2 Hz, 1H), 8.43 (s, 1H). MS (FABH⁺) m/e (relative intensity) 489.6 (100), 471.5 (25). HRMS calculated for M+H 489.2423. Found 489.2456.

25

300

Step 5. Preparation of 5




To a suspension of 1.0 g (2.1 mmol) of **4** in 100 ml ethanol in a stainless steel Parr reactor was added 1 g 10% palladium on carbon. The reaction vessel was sealed, purged twice with H_2 , then charged with H_2 (100 psi) and heated to 45 °C for six hours. The reaction vessel was cooled to ambient temperature and the contents filtered in vacuo to remove the catalyst. The filtrate was concentrated in vacuo to give 0.9 g (96%) of **5**. 1H NMR ($CDCl_3$) δ 0.80-0.98 (m, 6H), 1.00-1.52 (m, 10H), 1.52-1.69 (m, 1H), 2.15-2.29 (m, 1H), 2.83 (s, 6H), 3.07 (q_{AB} , J_{AB} = 15.1 Hz, DV = 44.2 Hz, 2H), 3.70 (s, 2H), 4.14 (s, 1H), 5.43 (s, 1H), 6.09 (d, J = 2.4 Hz, 1H), 6.52 (dd, J = 12.2 and 2.6 Hz, 1H), 6.65 (dd, J = 7.8 and 1.8 Hz, 1H), 6.83 (s, 1H), 6.93 (d, J = 7.50 Hz, 1H), 7.19 (t, J = 7.6 Hz, 1H), 7.89 (d, J = 8.9 Hz, 1H). MS ($FABH^+$) m/e (relative intensity) 459.7 (100). HRMS calculated for $M+H$ 459.2681. Found 459.2670.

361

Step 6. Preparation of 6

To a solution of 914 mg (2.0 mmol) of **5** in 50 ml THF was added 800 mg (4.0 mmol) 5-bromovaleryl chloride. Next was added 4 g (39.6 mmol) TEA. The reaction was stirred 10 minutes, then partitioned between ethyl acetate and brine. The organic layer was dried ($MgSO_4$) and concentrated in vacuo. Purification by silica gel chromatography through a 70 ml MPLC column using a gradient of ethyl acetate (20-50%) in hexane as eluent yielded 0.9 g (73%) of **6** as a pale yellow oil. 1H NMR ($CDCl_3$) δ 0.84-0.95 (m, 6H), 1.02-1.53 (m, 10H), 1.53-1.68 (m, 1H), 1.80-2.00 (m, 4H), 2.12-2.26 (m, 4H), 2.38 (t, J = 6.9 Hz, 2H), 2.80 (s, 6H), 3.07 (q_{AB} , J_{AB} = 15.6 Hz, DV = 40.4 Hz, 2H), 3.43 (t, J = 6.9 Hz, 2H), 4.10 (s, 1H), 5.51 (s, 1H), 5.95 (d, J = 2.4 Hz, 1H), 6.51 (dd, J = 9.3 and 2.7 Hz, 1H), 7.28 (s, 1H), 7.32-7.41 (m, 2H), 7.78 (d, J = 8.1 Hz, 1H), 7.90 (d, J = 9.0 Hz, 1H).

Step 7. Preparation of 7




To a solution of 0.9 g (1.45 mmol) of **6** in 25 ml acetonitrile add 18 g (178 mmol) TEA. Heat at 55 °C for 16 hours. The reaction mixture was cooled to ambient temperature and concentrated in vacuo.

Purification by reverse-phase silica gel chromatography (Waters Delta Prep 3000) using an acetonitrile /water


362

gradient containing 0.05% TFA (20-65% acetonitrile) gave 0.8 g (73%) of 7 as a white foam. ¹H NMR (CDCl₃) δ 0.80-0.96 (m, 6H), 0.99-1.54 (m, 19H), 1.59-1.84 (m, 3H), 2.09-2.24 (m, 1H), 2.45-2.58 (m, 2H), 2.81 (s, 6H), 3.09 (q_{AB}, J_{AB} = 15.6 Hz, DV = 18.5 Hz, 2H), 3.13-3.31 (m, 8H), 4.16 (s, 1H), 5.44 (s, 1H), 6.08 (d, J = 1.8 Hz, 1H), 6.57 (dd, J = 9.3 and 2.7 Hz, 1H), 7.24 (t, J = 7.5 Hz, 1H), 7.34 (t, J = 8.4 Hz, 1H), 7.56 (d, J = 8.4 Hz, 1H), 7.74 (s, 1H), 7.88 (d, J = 9.0 Hz, 1H), 9.22 (s, 1H). HRMS calcd 642.4304; observed 642.4343.

Example 1400**Step 1**C₁₄H₁₃O₂F fw=232.25

A 12-liter, 4-neck round-bottom flask was equipped with reflux condenser, N₂ gas adaptor, mechanical stirrer, and an addition funnel. The system was purged with N₂. A slurry of sodium hydride (126.0g/4.988mol) in toluene (2.5 L) was added, and the mixture was cooled to 6 C. A solution of 4-fluorophenol (560.5g/5.000mol) in toluene (2.5 L) was added via addition funnel over a period of 2.5 h. The reaction mixture was heated to reflux (100 C) for 1h. A solution of 3-methoxybenzyl chloride (783.0g/5.000mol) in toluene (750 mL) was added via addition funnel while maintaining reflux. After 15 h. refluxing, the mixture was cooled to room temperature and poured into H₂O (2.5 L). After 20 min. stirring, the layers were separated, and the organic layer was extracted with a solution of potassium

hydroxide (720g) in MeOH (2.5 L). The MeOH layer was added to 20% aqueous potassium hydroxide, and the mixture was stirred for 30 min. The mixture was then washed 5 times with toluene. The toluene washes were extracted with 20% aq. KOH. All 20% aq. KOH solutions were combined and acidified with concentrated HCl. The acidic solution was extracted three times with ethyl ether, dried (MgSO₄), filtered and concentrated in vacuo. The crude product was purified by Kugelrohr distillation to give a clear, colorless oil (449.0g/39% yield). b.p.: 120-130 C/50mmHg. ¹H NMR and MS (M + H)⁺ = 233 confirmed desired structure.


Step 2C₁₇H₁₈NO₂FS fw=319.39

A 12-liter, 3-neck round-bottom flask was fitted with mechanical stirrer and N₂ gas adaptor. The system was purged with N₂. 4-Fluoro-2-(3-methoxybenzyl)-phenol (455.5g/1.961mol) and dimethylformamide were added. The solution was cooled to 6 C, and sodium hydride (55.5g/2.197mol) was added slowly. After warming to room temperature, dimethylthiocarbamoyl chloride (242.4g/1.961mol) was added. After 15 h, the reaction mixture was poured into H₂O (4.0 L), and extracted two times with ethyl ether. The combined organic layers were washed with H₂O and saturated aqueous NaCl, dried (MgSO₄), filtered, and concentrated in vacuo to give

the product (605.3g, 97% yield). ^1H NMR and MS ($(\text{M}+\text{H})^+ = 320$) confirm desired structure.

5

Step 3




$\text{C}_{14}\text{H}_{13}\text{OFS}$ fw=248.32

A 12-liter, round-bottom flask was equipped with N_2 gas adaptor, mechanical stirrer, and reflux condenser. The system was purged with N_2 . 4-Fluoro-2-(3-methoxybenzyl)-phenyldimethylthiocarbamate (605.3g/1.895mol) and phenyl ether (2.0kg) were added, and the solution was heated to reflux for 2 h. The mixture was stirred for 64 h. at room temperature and then heated to reflux for 2 h. After cooling to room temperature, MeOH (2.0 L) and THF (2.0 L) were added, and the solution was stirred for 15 h. Potassium hydroxide (425.9g/7.590mol) was added, and the mixture was heated to reflux for 4 h. After cooling to room temperature, the mixture was concentrated by rotavap, dissolved in ethyl ether (1.0 L), and extracted with H_2O . The aqueous extracts were combined, acidified with concentrated HCl, and extracted with ethyl ether. The ether extracts were dried (MgSO_4), filtered, and concentrated in vacuo to give an amber oil (463.0g, 98% yield). ^1H NMR confirmed desired structure.

Step 4

305




$\text{C}_{25}\text{H}_{35}\text{O}_2\text{FS}$ fw=418.61

A 5-liter, 3-neck, round-bottom flask was equipped with N_2 gas adaptor and mechanical stirrer. The system was purged with N_2 . 4-Fluoro-2-(3-methoxybenzyl)-thiophenol (100.0g/403.2mmol) and 2-methoxyethyl ether (1.0 L) were added and the solution was cooled to 0 C. Sodium hydride (9.68g/383.2mmol) was added slowly, and the mixture was allowed to warm to room temperature. 2,2-Dibutylpropylene sulfate (110.89g/443.6mmol) was added, and the mixture was stirred for 64 h. The reaction mixture was concentrated by rotavap and dissolved in H_2O . The aqueous solution was washed with ethyl ether, and concentrated H_2SO_4 was added. The aqueous solution was heated to reflux for 30 min, cooled to room temperature, and extracted with ethyl ether. The ether solution was dried (MgSO_4), filtered, and conc'd in vacuo to give an amber oil (143.94g/85% yield). ^1H NMR and MS ($(\text{M}+\text{H})^+ = 419$) confirm the desired structure.

Step 5


306



$C_{25}H_{33}O_2FS$ fw=416.59

A 2-liter, 4-neck, round-bottom flask was equipped with N_2 gas adaptor, and mechanical stirrer. The system was purged with N_2 . The corresponding alcohol (143.94g/343.8mmol) and CH_2Cl_2 (1.0 L) were added and cooled to 0 C. Pyridinium chlorochromate (140.53g/651.6mmol) was added. After 6 h., CH_2Cl_2 was added. After 20 min, the mixture was filtered through silica gel, washing with CH_2Cl_2 . The filtrate was concentrated in vacuo to give a dark yellow-red oil (110.6g, 77% yield). 1H NMR and MS ($M + H^+ = 417$) confirm the desired structure.

Step 6



$C_{25}H_{33}O_4FS$ fw=448.59


A 2-liter, 4-neck, round-bottom flask was equipped with N_2 gas adaptor and mechanical stirrer. The system was

307

purged with N_2 . The corresponding sulfide (110.6g/265.5mmol) and CH_2Cl_2 (1.0 L) were added. The solution was cooled to 0 C, and 3-chloroperoxybenzoic acid (158.21g/531.7mmol) was added portionwise. After 30 min, the reaction mixture was allowed to warm to room temperature. After 3.5 h, the reaction mixture was cooled to 0 C and filtered through a fine fritted funnel. The filtrate was washed with 10% aqueous K_2CO_3 . An emulsion formed which was extracted with ethyl ether. The organic layers were combined, dried ($MgSO_4$), filtered, and concentrated in vacuo to give the product (93.2g, 78% yield). 1H NMR confirmed the desired structure.

308

Step 7




$C_{25}H_{33}O_4FS$ fw=448.59

- 5 A 2-liter, 4-neck, round-bottom flask was equipped with N_2 gas adaptor, mechanical stirrer, and a powder addition funnel. The system was purged with N_2 . The corresponding aldehyde (93.2g/208mmol) and THF (1.0 L) were added, and the mixture was cooled to 0 C.
- 10 Potassium tert-butoxide (23.35g/208.1mmol) was added via addition funnel. After 1h, 10% aq/HCl (1.0 L) was added. After 1 h, the mixture was extracted three times with ethyl ether, dried ($MgSO_4$), filtered, and concentrated in vacuo. The crude product was purified by recryst. from 80/20 hexane/ethyl acetate to give a white solid (32.18 g). The mother liquor was concentrated in vacuo and recrystelized from 95/5 toluene/ethyl acetate to give a white solid (33.60g/ combined yield: 71%). 1H NMR confirmed the desired product.
- 15
- 20

309

Step 8




$C_{27}H_{39}O_4NS$ fw=473.67

- 5 A Fisher porter bottle was fitted with N_2 line and magnetic stirrer. The system was purged with N_2 . The corresponding fluoro-compound (28.1g/62.6mmol) was added, and the vessel was sealed and cooled to -78 C.
- 10 Dimethylamine (17.1g/379mmol) was condensed via a CO_2 /acetone bath and added to the reaction vessel. The mixture was allowed to warm to room temperature and was heated to 60 C. After 20 h, the reaction mixture was allowed to cool and was dissolved in ethyl ether. The ether solution was washed with H_2O , saturated aqueous NaCl, dried ($MgSO_4$), filtered, and concentrated in vacuo to give a white solid (28.5g/96% yield). 1H NMR confirmed the desired structure.
- 15

310

Step 9




$C_{26}H_{37}O_4NS$ fw=459.64

A 250-ml, 3-neck, round-bottom flask was equipped with N_2 gas adaptor and magnetic stirrer. The system was purged with H_2 . The corresponding methoxy-compound (6.62g/14.0mmol) and $CHCl_3$ (150 mL) were added. The reaction mixture was cooled to $-78^\circ C$, and boron tribromide (10.50g/41.9mmol) was added. The mixture was allowed to warm to room temperature. After 4 h, the reaction mixture was cooled to $0^\circ C$ and was quenched with 10% K_2CO_3 (100 mL). After 10 min, the layers were separated, and the aqueous layer was extracted two times with ethyl ether. The $CHCl_3$ and ether extracts were combined, washed with saturated aqueous $NaCl$, dried ($MgSO_4$), filtered, and concentrated in vacuo to give the product (6.27g/98% yield). 1H NMR confirmed the desired structure.

311

Step 10




In a 250 ml single neck round bottom flask with stir bar place 2-diethylaminoethyl chloride hydrochloride (fw 172.10g/mole) Aldrich D8, 720-1 (2.4 mmol, 4.12g), 34 ml dry ether and 34 ml of 1N KOH(aqueous). Stir 15 minutes and then separate by ether extraction and dry over anhydrous potassium carbonate.

In a separate 2-necked 250 ml round bottom flask with stir bar add sodium hydride (60% dispersion in mineral oil, 100 mg, 2.6 mmol) and 34 ml of DMF. Cool to ice temperature. Next add phenol product (previous step) 1.1 g (2.4 mmol) in 5 ml DMF and the ether solution prepared above. Heat to $40^\circ C$ for 3 days. The product which contained no starting material by TLC was diluted with ether and extracted with 1 portion of 5% $NaOH$, followed by water and then brine. The ether layer was dried over magnesium sulfate and isolated by removing ether by rotary evaporation (1.3 gms). The product may be further purified by chromatography (SiO_2 98% ethyl acetate/1% NH_4OH at 5ml/min.). Isolated yield: 0.78 g (mass spec, and 1H NMR)

312

Step 11




5 The product from step 10 (0.57gms, 1.02 millimole fw 558.83 g/mole) and 1.6 gms iodoethane (10.02 mmol) was placed in 5 ml acetonitrile in a fischer-porter bottle and heated to 45 C for 3 days. The solution was evaporated to dryness and redissolved in 5 mls of chloroform. Next ether was added to the chloroform solution and the resulting mixture was chilled. The desired product is isolated as a precipitate 0.7272 gms. Mass spec M-I = 587.9 , H NMR).

15

Example 1401

Step 1



$C_{14}H_{13}O_2F$ fw=232.25

20

A 12-liter, 4-neck round-bottom flask was equipped with reflux condenser, N_2 gas adaptor, mechanical stirrer, and an addition funnel. The system was purged with N_2 .

5 A slurry of sodium hydride (126.0g/4.988mol) in toluene (2.5 L) was added, and the mixture was cooled to 6 C. A solution of 4-fluorophenol (560.5g/5.000mol) in toluene (2.5 L) was added via addition funnel over a period of 2.5 h. The reaction mixture was heated to reflux (100 C) for 1h. A solution of 3-methoxybenzyl chloride (783.0g/5.000mol) in toluene (750 mL) was added via addition funnel while maintaining reflux.


10 After 15 h. refluxing, the mixture was cooled to room temperature and poured into H_2O (2.5 L). After 20 min. stirring, the layers were separated, and the organic layer was extracted with a solution of potassium

15 hydroxide (720g) in MeOH (2.5 L). The MeOH layer was added to 20% aqueous potassium hydroxide, and the mixture was stirred for 30 min. The mixture was then washed 5 times with toluene. The toluene washes were extracted with 20% aq. KOH. All 20% aqueous KOH solutions were combined and acidified with concentrated HCl. The acidic solution was extracted three times with ethyl ether, dried over $MgSO_4$, filtered and concentrated in vacuo. The crude product was purified

25 by Kugelrohr distillation to give a clear, colorless oil (449.0g/39% yield). b.p.: 120-130 C/50mmHg.

1H NMR and MS [(M + H)⁺ = 233] confirmed desired structure.

30 Step 2


C₁₇H₁₈NO₂PS fw=319.39

A 12-liter, 3-neck round-bottom flask was fitted with mechanical stirrer and N₂ gas adaptor. The system was purged with N₂. 4-Fluoro-2-(3-methoxybenzyl)-phenol (455.5g/1.961mol) and dimethylformamide were added. The solution was cooled to 6 C, and sodium hydride (55.5g/2.197mol) was added slowly. After warming to room temperature, dimethylthiocarbamoyl chloride (242.4g/1.961mol) was added. After 15 h, the reaction mixture was poured into H₂O (4.0 L), and extracted two times with ethyl ether. The combined organic layers were washed with H₂O and saturated aqueous NaCl, dried over MgSO₄, filtered, and concentrated in vacuo to give the product (605.3g, 97% yield). ¹H NMR and MS [(M+H)⁺ = 320] confirm desired structure.

Step 3

20

315


C₁₄H₁₃OFS fw=248.32

A 12-liter, round-bottom flask was equipped with N₂ gas adaptor, mechanical stirrer, and reflux condenser. The system was purged with N₂. 4-Fluoro-2-(3-methoxybenzyl)-phenyldimethylthiocarbamate (605.3g/1.895mol) and phenyl ether (2.0kg) were added, and the solution was heated to reflux for 2 h. The mixture was stirred for 64 h, at room temperature and then heated to reflux for 2 h. After cooling to room temperature, MeOH (2.0 L) and T₂F (2.0 L) were added, and the solution was stirred for 15 h. Potassium hydroxide (425.9g/7.590mol) was added, and the mixture was heated to reflux for 4 h. After cooling to room temperature, the mixture was concentrated by rotavap, dissolved in ethyl ether (1.0 L), and extracted with H₂O. The aqueous extracts were combined, acidified with conc. HCl, and extracted with ethyl ether. The ether extracts were dried (MgSO₄), filtered, and concentrated in vacuo to give an amber oil (463.0g, 98% yield). ¹H NMR confirmed desired structure.

Step 4

20


316



$C_{25}H_{35}O_2FS$ fw=418.61

- 5 A 5-liter, 3-neck, round-bottom flask was equipped with N_2 gas adaptor and mechanical stirrer. The system was purged with N_2 . 4-Fluoro-2-(3-methoxybenzyl)-thiophenol (100.0g/403.2mmol) and 2-methoxyethyl ether (1.0 L) were added and the solution was cooled to 0 C. Sodium hydride (9.68g/383.2mmol) was added slowly, and the mixture was allowed to warm to room temperature. 2,2-Dibutylpropylene sulfide (110.89g/443.6mmol) was added, and the mixture was stirred for 64 h. The reaction mixture was concentrated by rotavap and dissolved in H_2O . The aqueous solution was washed with ethyl ether, and conc. H_2SO_4 was added. The aqueous solution was heated to reflux for 30 min, cooled to room temperature, and extracted with ethyl ether. The ether solution was dried ($MgSO_4$), filtered, and concentrated in vacuo to give an amber oil (143.94g/85% yield). 1H NMR and MS ($M + H^+$ = 419) confirm the desired structure.
- 10
- 15
- 20

317




$C_{25}H_{33}O_2FS$ fw=416.59

- 5 A 2-liter, 4-neck, round-bottom flask was equipped with N_2 gas adaptor, and mechanical stirrer. The system was purged with N_2 . The corresponding alcohol (143.94 g/343.8 mmol) and CH_2Cl_2 (1.0 L) were added and cooled to 0 C. Pyridinium chlorochromate (140.53g/651.6mmol) was added. After 6 h., CH_2Cl_2 was added. After 20 min, the mixture was filtered through silica gel, washing with CH_2Cl_2 . The filtrate was concentrated in vacuo to give a dark yellow-red oil (110.6g, 77% yield). 1H NMR and MS ($M + H^+$ = 417) confirm the desired structure.
- 10
- 15

318


Step 6



$$C_{25}H_{33}O_4FS \text{ fw}=448.59$$

A 2-liter, 4-neck, round-bottom flask was equipped with N_2 gas adaptor and mechanical stirrer. The system was purged with N_2 . The corresponding sulfide (110.6g/265.5mmol) and CH_2Cl_2 (1.0 L) were added. The solution was cooled to 0 C, and 3-chloroperbenzoic acid (158.21g/531.7mmol) was added portionwise. After 30 min, the reaction mixture was allowed to warm to room temperature. After 3.5 h, the reaction mixture was cooled to 0 C and filtered through a fine fritted funnel. The filtrate was washed with 10% aqueous K_2CO_3 . An emulsion formed which was extracted with ethyl ether. The organic layers were combined, dried ($MgSO_4$), filtered, and concentrated in vacuo to give the product (93.2g, 78% yield). 1H NMR confirmed the desired structure.


Step 7



$$C_{25}H_{33}O_4FS \text{ fw}=448.59$$

A 2-liter, 4-neck, round-bottom flask was equipped with N_2 gas adaptor, mechanical stirrer, and a powder addition funnel. The system was purged with N_2 . The corresponding aldehyde (93.2g/208mmol) and THF (1.0 L) were added, and the mixture was cooled to 0 C. Potassium tert-butoxide (23.35g/208.1mmol) was added via addition funnel. After 1h, 10% aq/ HCl (1.0 L) was added. After 1 h, the mixture was extracted three times with ethyl ether, dried ($MgSO_4$), filtered, and concentrated in vacuo. The crude product was purified by recrystallized from 80/20 hexane/ethyl acetate to give a white solid (32.18g). The mother liquor was concentrated in vacuo and recrystallized from 95/5 toluene/ethyl acetate to give a white solid (33.60g, combined yield: 71%). 1H NMR confirmed the desired product.

Step 8




$C_{27}H_{39}O_4NS$ fw=473.67

- 5 A Fisher porter bottle was fitted with N_2 line and magnetic stirrer. The system was purged with N_2 . The corresponding fluoro-compound (28.1g/62.6mmol) was added, and the vessel was sealed and cooled to $-78^\circ C$. Dimethylamine (17.1g/379mmol) was condensed via a CO_2 /acetone bath and added to the reaction vessel. The mixture was allowed to warm to room temperature and was heated to $60^\circ C$. After 20 h, the reaction mixture was allowed to cool and was dissolved in ethyl ether. The ether solution was washed with H_2O , saturated aqueous NaCl, dried over $MgSO_4$, filtered, and concentrated in *vacuo* to give a white solid (28.5g/96% yield). 1H NMR confirmed the desired structure.
- 10
- 15

321

Step 9




$C_{26}H_{37}O_4NS$ fw=459.64

- 5 A 250-mL, 3-neck, round-bottom flask was equipped with N_2 gas adaptor and magnetic stirrer. The system was purged with N_2 . The corresponding methoxy-compound (6.62g/14.0mmol) and $CHCl_3$ (150 mL) were added. The reaction mixture was cooled to $-78^\circ C$, and boron tribromide (10.50g/41.9mmol) was added. The mixture was allowed to warm to room temperature. After 4 h, the reaction mixture was cooled to $0^\circ C$ and was quenched with 10% K_2CO_3 (100 mL). After 10 min, the layers were separated, and the aqueous layer was extracted two times with ethyl ether. The $CHCl_3$ and ether extracts were combined, washed with saturated aqueous NaCl, dried over $MgSO_4$, filtered, and concentrated in *vacuo* to give the product (6.27g/98% yield). 1H NMR confirmed the desired structure.
- 10
- 15
- 20

322

Step 10




5 In a 250 ml single neck round bottom flask with stir bar place 2-diethylaminoethyl chloride hydrochloride (fw 172.10g/mole) Aldrich D8, 720-1 (2.4 millimoles, 4.12g), 34 ml dry ether and 34 ml of 1N KOH (aqueous). Stir 15 minutes and then separate by ether extraction and dry over anhydrous potassium carbonate.

10 In a separate 2-necked 250 ml round bottom flask with stir bar add sodium hydride (60% dispersion in mineral oil, 100 mg, (2.6 mmol) and 34 ml of DMF. Cool to ice temperature. Next add phenol product (previous step) 1.1 g (2.4 mmol) in 5 ml DMF and the ether solution prepared above. Heat to 40C for 3 days. The product which contained no starting material by TLC was diluted with ether and extracted with 1 portion of 5% NaOH, followed by water and then brine. The ether layer was dried over Magnesium sulfate and isolated by removing ether by rotary evaporation (1.3 gms). The product may be further purified by chromatography (silica 98 ethyl acetate/1% NH₄OH at 5ml/min.).

25 Isolated yield: 0.78 g (mass spec, and H¹ NMR)

323

Step 11



5 The product from step 10 (0.57gms, 1.02 millimole fw 558.83 g/mole) and iodoethane (1.6 gms (10.02 millimoles) was place in 5 ml acetonitrile in a Fischer-Porter bottle and heated to 45 C for 3 days. The solution was evaporated to dryness and redissolved in 5 mls of chloroform. Next ether was added to the chloroform solution and the resulting mixture was chilled. The desired product is isolated as a precipitate 0.7272 gms. Mass spec M-I = 587.9, ¹H NMR).

BIOLOGICAL ASSAYS

15 The utility of the compounds of the present invention is shown by the following assays. These assays are performed in vitro and in animal models essentially using a procedure recognized to show the utility of the present invention.

20 In Vitro Assay of compounds that inhibit IBAT-mediated uptake of [¹⁴C]-Taurocholate (TC) in H14 cells

Baby hamster kidney cells (BHK) transfected with the cDNA of human IBAT (H14 cells) are seeded at 60,000

324

cells/well in 96 well Top-Count tissue culture plates for assays run within in 24 hours of seeding, 30,000 cells/well for assays run within 48 hours, and 10,000 cells/well for assays run within 72 hours.

5 On the day of assay, the cell monolayer is gently washed once with 100 ml assay buffer (Dulbecco's Modified Eagle's medium with 4.5 g/L glucose + 0.2% (w/v) fatty acid free bovine serum albumin- (FAF)BSA).

10 To each well 50 ml of a two-fold concentrate of test compound in assay buffer is added along with 50 ml of 6 mM [¹⁴C]-taurocholate in assay buffer (final concentration of 3 mM [¹⁴C]-taurocholate). The cell culture plates are incubated 2 hours at 37° C prior to gently washing each well twice with 100 ml 4° C

15 Dulbecco's phosphate-buffered saline (PBS) containing 0.2% (w/v) (FAF)BSA. The wells are then gently washed once with 100 ml 4° C PBS without (FAF)BSA. To each 200 ml of liquid scintillation counting fluid is added, the plates are heat sealed and shaken for 30 minutes at room temperature prior to measuring the amount of radioactivity in each well on a Packard Top-Count instrument.

20

In Vitro Assay of compounds that inhibit uptake of [¹⁴C]-Alanine

The alanine uptake assay is performed in an identical fashion to the taurocholate assay, with the exception that labeled alanine is substituted for the labeled taurocholate.

5

In Vivo Assay of compounds that inhibit Rat Ileal uptake of [¹⁴C]-Taurocholate into Bile

10 (See "Metabolism of 3a,7b-dihydroxy-7a-methyl-5b-cholanoic acid and 3a,7b-dihydroxy-7a-methyl-5b-cholanoic acid in hamsters" in Biochimica et Biophysica Acta 833 (1985) 196-202 by Uneyama et al.)

Male wistar rats (200-300 g) are anesthetized with inactin @100 mg/kg. Bile ducts are cannulated with a 10" length of PE10 tubing. The small intestine is exposed and laid out on a gauze pad. A cannulae (1/8" luer lock, tapered female adapter) is inserted at 12 cm from the junction of the small intestine and the cecum.

20 A slit is cut at 4 cm from this same junction (utilizing a 8 cm length of ileum). 20 ml of warm Dulbecco's phosphate buffered saline, pH 6.5 (PBS) is used to flush out the intestine segment. The distal opening is cannulated with a 20 cm length of silicone tubing (0.02" I.D. x 0.037" O.D.). The proximal cannulae is hooked up to a peristaltic pump and the intestine is washed for 20 min with warm PBS at 0.25 ml/min. Temperature of the gut segment is monitored continuously. At the start of the experiment, 2.0 ml of control sample ([¹⁴C]-taurocholate @ 0.05 mi/ml with 5 mM cold taurocholate) is loaded into the gut segment with a 3 ml syringe and bile sample collection is begun. Control sample is infused at a rate of 0.25 ml/min for 21 min. Bile samples fractions are collected every 3 minute for the first 27 minutes of the procedure. After the 21 min of sample infusion, the ileal loop is washed out with 20 ml of warm PBS

25

30

35

(using a 30 ml syringe), and then the loop is washed out for 21 min with warm PBS at 0.25 ml/min. A second perfusion is initiated as described above but this with test compound being administered as well (21 min administration followed by 21 min of wash out) and bile sampled every 3 min for the first 27 min. If necessary, a third perfusion is performed as above that typically contains the control sample.

10 Measurement of Hepatic Cholesterol Concentration (HEPATIC CHOL)

Liver tissue was weighed and homogenized in chloroform:methanol (2:1). After homogenization and centrifugation the supernatant was separated and dried under nitrogen. The residue was dissolved in isopropanol and the cholesterol content was measured enzymatically, using a combination of cholesterol oxidase and peroxidase, as described by Allain, C. A., et al. (1974) Clin. Chem. 20, 470.

20 Measurement of Hepatic HMG CoA-Reductase Activity (HMG COA)

Hepatic microsomes were prepared by homogenizing liver samples in a phosphate/sucrose buffer, followed by centrifugal separation. The final pelleted material was resuspended in buffer and an aliquot was assayed for HMG CoA reductase activity by incubating for 60 minutes at 37° C in the presence of ¹⁴C-HMG-CoA (Dupont-NEN). The reaction was stopped by adding 6N HCl followed by centrifugation. An aliquot of the supernatant was separated, by thin-layer chromatography, and the spot corresponding to the enzyme product was scraped off the plate, extracted and radioactivity was determined by scintillation counting. (Reference: Akerlund, J. and Bjorkhem, I. (1990) J. Lipid Res. 31, 2159).

327

5 Determination of Serum Cholesterol (SER. CHOL, HDL-CHOL, VLDL and VLDL + LDL)

Total serum cholesterol (SER. CHOL) was measured enzymatically using a commercial kit from Wako Fine Chemicals (Richmond, VA); Cholesterol CII, Catalog No. 276-64309. HDL cholesterol (HDL-CHOL) was assayed using this same kit after precipitation of VLDL and LDL with Sigma Chemical Co. HDL Cholesterol reagent, Catalog No. 352-3 (dextran sulfate method). Total serum triglycerides (blanked) (TG1) were assayed enzymatically with Sigma Chemical Co. GPO-Trinder, Catalog No. 337-B. VLDL and LDL (VLDL + LDL) cholesterol concentrations were calculated as the difference between total and HDL cholesterol.

15 Measurement of Hepatic Cholesterol 7- α -Hydroxylase Activity (7 α -OHase)

Hepatic microsomes were prepared by homogenizing liver samples in a phosphate/sucrose buffer, followed by centrifugal separation. The final pelleted material was resuspended in buffer and an aliquot was assayed for cholesterol 7- α -hydroxylase activity by incubating for 5 minutes at 37° C in the presence of NADPH. Following extraction into petroleum ether, the organic solvent was evaporated and the residue was dissolved in acetonitrile/methanol. The enzymatic product was separated by injecting an aliquot of the extract onto a C₁₈ reversed phase HPLC column and quantitating the eluted material using UV detection at 240nm. (Reference: Horcon, J. D., et al. (1994) J. Clin. Invest. 93, 2084).

30 Measurement of Fecal Bile Acid Concentration (FBA)

Total fecal output from individually housed hamsters was collected for 24 or 48 hours, dried under a stream of nitrogen, pulverized and weighed. Approximately 0.1 gram was weighed out and extracted

328

into an organic solvent (butanol/water). Following separation and drying, the residue was dissolved in methanol and the amount of bile acid present was measured enzymatically using the 3 α -hydroxysteroid steroid dehydrogenase reaction with bile acids to reduce NAD. (Reference: Mashige, P., et al. (1981) Clin. Chem. 27, 1352).

10 [³H]-taurocholate Uptake in Rabbit Brush Border Membrane Vesicles (BBMV)

Rabbit ileal brush border membranes were prepared from frozen ileal mucosa by the calcium precipitation method describe by Malathi et al. (Reference: (1979) Biochimica Biophysica Acta, 556, 259). The method for measuring taurocholate was essentially as described by Kramer et al. (Reference: (1992) Biochimica Biophysica Acta, 1111, 93) except the assay volume was 200 μ l instead of 100 μ l. Briefly, at room temperature a 190 μ l solution containing 2 μ M [³H]-taurocholate (0.75 μ Ci), 20 mM tris, 100 mM NaCl, 100 mM mannitol pH 7.4 was incubated for 5 sec with 10 μ l of brush border membrane vesicles (60-120 μ g protein). The incubation was initiated by the addition of the BBMV while vortexing and the reaction was stopped by the addition of 5 ml of ice cold buffer (20 mM Hepes-tris, 150 mM KCl) followed immediately by filtration through a nylon filter (0.2 μ m pore) and an additional 5 ml wash with stop buffer.

30 Acyl-CoA:cholesterol Acyl Transferase (ACAT)

Hamster liver and rat intestinal microsomes were prepared from tissue as described previously (Reference: (1980) J. Biol. Chem. 255, 9098) and used as a source of ACAT enzyme. The assay consisted of a 2.0 ml incubation containing 24 μ M Oleoyl-CoA (0.05 μ Ci) in a 50 mM sodium phosphate, 2 mM DTT pH 7.4 buffer containing 0.25 % BSA and 200 μ g of microsomal

protein. The assay was initiated by the addition of oleoyl-CoA. The reaction went for 5 min at 37 $^{\circ}$ C and was terminated by the addition of 8.0 ml of chloroform/methanol (2:1). To the extraction was added 125 μ g of cholesterol oleate in chloroform methanol to act as a carrier and the organic and aqueous phases of the extraction were separated by centrifugation after thorough vortexing. The chloroform phase was taken to dryness and then spotted on a silica gel 60 TLC plate and developed in hexane/ethyl ether (9:1). The amount of cholesterol ester formed was determined by measuring the amount of radioactivity incorporated into the cholesterol oleate spot on the TLC plate with a Packard instainager.

Data from each of the noted compounds in the assays described above is as set forth in TABLES 5, 6, 7, and 8 as follows:

TABLE 5

COMPOUND	IC50 μM*	In vitro % Inhibition of TC Uptake @ 100 μM #	% Inhibition of Alanine Uptake @ 100 μM #	% of Control Transport of TC in Rat Ileum @ 0.1mM #
Benzothiazepine=	2		0	45.4 +/- 0.7
12		25		
3		0		
4a		3		
5a		34		
5b	40		0	72.9 ± 5.4 @ 0.5 mM
4b		9		
18		6		
14b		18		
14a		13		
13		23		
15	60			
19a		0		
19b		15		
8a		41		
Mixture of 8a and 8b		69		
Mixture of 9a and 9b	6			
6a	5			

331

6b		85		
9a	5		0% @ 25 mM	53.7 +/- 3.9
Mixture of 6a and 20	13			
Mixture of 6d and 10a	0.8		14% @ 25 mM	
21a		37		
21c		52		
21b		45		
6c	2		58.5 mM	68.8 +/- 5.7 at 0.4 mM
6d	0.6		77.7	16.4 +/- 1.1 @ 0.5 mM 30.2 +/- 0.9 @ 0.15 mM
17		10		
7	50		49.3	
10a	7		77.6	62.4 +/- 2.5 @ 0.2 mM
10b	15		68.6	
25	0.1		4% @ 10 mM	26.0 +/- 3.3
26	2		31% @ 25 mM	87.9 +/- 1.5
27	5		7% @ 20 mM	
28	8		31% @ 20mM	
29		88 @ 50 mM		
30		96 @ 50 mM		
31		41 @ 50 mM		
37	3		0% @ 5 mM	

332

38		0.3		11% @ 5mM	20.6 +/- 5.7
40			49 @ 50 mM		
41		2		0% @ 20 mM	
42		1.5			
43		1.5		16% @ 25 mM	
48		2		22% @ 20 mM	
49		0.15		21% @ 200 mM	21.2 +/- 2.7
57			51 @ 50 mM		
58			20 @ 50 mM		
59		70			
60		9		59	
61		30		175	
62		10			
63			90 @ 6 mM		
64			100 @ 6 mM		

* In vitro Taurocholate Cell Uptake

Unless otherwise noted

= Comparative Example is Example No. 1 in WO 93/16055

5

TABLE 6

Comp ound	TC-uptake (H14 cells)	TC-uptake Ileal Loop	TC-uptake IC(50)	TC-uptake (BBMV)	ACAT (liver)	ACAT intestine
	IC(50)	EC(50)	IC(50)	IC(50)	IC(50)	IC(50)
COMP. EXAMPLE*	1 mM	74 mM	3 mM	20 mM	20 mM	20 mM
6d	0.6 mM	31 mM	1.5 mM	25 mM	20 mM	20 mM
* 38	0.3 mM	12 mM	2 mM	15 mM	N.D.	N.D.
49	0.1 mM	12 mM	N.D.	6 mM	N.D.	N.D.
25	0.1 mM	20 mM	0.8 mM	8 mM	8 mM	8 mM

Comparative Example is Example No. 1 in WO 93/16055

5

TABLE 7 EFFICACY OF COMPOUND NO. 25 IN CHOLESTEROL-FED HAMSTERS				
PARAMETER	CONTROL	4% CHOLESTEROL TYRAMINE	0.2% CPD. NO. 25	
WEIGHT (G)	(mean ± SEM, *p<0.05, A-Student's t, B-Dunnett's)			
day 1	117	114(6)	117(5)	
day 14	(2)	127(3)	132(4)	
LIVER WEIGHT (G)	127(3)	4.9(0.4)	5.8(0.2)	
SER. CHOL(mg%)			126(2)*A	
HDL-CHOL(mg%)	5.4(0.3)	119(4)*A,B	76(1)*A,B	
VLDL + LDL				
TGI(mg%)	143(7)	76(3)*A,B		

333

334

HEPATIC CHOL (mg/g)	A	B	50 (3)
HMG COA (pm/mg/min.)	89 (4)	42 (3) *A	175 (11)
7a-OHase (pm/mg/min.)	54 (7)	190 (15)	1.9 (0.1)
24 HR. FECAL WT (g)	203 (3)	1.9 (0.1)	*A, B
FBA (mm/24H/100g)	2)	*A, B	312.9 (37.5) *A
	2.5 (0)	448.8 (2)	, B
	.3)	1.6) *A, B	
	15.8 (291.0 (6.
	7.6)	357.2 (2	0) *A
	235.3 (25.1	8.3) *A, B	2.4 (0.04
)	2.7 (0.1)
	2.3 (0	*A, B	11.9 (0.5
	.1)	12.3 (1.) *A, B
	6.2 (0	5) *A, B	
	.8)		

TABLE 8 EFFICACY OF COMPOUND NO. 25 IN RAT ALZET MINIPUMP MODEL		
PARAMETER	CONTROL	20 MP/L/DAY CPD. NO. 25
WEIGHT (g)	(mean \pm SEM, *p<0.05, A-Student's t, B-Dunnett's)	
day 1	307 (4)	307 (3)
day 8	330 (4)	310 (4) *A, B
LIVER WEIGHT (g)	15.5 (0.6)	14.6 (0.4)
SER. CHOL (mg%)	85 (3)	84 (3)
HEPATIC CHOL (mg/g)	21 (0.03)	2.0 (0.03)
HMG COA pm/mg/min	75.1 (6.4)	318.0
7a-OHase (pm/mg/min)	281.9 (13.9)	(40.7) *A, B
24 HR. FECAL WT (g)	5.8 (0.1)	535.2
FBA (mm/24H/100g)	17.9 (0.9)	(35.7) *A, B
		5.7 (0.4)
		39.1 (4.5) *A, B

Additional taurocholate uptake tests were conducted in 5 the following compounds listed in Table 9.

TABLE 9

Biological Assay Data for Some Compounds
of the Present Invention

Compound Number	Human TC IC ₅₀ (μ M)	Alanine Uptake Percent Inhibition @ μ M
101		0 @ 1.0
102	0.083	
103		13 @ 0.25
104	0.0056	
105	0.6	
106	0.8	
107		14.0 @ 0.063
108	0.3	
109		2.0 @ 0.063
110	0.09	
111	2.5	
112	3.0	
113	0.1	
114	0.19	
115	8.0	
116	0.3	
117		12.0 @ 0.625
118	0.4	
119	1.3	
120		34.0 @ 5.0
121	0.068	
122	1.07	
123	1.67	
124		14.0 @ 6.25
125	18.0	
126		18 @ 1.25
127	0.55	
128	0.7	
129	0.035	
131	1.28	
132		5.4 @ 0.063
133	16.0	
134	0.3	
135	22.0	
136	0.09	

337

137	2.4	
138	3.0	
139	>25.0	
142	0.5	
143	0.03	
144	0.053	
262	0.07	
263	0.7	
264	0.2	
265	2.0	
266	0.5	
267	0.073	
268	0.029	
269	0.08	
270	0.12	
271	0.07	
272	0.7	
273	1.9	
274	0.18	
275		5.0 @ 0.25
276	0.23	
277	0.04	
278	3.0	
279	0.4	
280	0.18	
281	0.019	
282	0.021	
283	0.35	
284	0.08	
286	19.0	
287	4.0	
288		10.0 @ 6.25
289	0.23	
290	0.054	
291	0.6	
292	0.046	
293	1.9	
294	0.013	
295	1.3	
296	1.6	
1005	0.0004	
1006	0.001	

338

1007	0.001	
1008	0.001	
1009	0.001	
1010	0.001	
1011	0.001	
1012	0.0015	
1013	0.002	
1014	0.002	
1015	0.002	
1016	0.002	
1017	0.002	
1018	0.002	
1019	0.002	
1020	0.002	
1021	0.002	
1022	0.002	
1023	0.002	
1024	0.002	
1025	0.002	
1026	0.002	
1027	0.002	
1028	0.002	
1029	0.002	
1030	0.002	
1031	0.002	
1032	0.002	
1033	0.002	
1034	0.002	
1035	0.002	
1036	0.002	
1037	0.0022	
1038	0.0025	
1039	0.0026	
1040	0.003	
1041	0.003	
1042	0.003	
1043	0.003	
1044	0.003	
1045	0.003	
1046	0.003	
1047	0.003	
1048	0.003	

339

1049	0.003	
1050	0.003	
1051	0.003	
1052	0.003	
1053	0.003	
1054	0.003	
1055	0.003	
1056	0.003	
1057	0.003	
1058	0.003	
1059	0.003	
1060	0.0036	
1061	0.004	
1062	0.004	
1063	0.004	
1064	0.004	
1065	0.004	
1066	0.004	
1067	0.004	
1068	0.004	
1069	0.004	
1070	0.004	
1071	0.004	
1072	0.004	
1073	0.004	
1074	0.004	
1075	0.0043	
1076	0.0045	
1077	0.0045	
1078	0.0045	
1079	0.005	
1080	0.005	
1081	0.005	
1082	0.005	
1083	0.005	
1084	0.005	
1085	0.005	
1086	0.005	
1087	0.005	
1088	0.0055	
1089	0.0057	
1090	0.006	

340

1091	0.006	
1092	0.006	
1093	0.006	
1094	0.006	
1095	0.006	
1096	0.006	
1097	0.006	
1098	0.006	
1099	0.0063	
1100	0.0068	
1101	0.007	
1102	0.007	
1103	0.007	
1104	0.007	
1105	0.007	
1106	0.0073	
1107	0.0075	
1108	0.0075	
1109	0.008	
1110	0.008	
1111	0.008	
1112	0.008	
1113	0.009	
1114	0.009	
1115	0.0098	
1116	0.0093	
1117	0.01	
1118	0.01	
1119	0.01	
1120	0.01	
1121	0.01	
1122	0.011	
1123	0.011	
1124	0.011	
1125	0.012	
1126	0.013	
1127	0.013	
1128	0.017	
1129	0.018	
1130	0.018	
1131	0.02	
1132	0.02	

341

1133	0.02	
1134	0.02	
1135	0.021	
1136	0.021	
1137	0.021	
1138	0.022	
1139	0.022	
1140	0.023	
1141	0.023	
1142	0.024	
1143	0.027	
1144	0.028	
1145	0.029	
1146	0.029	
1147	0.029	
1148	0.03	
1149	0.03	
1150	0.03	
1151	0.031	
1152	0.036	
1153	0.037	
1154	0.037	
1155	0.039	
1156	0.039	
1157	0.04	
1158	0.06	
1159	0.06	
1160	0.062	
1161	0.063	
1162	0.063	
1163	0.09	
1164	0.093	
1165	0.11	
1166	0.11	
1167	0.12	
1168	0.12	
1169	0.12	
1170	0.13	
1171	0.14	
1172	0.14	
1173	0.15	
1174	0.15	

342

1175	0.17	
1176	0.18	
1177	0.18	
1178	0.19	
1179	0.19	
1180	0.2	
1181	0.22	
1182	0.25	
1183	0.28	
1184	0.28	
1185	0.28	
1186	0.3	
1187	0.32	
1188	0.35	
1189	0.35	
1190	0.55	
1191	0.65	
1192	1.0	
1193	1.0	
1194	1.6	
1195	1.7	
1196	2.0	
1197	2.2	
1198	2.5	
1199	4.0	
1200	6.1	
1201	8.3	
1202	40.0	
1203		0 @ 0.063
1204	0.05	
1205	0.034	
1206	0.035	
1207	0.068	
1208	0.042	
1209		0 @ 0.063
1210	0.14	
1211	0.28	
1212	0.39	
1213	1.7	
1214	0.75	
1215	0.19	
1216	0.39	

343

1217	0.32	
1218	0.19	
1219	0.34	
1220	0.2	
1221	0.041	
1222	0.065	
1223	0.28	
1224	0.33	
1225	0.12	
1226	0.046	
1227	0.25	
1228	0.038	
1229	0.049	
1230	0.062	
1231	0.075	
1232	1.2	
1233	0.15	
1234	0.067	
1235	0.045	
1236	0.05	
1237	0.07	
1238	0.8	
1239	0.035	
1240	0.016	
1241	0.047	
1242	0.029	
1243	0.63	
1244	0.062	
1245	0.32	
1246	0.018	
1247	0.017	
1248	0.33	
1249	10.2	
1250	0.013	
1251	0.62	
1252	29.	
1253	0.3	
1254	0.85	
1255	0.69	
1256	0.011	
1257	0.1	
1258	0.12	

344

1259	16.5	
1260	0.012	
1261	0.019	
1262	0.03	
1263	0.079	
1264	0.21	
1265	0.24	
1266	0.2	
1267	0.29	
1268	0.035	
1269	0.026	
1270	0.026	
1271	0.011	
1272	0.047	
1273	0.029	
1274	0.028	
1275	0.024	
1276	0.029	
1277	0.018	
1278	0.017	
1279	0.028	
1280	0.76	
1281	0.055	
1282	0.17	
1283	0.17	
1284	0.011	
1285	0.027	
1286	0.068	
1287	0.071	
1288	0.013	
1289	0.026	
1290	0.017	
1291	0.013	
1292	0.025	
1293	0.019	
1294	0.011	
1295	0.014	
1296	0.063	
1297	0.029	
1298	0.018	
1299	0.012	
1300	1.0	

345

1301	0.15	
1302	1.4	
1303	0.26	
1304	0.25	
1305	0.25	
1306	1.2	
1307	3.1	
1308	0.04	
1309	0.24	
1310	1.16	
1311	3.27	
1312	5.0	
1313	6.1	
1314	0.26	
1315	1.67	
1316	3.9	
1317	21.0	
1319		11.0 @ 0.25
1321		11.1 @ 5.0
1322		3.0 @ 0.0063
1323		4.0 @ 0.0063
1324		43.0 @ 0.0008
1325		1.0 @ 0.0063
1326		36.0 @ 0.0008
1327		3.0 @ 0.0063
1328		68.0 @ 0.0063
1329		2.0 @ 0.0063
1330		9.0 @ 0.0063
1331		57.0 @ 0.0008
1332		43.0 @ 0.0008
1333		0 @ 0.0063
1334		50.0 @ 0.0008
1335		38.0 @ 0.0008
1336		45.0 @ 0.0008
1337		0 @ 0.0063
1338		1.0 @ 0.25
1339		0 @ 0.063
1340		9.0 @ 0.063
1341		1.0 @ 0.063
1342		1.0 @ 0.063
1345		13.0 @ 0.25
1347	0.0036	

346


1351	0.44	
1352	0.10	
1353	0.0015	
1354	0.006	
1355	0.0015	
1356	0.22	
1357	0.023	
1358	0.008	
1359	0.014	
1360	0.003	
1361	0.004	
1362	0.019	
1363	0.008	
1364	0.006	
1365	0.008	
1366	0.015	
1367	0.002	
1368	0.005	
1369	0.005	
1370	0.002	
1371	0.004	
1372	0.004	
1373	0.008	
1374	0.007	
1375	0.002	
1449	0.052	
1450	0.039	
1451	0.014	

The examples herein can be repeated with similar success by substituting the generically or specifically described reactants and/or operating conditions of this invention for those used in the preceding examples.

Novel compositions of the invention are further illustrated in attached Exhibits A and B.

The invention being thus described, it is apparent that the same can be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the present invention, and all such modifications and equivalents as would be obvious to one skilled in the art are intended to be included within the scope of the following claims.











Table C2: Alternative compounds #2 (Families F101-F123)



Family Cpd#	R ¹ =R ²	R ³	(R ³) _q
F101	CHOSEN FROM TABLE D *	Ph-	CHOSEN FROM TABLE D
F102	CHOSEN FROM TABLE D	p-F-Ph-	CHOSEN FROM TABLE D
F103	CHOSEN FROM TABLE D	m-F-Ph-	CHOSEN FROM TABLE D
F104	CHOSEN FROM TABLE D	p-CH ₃ O-Ph-	CHOSEN FROM TABLE D
F105	CHOSEN FROM TABLE D	m-CH ₃ O-Ph-	CHOSEN FROM TABLE D
F106	CHOSEN FROM TABLE D	p-(CH ₃) ₂ N-Ph-	CHOSEN FROM TABLE D
F107	CHOSEN FROM TABLE D	m-(CH ₃) ₂ N-Ph	CHOSEN FROM TABLE D
F108	CHOSEN FROM TABLE D	I ⁻ , p-(CH ₃) ₃ N ⁺ -Ph-	CHOSEN FROM TABLE D
F109	CHOSEN FROM TABLE D	I ⁻ , m-(CH ₃) ₃ N ⁺ -Ph-	CHOSEN FROM TABLE D
F110	CHOSEN FROM TABLE D	I ⁻ , p-(CH ₃) ₃ N ⁺ -CH ₂ CH ₂ -(OCH ₂ CH ₂) ₂ O-Ph-	CHOSEN FROM TABLE D
F111	CHOSEN FROM TABLE D	I ⁻ , m-(CH ₃) ₃ N ⁺ -CH ₂ CH ₂ -(OCH ₂ CH ₂) ₂ O-Ph-	CHOSEN FROM TABLE D
F112	CHOSEN FROM TABLE D	I ⁻ , p-(N,N-dimethylpiperazine)-(N')-CH ₂ -(OCH ₂ CH ₂) ₂ O-Ph-	CHOSEN FROM TABLE D

F113	CHOSEN FROM TABLE D	I ⁻ , m-(N,N-dimethylpiperazine)-(N')-CH ₂ -(OCH ₂ CH ₂) ₂ O-Ph-	CHOSEN FROM TABLE D
F114	CHOSEN FROM TABLE D	m-F-Ph-p-CH ₃ O-	CHOSEN FROM TABLE D
F115	CHOSEN FROM TABLE D	3,4,dioxy-methylene-Ph-	CHOSEN FROM TABLE D
F116	CHOSEN FROM TABLE D	m-F-Ph-p-F-Ph-	CHOSEN FROM TABLE D
F117	CHOSEN FROM TABLE D	m-CH ₃ O-p-F-Ph-	CHOSEN FROM TABLE D
F118	CHOSEN FROM TABLE D	4-pyridine	CHOSEN FROM TABLE D
F119	CHOSEN FROM TABLE D	N-methyl-4-pyridinium	CHOSEN FROM TABLE D
F120	CHOSEN FROM TABLE D	3-pyridine	CHOSEN FROM TABLE D
F121	CHOSEN FROM TABLE D	N-methyl-3-pyridinium	CHOSEN FROM TABLE D
F122	CHOSEN FROM TABLE D	2-pyridine	CHOSEN FROM TABLE D
F123	CHOSEN FROM TABLE D	p-CH ₃ O ₂ C-Ph-	CHOSEN FROM TABLE D

Similar families can be generated where R¹<R², such as R¹ = Et and R² = n-Bu, but (R³)_q is chosen from table C1.



R^1 and R^2 taken together with the carbon to which they are attached form C_{10} cycloalkylidene;

R^3 and R^4 are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, acyloxy, aryl, heterocycle, OR^9 , NR^9R^{10} , SR^9 , $S(O)R^9$, SO_2R^9 , and SO_3R^9 , wherein R^9 and R^{10} are as defined above; or

R^3 and R^4 together form $=O$, $=NOR^{11}$, $=S$, $=NNR^{11}R^{12}$, $=NR^9$, or $=CR^{11}R^{12}$,

wherein R^{11} and R^{12} are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl, heterocycle, carboxyalkyl, carbalkoxyalkyl, cycloalkyl, cyanoalkyl, OR^9 , NR^9R^{10} , SR^9 , $S(O)R^9$, SO_2R^9 , SO_3R^9 , CO_2R^9 , CN, halogen, oxo, and $CONR^9R^{10}$,

wherein R^9 and R^{10} are as defined above, provided that both R^3 and R^4 cannot be OH, NH_2 , and SH, or

R^{11} and R^{12} together with the nitrogen or carbon atom to which they are attached form a cyclic ring;

R^5 and R^6 are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, OR^9 , SR^9 , $S(O)R^9$, SO_2R^9 , and SO_3R^9 ,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, and quaternary heteroaryl can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, halogen, oxo, OR^{13} , $NR^{13}R^{14}$, SR^{13} , $S(O)R^{13}$, SO_2R^{13} , SO_3R^{13} , $NR^{13}OR^{14}$, $NR^{13}NR^{14}R^{15}$, NO_2 , CO_2R^{13} , CN, OM, SO_2OM , $SO_2NR^{13}R^{14}$, $C(O)NR^{13}R^{14}$, $C(O)OM$, COR^{13} ,

355

$P(O)R^{13}R^{14}$, $P^+R^{13}R^{14}R^{15}A^-$, $P(OR^{11})OR^{11}$, $SR^{11}R^{11}A^-$, and $N^+R^9R^{11}R^{12}A^-$,

wherein:

A^- is a pharmaceutically acceptable anion and M is a pharmaceutically acceptable cation,

said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of OR^7 , NR^7R^8 , SR^7 , $S(O)R^7$, SO_2R^7 , SO_3R^7 , CO_2R^7 , CN, oxo, $CONR^7R^8$, $N^+R^7R^8R^9A^-$, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, $P(O)R^7R^8$, $P^+R^7R^8R^9A^-$, and $P(O)(OR^7)OR^8$, and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O, NR^7 , $N^+R^7R^8A^-$, S, SO, SO_2 , $S^+R^7A^-$, PR^7 , $P(O)R^7$, $P^+R^7R^8A^-$, or phenylene, and R^{13} , R^{14} , and R^{15} are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, polyalkyl, aryl, arylalkyl, cycloalkyl, heterocycle, heterocycle, quaternary heterocycle, quaternary heteroaryl, and quaternary heteroarylalkyl.

wherein alkyl, alkenyl, alkynyl, arylalkyl, heterocycle, and polyalkyl optionally have one or more carbons replaced by O, NR^9 , $N^+R^9R^{10}A^-$, S, SO, SO_2 , $S^+R^9A^-$, PR^9 , $P^+R^9R^{10}A^-$, $P(O)R^9$, phenylene, carbonyl, amino acid, peptide, or polypeptide, and

R^{13} , R^{14} , and R^{15} are optionally substituted with one or more groups selected from the group consisting of sulfoalkyl, quaternary heterocycle, quaternary heteroaryl, OR^9 , NR^9R^{10} , $N^+R^9R^{11}R^{12}A^-$, SR^9 , $S(O)R^9$,

356

SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰, SO₂OM, SO₂NR⁹R¹⁰, PO(OR¹⁶)OR¹⁷, P⁺R⁹R¹⁰A⁻, S⁺R⁹A⁻, and C(O)OM, wherein R¹⁶ and R¹⁷ are independently selected

from the substituents constituting R⁹ and M; or

R¹⁴ and R¹⁵, together with the nitrogen atom to

which they are attached, form a cyclic ring;

R⁷ and R⁸ are independently selected from the group consisting of hydrogen and alkyl; and

one or more R^x are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl,

polyalkyl, acyloxy, aryl, arylalkyl, halogen,

haloalkyl, cycloalkyl, heterocycle, heterocycle,

polyether, quaternary heterocycle, quaternary

heteroaryl, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, S(O)₂R¹³,

SO₃R¹³, S⁺R¹³A⁻, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, NR¹³C(O)R¹⁴, C(O)NR¹³R¹⁴,

NR¹⁴C(O)R¹³, C(O)OM, COR¹³, OR¹⁸, S(O)NR¹⁸, NR¹³R¹⁸,

NR¹⁸OR¹⁴, N⁺R⁹R¹²A⁻, P⁺R⁹R¹²A⁻, amino acid,

peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl,

polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl,

polyether, quaternary heterocycle, and quaternary

heteroaryl can be further substituted with OR⁹, NR⁹R¹⁰,

N⁺R⁹R¹²A⁻, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹,

CN, halogen, CONR⁹R¹⁰, SO₂OM, SO₂NR⁹R¹⁰, PO(OR¹⁶)OR¹⁷,

P⁺R⁹R¹²A⁻, S⁺R⁹A⁻, or C(O)OM, and

wherein R¹⁸ is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heterocycle, alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl,

heterocycle, heterocycle, alkyl quaternary heterocycle,

and quaternary heteroaryl optionally are substituted

with one or more substituent selected from the group

357

consisting of OR⁹, NR⁹R¹⁰, N⁺R⁹R¹²A⁻, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰, SO₃R⁹, SO₂OM, SO₂NR⁹R¹⁰, PO(OR¹⁶)OR¹⁷, and C(O)OM,

wherein in R^x, one or more carbons are optionally

replaced by O, NR¹³, N⁺R¹³R¹⁴A⁻, S, SO, SO₂, S⁺R¹³A⁻,

PR¹³, P(OR¹³), P⁺R¹³R¹⁴A⁻, phenylene, amino acid,

peptide, polypeptide, carbohydrate, polyether, or polyalkyl,

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more

carbons are optionally replaced by O, NR⁹, N⁺R⁹R¹⁰A⁻,

S, SO, SO₂, S⁺R⁹A⁻, PR⁹, P⁺R⁹R¹⁰A⁻, or P(O)R⁹;

wherein quaternary heterocycle and quaternary

heteroaryl are optionally substituted with one or more groups selected from the group consisting of alkyl,

alkenyl, alkynyl, polyalkyl, polyether, aryl,

haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen,

oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, SO₃R¹³,

NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM,

SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴,

P⁺R¹³R¹⁴R¹⁵A⁻, P(OR¹³)OR¹⁴, S⁺R¹³R¹⁴A⁻, and N⁺R⁹R¹⁰R¹²A⁻,

provided that both R⁵ and R⁶ cannot be hydrogen,

OH, or SH and when R⁵ is OH, R¹, R², R³, R⁴, R⁷ and R⁸ cannot be all hydrogen;

provided that when R¹ or R⁴ is phenyl, only one of R¹ or R⁴ is H;

provided that when q = 1 and R¹ is styryl,

anilido, or anilino, only one of R¹ or R⁴ is alkyl.

2. A compound of claim 1, wherein R⁵ and R⁶ are independently selected from the group consisting of H,

358

aryl, heterocycle, quaternary heterocycle, and quaternary heteroaryl,

wherein said aryl, heteroaryl, quaternary heterocycle, and quaternary heteroaryl can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen,

oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴, P⁺R¹³R¹⁴R¹⁵A⁻, P(OR¹³)OR¹⁴, S⁺R¹³R¹⁴A⁻, and N⁺R⁷R⁸R⁹R¹²A⁻,

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O, NR⁷, N⁺R⁷R⁸A⁻, S, SO, SO₂, S⁺R⁷A⁻, PR⁷, P(O)R⁷, P⁺R⁷R⁸A⁻, or phenylene,

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of OR⁷, NR⁷R⁸, SR⁷, S(O)R⁷, SO₂R⁷, SO₃R⁷, CO₂R⁷, CN, oxo,

CONR⁷R⁸, N⁺R⁷R⁸R⁹A⁻, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, P(O)R⁷R⁸, P⁺R⁷R⁸A⁻, and P(O)(OR⁷)OR⁸.

3. A compound of claim 2, wherein R¹ or R¹ has the formula



wherein:

t is an integer from 0 to 5;

359

Ar is selected from the group consisting of phenyl, thiophenyl, pyridyl, piperazinyl, piperonyl, pyrrolyl, naphthyl, furanyl, anthracenyl, quinolinyl, isochinolinyl, quinoxalinyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, pyrimidinyl, thiazolyl, triazolyl, isothiazolyl, indolyl, benzimidazolyl, benzoxazolyl, benzothiazolyl, and benzoisothiazolyl; and

one or more R¹ are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, OR⁹, SR⁹, S(O)R⁹, SO₂R⁹, and SO₃R⁹,

wherein said alkyl, alkenyl, alkynyl, aryl, cycloalkyl, and heterocycle can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴, P⁺R¹³R¹⁴R¹⁵A⁻, P(OR¹³)OR¹⁴, S⁺R¹³R¹⁴A⁻, and N⁺R⁹R¹¹R¹²A⁻,

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of OR⁷, NR⁷R⁸, SR⁷, S(O)R⁷, SO₂R⁷, SO₃R⁷, CO₂R⁷, CN, oxo, CONR⁷R⁸, N⁺R⁷R⁸R⁹A⁻, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, P(O)R⁷R⁸, P⁺R⁷R⁸A⁻, and P(O)(OR⁷)OR⁸, and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O,

360

NR⁷, N⁷R⁷R⁸A⁻, S, SO, SO₂, S⁷R⁷A⁻, PR⁷, P(O)R⁷,
P⁷R⁷R⁸A⁻, or phenylene.

4. A compound of claim 3, wherein R⁵ or R⁶ has
the formula (II)



5. A compound of claim 4, wherein n is 1 or 2.

6. A compound of claim 5, wherein one of R⁷ or
R⁸ is H and the other of R⁷ or R⁸ is alkyl.

7. A compound of claim 5, wherein both R⁷ and R⁸
are H.

8. A compound of claim 7, wherein R¹ and R² are
independently selected from the group consisting of H
and alkyl.

9. A compound of claim 8, wherein said alkyl is
a C₁-C₁₀ alkyl.

10. A compound of claim 8, wherein R¹ and R² are
both alkyl.

11. A compound of claim 10, wherein said alkyl is
a C₁-C₁₀ alkyl.

12. A compound of claim 11, wherein said alkyl is
a C₁-C₁₀ alkyl.

13. A compound of claim 12, wherein said alkyl is
a C₁-C₁₀ alkyl.

14. A compound of claim 13, wherein said alkyl is
independently selected from the group consisting of
ethyl, n-propyl, n-butyl, and isobutyl.

15. A compound of claim 8, wherein R¹ and R² are
each n-butyl.

16. A compound of claim 8, wherein one of R¹ and
R² is ethyl and the other of R¹ and R² is n-butyl.

17. A compound of claim 15, wherein q is 1, 2, or
3.

18. A compound of claim 16, wherein q is 1, 2, or
3.

19. A compound of claim 17, wherein q is 1 or 2.

20. A compound of claim 19, wherein q is 1.

21. A compound of claim 18, wherein q is 1 or 2.

22. A compound of claim 21, wherein q is 1.

23. A compound of claim 19, wherein R¹ and R² are
independently selected from the group consisting of H
and OR¹.

24. A compound of claim 21, wherein R¹ and R² are
independently selected from the group consisting of H
and OR¹.

25. A compound of claim 23, wherein R' is H.

26. A compound of claim 24, wherein R' is H.

27. A compound of claim 25, wherein one or more R^x are in the 7-, 8-, or 9-position of the benzo ring of formula (I).

28. A compound of claim 26, wherein said R' is in the 7-, 8-, or 9- position of the benzo ring of formula (I).

29. A compound of claim 27, wherein said R' are in the 7- and 9- positions of the benzo ring of formula (I).

30. A compound of claim 28, wherein said R^x is in the 7-position of the benzo ring of formula (I).

31. A compound of claim 29, wherein said one or more R^x are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen, OR¹³, NR¹³R¹⁴, NR¹³NR¹⁴R¹⁵, N⁺R⁹R¹¹R¹²A⁻, SR¹³, S⁺R¹³R¹⁴, CO₂R¹³, NR¹³C(O)R¹⁴, and NR¹³C(O)R¹⁵,

wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with OR⁹, NR⁹R¹⁰, N⁺R⁹R¹¹R¹²A⁻, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰SO₂OM, SO₂NR⁹R¹⁰, PO(OR¹⁰)OR⁹, P⁺R⁹R¹¹R¹²A⁻, S⁺R⁹R¹⁰A⁻, or C(O)OM, and

wherein in R^x, one or more carbons are optionally replaced by O, NR¹³, N⁺R¹³R¹⁴A⁻, S, SO, SO₂, S⁺R¹³A⁻, PR¹³, P(O)R¹³, P⁺R¹³R¹⁴A⁻, phenylene, amino acid,

363

peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR⁹, N⁺R⁹R¹⁰A⁻, S, SO, SO₂, S⁺R⁹A⁻, PR⁹, P⁺R⁹R¹⁰A⁻, or P(O)R⁹.

32. A compound of claim 30, wherein said R^x is selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen, OR¹³, NR¹³R¹⁴, NR¹³NR¹⁴R¹⁵, N⁺R⁹R¹¹R¹²A⁻, SR¹³, S⁺R¹³R¹⁴, CO₂R¹³, NR¹³C(O)R¹⁴, and NR¹³C(O)R¹⁵, wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with OR⁹, NR⁹R¹⁰, N⁺R⁹R¹¹R¹²A⁻, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰SO₂OM, SO₂NR⁹R¹⁰, PO(OR¹⁰)OR⁹, P⁺R⁹R¹¹R¹²A⁻, S⁺R⁹R¹⁰A⁻, or C(O)OM, and

wherein in R^x, one or more carbons are optionally replaced by O, NR¹³, N⁺R¹³R¹⁴A⁻, S, SO, SO₂, S⁺R¹³A⁻, PR¹³, P(O)R¹³, P⁺R¹³R¹⁴A⁻, phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR⁹, N⁺R⁹R¹⁰A⁻, S, SO, SO₂, S⁺R⁹A⁻, PR⁹, P⁺R⁹R¹⁰A⁻, or P(O)R⁹.

33. A compound of claim 31, wherein said one or more R^x are independently selected from the group consisting of polyether, OR¹³, NR¹³R¹⁴, and N⁺R⁹R¹¹R¹²A⁻.

364

34. A compound of the claim 32, wherein said Rx is selected from the group consisting of polyether, OR", NR"R", and N⁺R⁹R¹¹R¹²A⁻.

35. A compound of claim 33, wherein said one or more Rx are independently selected from the group consisting of OR" and NR"R".

36. A compound of claim 34, wherein said R' is independently selected from the group consisting of OR" and NR"R".

37. A compound of claim 35, wherein R" and R" each methyl.

38. A compound of the claim 36, wherein R" and R" each methyl.

39. A compound of claim 31, wherein one or more R^y are independently in the 3- or the 4-position of the phenyl ring of formula (II).

40. A compound of claim 32, wherein one or more Ry are independently in the 3- or the 4- position of the phenyl ring of formula (II).

41. A compound of claim 39, wherein t is 1 or 2.

42. A compound of claim 40, wherein t is 1 or 2.

43. A compound of claim 41, wherein said one or more R^y are independently selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide, NR¹³R¹⁴, NR"C(O)R", and OR",

wherein alkyl and polyether can be further substituted with SO₃R⁹, N⁺R⁹R¹¹R¹²A⁻, and quaternary heteroaryl.

44. A compound of claim 42, wherein said R^y is independently selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide, NR¹³R¹⁴, NR"C(O)R", and OR",

wherein alkyl and polyether can be further substituted with SO₃R⁹, N⁺R⁹R¹¹R¹²A⁻, and quaternary heteroaryl.

45. A compound of claim 43, wherein said one or more R^y are independently selected from the group consisting of alkyl, polyether, fluoride, NR¹³R¹⁴, NR"C(O)R", and OR",

wherein alkyl and polyether can be further substituted with SO₃R⁹, N⁺R⁹R¹¹R¹²A⁻, and quaternary heteroaryl.

46. A compound of claim 44 wherein said R^y is independently selected from the group consisting of alkyl, polyether, fluoride, NR¹³R¹⁴, NR"C(O)R", and OR",

wherein alkyl and polyether can be further substituted with SO₃R⁹, N⁺R⁹R¹¹R¹²A⁻, and quaternary heteroaryl.

47. A compound of claim 45, wherein said R" and R" are alkyl,

wherein alkyl can be further substituted with SO₃R⁹, N⁺R⁹R¹¹R¹²A⁻, and quaternary heteroaryl.

48. A compound of claim 46, wherein said R' and R' are alkyl,

wherein alkyl can be further substituted with SO₃R⁹, N⁺R⁹R¹¹R¹²A⁻, and quaternary heteroaryl.


49. A compound of claim 47, wherein n is 2.

50. A compound of claim 48, wherein n is 2.


51. A compound of claim 49, wherein said OH group is in a syn relationship to said structure of formula (II).

52. A compound of claim 50, wherein said OH group is in a syn relationship to said structure of formula (II).


53. A compound of claim 51, having the formula:




54. A compound of claim 51, having the formula:



55. A compound of claim 51, having the formula:



56. A compound of claim 51, having the formula:




57. A compound of claim 51, having the formula:


10

369


58. A compound of claim 52, having the formula:



59. A compound of claim 52, having the formula:




60. A compound of claim 52, having the formula:




10


370



61. A compound of claim 52, having the formula:




62. A compound of claim 52, having the formula:



63. A compound of claim 31, wherein n is 1.

64. A compound of claim 63, wherein R' is H.

65. A compound of claim 64, having the formula



66. A compound of claim 4, wherein R' and R' are independently selected from the group consisting of H and alkyl.

67. A compound of claim 66, wherein said alkyl is C₁-C₆, alkyl.

68. A compound of claim 67, wherein said alkyl is C₂-C₆, alkyl.

69. A compound of claim 68, wherein said alkyl is C₁-C₄ alkyl.

70. A compound of claim 69, wherein R' and R' are independently selected from the group consisting of ethyl, n-propyl, n-butyl, and isobutyl.

71. A compound of claim 4, wherein R' and R' are independently selected from the group consisting of H and OR'.

72. A compound of claim 71, wherein R' is H.

73. A compound of claim 4, wherein n is 2.

74. A compound of claim 3, wherein R' and R' are independently selected from the group consisting of H and OR'.

75. A compound of claim 74, wherein R⁹ is H.

76. A compound of claim 3, wherein one of R⁷ or R⁸ is H.

77. A compound of claim 76, wherein both R' and R' are H.

78. A compound of claim 3, wherein said one or more R^x are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen, OR¹³, NR¹³, NR¹⁴, NR¹⁵, NR⁹, NR¹¹, NR¹², SR¹³, SR¹⁴, SR¹⁵, CO₂R¹³, NR¹³(O)R', and NR¹³(O)R'',

wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with OR⁹, NR⁹, NR¹⁰, NR¹¹, NR¹², SR⁹,

S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹, SO₂OM, SO₂NR⁹, PO(OR⁹)OR'', P⁹R⁹R¹¹R¹², S⁹R⁹A', or C(O)OM, and

wherein in R^x, one or more carbons are optionally replaced by O, NR¹³, NR¹³R¹⁴A-, S, SO, SO₂, S⁹R¹³A-, PR¹³, P(O)R'', P⁹R¹³R¹⁴A', phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR⁹, NR⁹R¹⁰A-, S, SO, SO₂, S⁹R⁹A-, PR⁹, P⁹R⁹R¹⁰A-, or P(O)R'.

79. A compound of claim 78, wherein said one or more R' are independently selected from the group consisting of polyether, OR'', NR''R'', and N⁹R⁹R¹¹R¹²A-.

80. A compound of claim 79, wherein said one or more R' are independently selected from the group consisting of OR'' and NR''R''.

81. A compound of claim 80, wherein R'' and R'' are each methyl.

82. A compound of claim 3, wherein one or more R^y are independently in the 3- or the 4-position of the phenyl ring of formula (II).

83. A compound of claim 82, wherein one or more R^y is selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide, NR⁹, and NC(O)R⁹,

wherein alkyl and polyether can be substituted with SO₃R⁹, NR⁹R¹¹R¹²A-, and quaternary heteroaryl.

84. A compound of claim 83, wherein R^9 and R^{10} are alkyl.

85. A compound of claim 84, wherein one or more R^9 is selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide, NR^9R^{10} , and $NC(O)R^9$.

86. A compound of claim 1, wherein said one or more R^X are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen, OR^{13} , $NR^{13}R^{14}$, $NR^{13}NR^{14}R^{15}$, $N^+R^9R^{11}R^{12}A^-$, SR^{13} , $S^+R^{13}R^{14}$, CO_2R^{13} , $NR^9C(O)R^9$, and $NR^9C(O)R^9$,

wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with OR^9 , NR^9R^{10} , $N^+R^9R^{11}R^{12}A^-$, SR^9 , $S(O)R^9$, SO_2R^9 , SO_3R^9 , oxo, CO_2R^9 , CN, halogen, $CONR^9R^{10}$, SO_2OM , $SO_2NR^9R^{10}$, $PO(OR^9)OR^9$, $P^+R^9R^{11}R^{12}A^-$, $S^+R^9R^{14}A^-$, or $C(O)OM$, and

wherein in R^X , one or more carbons are optionally replaced by O, NR^{13} , $N^+R^{13}R^{14}A^-$, S, SO, SO_2 , $S^+R^{13}A^-$, PR^{13} , $P(O)R^{13}$, $P^+R^{13}R^{14}A^-$, phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR^9 , $N^+R^9R^{10}A^-$, S, SO, SO_2 , $S^+R^9A^-$, PR^9 , $P^+R^9R^{10}A^-$, or $P(O)R^9$.

87. A compound of claim 1, wherein n is 1 or 2.

375

88. A compound of claim 87, wherein n is 2.

89. A compound of claim 1, wherein R^1 and R^2 are independently selected from the group consisting of H and alkyl.

90. A compound of claim 89, wherein said alkyl is C_1-C_{10} alkyl.

91. A compound of claim 90, wherein said alkyl is C_1-C_2 alkyl.

92. A compound of claim 91, wherein said alkyl is C_1-C_2 alkyl.

93. A compound of claim 92, wherein R^1 and R^2 are independently selected from the group consisting of ethyl, n-propyl, n-butyl, and isobutyl.

94. A compound of claim 1, wherein R^1 and R^2 are independently selected from the group consisting of H and OR^1 .


95. A compound of claim 94, wherein R^1 is H.

96. A compound of claim 1, wherein one of R^7 or R^8 is H.

97. A compound of claim 96, wherein both R^7 and R^8 are H.

98. A compound of the formula (III)

376



wherein :

q and r are independently integers from 0 to 4;

d and e are independently integers from 0 to 2;

t and u are independently integers from 0 to 4;

R¹, R², R³, and R⁴ are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl.

10

wherein alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more

15

substituent selected from the group consisting of OR⁹, NR⁹R¹⁰, N⁺R⁹R¹⁰, SR⁹, S⁺R⁹, P⁺R⁹R¹⁰, S(O)R⁹, SO₂R⁹, CO₂R⁹, CN, halogen, oxo, and CONR⁹R¹⁰, wherein alkyl, alkenyl, alkynyl, alkylaryl, alkoxy, alkoxyalkyl, polyalkyl, aryl, and cycloalkyl

20

optionally have one or more carbons replaced by O, NR⁹, N⁺R⁹R¹⁰, S, SO, SO₂, S⁺R⁹, P⁺R⁹R¹⁰, or phenylene, wherein R⁹, R¹⁰, and R^w are independently selected

from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle,

25

ammoniumalkyl, alkylammoniumalkyl, and arylalkyl; or R¹ and R² taken together with the carbon to which they are attached form C₁-C₁₀ cycloalkylidene, or R³ and R⁴ taken together with the carbon to which they are attached form C₁-C₁₀ cycloalkylidene;

377

R¹, R², R³, and R⁴ are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, acyloxy, aryl, heterocycle, OR⁹, NR⁹R¹⁰, SR⁹, S(O)R⁹, SO₂R⁹, and SO₃R⁹, wherein R¹ and R² are as defined above; or

5

R³ and R⁴ together form =O, =NOR¹¹, =S, =NNR¹¹R¹², =NR⁹, or =CR¹¹R¹², or

R^{3A} and R^{4A} together form =O, =NOR¹¹, =S, =NNR¹¹R¹², =NR⁹, or =CR¹¹R¹²,

10

wherein R¹¹ and R¹² are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl, heterocycle, carboxyalkyl, carboalkoxyalkyl, cycloalkyl, cyanoalkyl, OR⁹, NR⁹R¹⁰, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, CO₂R⁹, CN, halogen, oxo, and CONR⁹R¹⁰,

15

wherein R⁹ and R¹⁰ are as defined above, provided that both R³ and R⁴ cannot be OH, NH₂, and SH, or

20

R¹¹ and R¹² together with the nitrogen or carbon atom to which they are attached form a cyclic ring;

wherein A⁻ is a pharmaceutically acceptable anion and M is a pharmaceutically acceptable cation;

R⁷, R⁸, R⁹, and R¹⁰ are independently selected from the group consisting of hydrogen and alkyl; and

25

one or more R^x and R^y are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heterocycle, polyether, quaternary heterocycle, quaternary heteroaryl, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, S⁺R¹³R¹⁴, NR¹³OR¹⁴, NR¹³NR¹⁴, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, NR¹⁴C(O)R¹³, C(O)NR¹³R¹⁴, NR¹⁴C(O)R¹³, C(O)OM, COR¹³, OR¹⁸, S(O)NR¹⁸, NR¹³-R¹⁸,

30

378

NR¹⁸OR¹⁴, N⁺R⁹R¹¹R¹²A⁻, P⁺R⁹R¹¹R¹²A⁻, amino acid, peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary heteroaryl can be further substituted with OR⁹, NR⁹R¹⁰, N⁺R⁹R¹¹R¹²A⁻, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰, SO₂OM, SO₂NR⁹R¹⁰, PO(OR⁹)OR⁹, P⁺R⁹R¹¹R¹²A⁻, S⁺R⁹R¹¹A⁻, or C(O)OM, and

wherein R¹⁸ is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heterocycle, alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heterocycle, alkyl quaternary heterocycle, and quaternary heteroaryl optionally are substituted with one or more substituent selected from the group consisting of OR⁹, NR⁹R¹⁰, N⁺R⁹R¹¹R¹²A⁻, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰, SO₃R⁹, SO₂OM, SO₂NR⁹R¹⁰, PO(OR⁹)OR⁹, and C(O)OM,

wherein in R¹ and R², one or more carbons are optionally replaced by O, NR¹³, N⁺R¹³R¹⁴A⁻, S, SO, SO₂, S⁺R¹³A⁻, PR¹³, P(O)R¹³, P⁺R¹³R¹⁴A⁻, phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl,

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR⁹, N⁺R⁹R¹⁰A⁻, S, SO, SO₂, S⁺R⁹A⁻, PR⁹, P⁺R⁹R¹⁰A⁻, or P(O)R⁹;

wherein quaternary heterocycle and quaternary heteroaryl are optionally substituted with one or more groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen,

oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, SO₃R¹³, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴, P⁺R¹³R¹⁴R¹⁵A⁻, P(OR¹³)OR¹³, S⁺R¹³R¹⁴A⁻, and N⁺R⁹R¹¹R¹²A⁻,

R¹ is selected from the group consisting of alkane diyl, alkene diyl, alkene diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, and peptide, polypeptide, wherein alkane diyl, alkene diyl, alkene diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, and peptide polypeptide, can optionally have one or more carbon replaced by O, NR⁷, N⁺R⁷R⁸, S, SO, SO₂, S⁺R⁷R⁸, PR⁷, P⁺R⁷R⁸, phenylene, heterocycle, quaternary heterocycle, quaternary heteroaryl, or aryl,

wherein alkane diyl, alkene diyl, alkene diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, peptide, and polypeptide can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, SO₃R¹³, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴, P⁺R¹³R¹⁴R¹⁵A⁻, P(OR¹³)OR¹³, S⁺R¹³R¹⁴A⁻, and N⁺R⁹R¹¹R¹²A⁻;

wherein one or more R¹ and R² are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, OR⁹, SR⁹, S(O)R⁹, SO₂R⁹, and SO₃R⁹,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl, and heterocycle can be substituted with one or more substituent groups independently selected from the

group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo, OR¹³, NR^{13,14}, SR¹³, S(O)R¹³, SO₂R¹³, SO₃R¹³, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R^{13,14}, P⁺R¹³R¹⁴, P(O)R¹³R¹⁴SR¹⁵A⁻, P(O)R¹³OR¹⁴, SR¹³R¹⁴A⁻, and NR⁹R¹¹R¹²A⁻,

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of OR⁷,

NR^{7,8}, SR⁷, S(O)R⁷, SO₂R⁷, SO₃R⁷, CO₂R⁷, CN, oxo, CONR^{7,8}, N⁺R^{7,8}R⁹A⁻, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary

heterocycle, quaternary heteroaryl, P(O)R^{7,8}, P⁺R^{7,8}A⁻, and P(O)OR⁷, and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O,

NR⁷, N⁺R^{7,8}A⁻, S, SO, SO₂, S⁺R⁷A⁻, PR⁷, P(O)R⁷, P⁺R^{7,8}A⁻, or phenylene.

99. A compound of claim 98, wherein R¹, R¹⁴, R¹, and R¹⁴ are independently selected from the group consisting of H and alkyl.

100. A compound of claim 99, wherein R¹, R¹⁴, R¹, and R¹⁴ are independently selected from the group consisting of H and C₁-C₁₀ alkyl.

101. A compound of claim 100, wherein said alkyl is a C₁-C₁₀ alkyl.

102. A compound of claim 101, wherein R¹, R¹⁴, R¹, and R¹⁴ are independently C₁-C₁₀ alkyl.

381

103. A compound of claim 102, wherein R¹, R¹⁴, R¹, and R¹⁴ are independently selected from the group consisting of ethyl, n-propyl, n-butyl, and isobutyl.

104. A compound of claim 98, wherein R¹, R¹⁴, R¹, and R¹⁴ are independently selected from the group consisting of H and OR⁷.

105. A compound of claim 104, wherein R⁷ is H.

106. A compound of claim 98, wherein R⁷, R¹⁴, R¹, and R¹⁴ are H.

107. A compound of claim 98, wherein d and e are independently 1 or 2.

108. A compound of claim 107, wherein d and e are both 2.

109. A compound of claim 98, wherein one or more R^x and one or more R¹⁴ are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen, OR¹³, NR^{13,14}, NR¹³NR¹⁴R¹⁵, N⁺R⁹R^{11,12}A⁻, SR¹³, S⁺R^{13,14}, CO₂R¹³, NR¹³C(O)R¹⁴, and NR¹³C(O)R¹⁴,

wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with OR⁹, NR^{9,10}, N⁺R⁹R^{11,12}A⁻, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰, SO₂NR⁹, PO(OR¹¹)OR¹¹, P⁺R⁹R^{11,12}A⁻, S⁺R⁹R¹¹A⁻, or C(O)OM, and

wherein in R^x, one or more carbons are optionally replaced by O, NR¹³, N⁺R^{13,14}A⁻, S, SO, SO₂, S⁺R¹³A⁻,

PR¹³, P(O)R¹³, P⁺R^{13,14}A⁻, phenylene, amino acid,

382

peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR⁹, N⁺R⁹R¹⁰A⁻, S, SO, SO₂, S⁺R⁹A⁻, PR⁹, P⁺R⁹R¹⁰A⁻, or P(O)R⁹.

110. A compound of claim 98, wherein one or more R^y and one or more Rⁿ are independently selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide, NR¹³R¹⁴, NR¹⁴C(O)Rⁿ, and ORⁿ, wherein alkyl and polyether can be further substituted with SO₂R⁹, N⁺R⁹R¹¹R¹²A⁻, and quaternary heteroaryl.

111. A compound of claim 98, wherein Rⁿ is selected from the group consisting of alkane diyl, polyalkane diyl, alkoxy diyl, and polyalkoxy diyl, wherein alkane diyl and polyalkane diyl can optionally have one or more carbon replaced by O, NR⁷, N⁺RR⁸, S, SO, SO₂, S⁺RR⁸, PR⁷, P⁺RR⁸, or phenylene.

112. A compound of claim 111, wherein Rⁿ is selected from the group consisting of alkoxy diyl and polyalkoxydiyl wherein one or more carbons are optionally replaced by O, NR⁹, N⁺R⁹R¹⁰, S, SO, SO₂, S⁺R⁹R¹⁰, PR⁹, P⁺R⁹R¹⁰, phenylene, amino acid, peptide, polypeptide, carbohydrate, or polyalkyl.

113. A compound of claim 112, wherein R¹, Rⁿ, R¹, and Rⁿ are independently selected from the group consisting of H and alkyl.

114. A compound of claim 113, wherein R¹, Rⁿ, R¹, and Rⁿ are independently selected from the group consisting of H and OR¹.

115. A compound of claim 114, wherein R¹ is H.

116. A compound of claim 115, wherein R¹, Rⁿ, R¹, and Rⁿ are each H.

117. A compound of claim 116, wherein d and e are independently 1 or 2.

118. A compound of claim 117, wherein one or more R^x and one or more Rⁿ are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen, OR¹³, NR¹³R¹⁴, NR¹³NR¹⁴R¹⁵, N⁺R⁹R¹¹R¹²A⁻, SR¹³, S⁺R¹³R¹⁴, CO₂R¹³, NR¹³C(O)Rⁿ, and NR¹⁴C(O)Rⁿ,

wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with OR⁹, NR⁹R¹⁰, N⁺R⁹R¹¹R¹²A⁻, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰, SO₂OM, SO₂NR⁹Rⁿ, PO(OR¹)ORⁿ, P⁺R⁹R¹¹R¹²A⁻, S⁺R⁹Rⁿ, or C(O)OM, and

wherein in R^x, one or more carbons are optionally replaced by O, NR¹³, N⁺R¹³R¹⁴A⁻, S, SO, SO₂, S⁺R¹³A⁻, PR¹³, P(O)R¹³, P⁺R¹³R¹⁴A⁻, phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and


wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR⁹, N⁺R⁹R¹⁰A⁻, S, SO, SO₂, S⁺R⁹A⁻, PR⁹, P⁺R⁹R¹⁰A⁻, or P(O)R⁹.

119. A compound of claim 118, wherein one or more R^y and one or more Rⁿ are independently selected from

the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide, $\text{NR}^{\text{13}}\text{R}^{\text{14}}$, $\text{NR}^{\text{14}}\text{C}(\text{O})\text{R}^{\text{15}}$, and OR^{15} , wherein alkyl and polyether can be further substituted with SO_3R^9 , $\text{N}^+\text{R}^9\text{R}^{\text{11}}\text{R}^{\text{12}}\text{A}^-$, and quaternary heteroaryl.

5

120. A compound of claim 119, having the formula:




PEG = 3400 molecular weight polyethylene glycol polymer chain

10

121. A compound of the formula (IV)

385



wherein :

q and r are independently integers from 0 to 3;
d and e are independently integers from 0 to 2;
t and u are independently integers from 0 to 5;
 R^1 , R^{11} , R^{12} , and R^{13} are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkoalkyl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl,

wherein alkyl, alkenyl, alkynyl, haloalkyl, alkoalkyl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl optionally are substituted with one or more substituent selected from the group consisting of OR^9 , $\text{NR}^9\text{R}^{\text{10}}$, $\text{N}^+\text{R}^9\text{R}^{\text{11}}\text{R}^{\text{12}}\text{A}^-$, SR^9 , $\text{S}^+\text{R}^9\text{R}^{\text{11}}\text{R}^{\text{12}}\text{A}^-$, $\text{S}(\text{O})\text{R}^9$, SO_2R^9 , SO_3R^9 , CO_2R^9 , CN, halogen, oxo, and $\text{CONR}^9\text{R}^{\text{10}}$,

wherein alkyl, alkenyl, alkynyl, alkoalkyl, alkoxy, alkoxyalkyl, polyalkyl, aryl, and cycloalkyl optionally have one or more carbons replaced by O, NR^9 , $\text{N}^+\text{R}^9\text{R}^{\text{10}}$, S, SO, SO_2 , $\text{S}^+\text{R}^9\text{R}^{\text{10}}$, $\text{P}^+\text{R}^9\text{R}^{\text{10}}\text{A}^-$, or phenylene,

wherein R^9 , R^{10} , and R^{11} are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle, ammoniumalkyl, alkylammoniumalkyl, and arylalkyl; or

25

386

R¹ and R² taken together with the carbon to which they are attached form C₁-C₂, cycloalkylidene, or R³ and R⁴ taken together with the carbon to which they are attached form C₃-C₄, cycloalkylidene;

R¹, R³, R⁵, and R⁶ are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, acyloxy, aryl, heterocycle, OR⁹, NR⁹R¹⁰, SR⁹, S(O)R⁹, SO₂R⁹, and SO₂R⁹, wherein R⁹ and R¹⁰ are as defined above; or

R³ and R⁴ together form =O, =NOR¹¹, =S, =NNR¹¹R¹², =NR⁹, or =CR¹¹R¹², or

R^{3A} and R^{4A} together form =O, =NOR¹¹, =S, =NNR¹¹R¹², =NR⁹, or =CR¹¹R¹²,

wherein R¹¹ and R¹² are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl, heterocycle, carboxyalkyl, carboalkoxyalkyl, cycloalkyl, cyanoalkyl, OR⁹, NR⁹R¹⁰, SR⁹, S(O)R⁹, SO₂R⁹, CO₂R⁹, CN, halogen, oxo, and CONR⁹R¹⁰,

wherein R⁹ and R¹⁰ are as defined above, provided that both R³ and R⁴ cannot be OH, NH₂, and SH, or

R¹¹ and R¹² together with the nitrogen or carbon atom to which they are attached form a cyclic ring;

wherein A⁻ is a pharmaceutically acceptable anion and M is a pharmaceutically acceptable cation;

R⁷, R⁸, R⁹, and R¹⁰ are independently selected from the group consisting of hydrogen and alkyl; and one or more R^X and R^Y are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen, haloalkyl, cycloalkyl, heterocycle, heterocycle, polyether, quaternary heterocycle, quaternary heteroaryl, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, S(O)₂R¹³,

387

SO₂R¹³, S⁺R¹³R¹⁴A⁻, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, NR¹³C(O)R¹⁴, C(O)NR¹³R¹⁴, NR¹⁴C(O)R¹³, C(O)OM, COR¹³, OR¹⁸, S(O)NR¹⁸, NR¹³R¹⁸, NR¹⁸OR¹⁴, N⁺R⁹R¹¹R¹²A⁻, P⁺R⁹R¹¹R¹²A⁻, amino acid, peptide, polypeptide, and carbohydrate.

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl, polyether, quaternary heterocycle, and quaternary heteroaryl can be further substituted with OR⁹, NR⁹R¹⁰, N⁺R⁹R¹¹R¹²A⁻, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰, SO₂OM, SO₂NR⁹R¹⁰, PO(OR⁹)OR⁹, P⁺R⁹R¹¹R¹²A⁻, S⁺R⁹R¹¹A⁻, or C(O)OM, and

wherein R¹⁸ is selected from the group consisting of acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heterocycle, alkyl,

wherein acyl, arylalkoxycarbonyl, arylalkyl, heterocycle, heterocycle, alkyl quaternary heterocycle, and quaternary heteroaryl optionally are substituted with one or more substituent selected from the group consisting of OR⁹, NR⁹R¹⁰, N⁺R⁹R¹¹R¹²A⁻, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰, SO₃R⁹, SO₂OM, SO₂NR⁹R¹⁰, PO(OR⁹)OR⁹, and C(O)OM,

wherein in R⁸ and R⁹, one or more carbons are optionally replaced by O, NR¹³, N⁺R¹³R¹⁴A⁻, S, SO, SO₂, S⁺R¹³A⁻, PR¹³, P(O)R¹³, P⁺R¹³R¹⁴A⁻, phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl,

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR⁹, N⁺R⁹R¹⁰A⁻, S, SO, SO₂, S⁺R⁹A⁻, PR⁹, P⁺R⁹R¹⁰A⁻, or P(O)R⁹;

wherein quaternary heterocycle and quaternary heteroaryl are optionally substituted with one or more

388

groups selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴, P⁺R¹³R¹⁴R¹⁵A⁻, P(OR¹³)OR¹⁴, S⁺R¹³R¹⁴A⁻, and N⁺R⁹R¹¹R¹²A⁻.

R¹⁰ is selected from the group consisting of alkane diyl, alkene diyl, alkene diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, and peptide, polypeptide, wherein alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, and peptide polypeptide, can optionally have one or more carbon replaced by O, NR⁷, N⁺R⁷R⁸, S, SO, SO₂, S⁺R⁷R⁸, PR⁷, P⁺R⁷R⁸, phenylene, heterocycle, quaternary heterocycle, quaternary heteroaryl, or aryl, wherein alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, and peptide polypeptide can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, SO₃R¹³, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴, P⁺R¹³R¹⁴R¹⁵A⁻, P(OR¹³)OR¹⁴, S⁺R¹³R¹⁴A⁻, and N⁺R⁹R¹¹R¹²A⁻;

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of OR⁷, NR⁷R⁸, SR⁷, S(O)R⁷, SO₂R⁷, SO₃R⁷, CO₂R⁷, CN, oxo,

CONR⁷R⁸, N⁺R⁷R⁸A⁻, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, P(O)R⁷R⁸, P⁺R⁷R⁸A⁻, and P(O)(OR⁷)OR⁸, and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O, NR⁷, N⁺R⁷R⁸A⁻, S, SO, SO₂, S⁺R⁷R⁸A⁻, PR⁷, P(O)R⁷, P⁺R⁷R⁸A⁻, or phenylene.

122. A compound of claim 121, wherein R¹, R^{1a}, R¹, and R^{1a} are independently selected from the group consisting of H and alkyl.

123. A compound of claim 122, wherein R¹, R^{1a}, R¹, and R^{1a} are independently selected from the group consisting of H and C₁-C₁₀ alkyl.

124. A compound of claim 123, wherein said alkyl is a C₁-C₁₀ alkyl.

125. A compound of claim 124, wherein R¹, R^{1a}, R¹, and R^{1a} are independently C₁-C₁₀ alkyl.

126. A compound of claim 125, wherein R¹, R^{1a}, R¹, and R^{1a} are independently selected from the group consisting of ethyl, n-propyl, n-butyl, and isobutyl.

127. A compound of claim 125, wherein R¹, R^{1a}, R¹, and R^{1a} are independently selected from the group consisting of H and OR¹.

128. A compound of claim 127, wherein R¹ is H.

129. A compound of claim 121, wherein R¹, R^{1a}, R¹, and R^{1a} are H.

130. A compound of claim 121, wherein d and e are independently 1 or 2.

131. A compound of claim 130, wherein d and e are both 2.

132. A compound of claim 121, wherein one or more R^x and one or more R^m are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen, OR^{13} , $NR^{13}R^{14}$, $NR^{13}NR^{14}R^{15}$, $N^+R^9R^{11}R^{12}A^-$, SR^{13} , $S^+R^{13}R^{14}$, CO_2R^{13} , $NR^xC(O)R^x$, and $NR^xC(O)R^x$,

wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with OR^9 , NR^9R^{10} , $N^+R^9R^{11}R^{12}A^-$, SR^9 , $S(O)R^9$, SO_2R^9 , SO_3R^9 , oxo, CO_2R^9 , CN, halogen, $CONR^9R^{10}$, $SO_2NR^9R^{10}$, $PO(OR^9)OR^9$, $P^+R^9R^{11}R^{12}A^-$, $SR^9R^9A^-$, or $C(O)OM$, and

wherein in R^x , one or more carbons are optionally replaced by O, NR^{13} , $N^+R^{13}R^{14}A^-$, S, SO, SO_2 , $S^+R^{13}A^-$, PR^{13} , $P(O)R^9$, $P^+R^{13}R^{14}A^-$, phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR^9 , $N^+R^9R^{10}A^-$, S, SO, SO_2 , $S^+R^9A^-$, PR^9 , $P^+R^9R^{10}A^-$, or $P(O)R^9$.

133. A compound of claim 121, wherein one or more R^y and one or more R^m are independently selected from the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide, $NR^{13}R^{14}$, $NR^xC(O)R^x$, and OR^x ,

wherein alkyl and polyether can be further substituted with SO_3R^9 , $N^+R^9R^{11}R^{12}A^-$, and quaternary heteroaryl.

134. A compound of claim 121, wherein R^m is selected from the group consisting of alkane diyl, polyalkane diyl, alkoxy diyl, and polyalkoxy diyl, wherein alkane diyl and polyalkane diyl can optionally have one or more carbon replaced by O, NR^7 , NR^7R^8 , S, SO, SO_2 , $S^+R^7R^8$, PR^7 , $P^+R^7R^8$, or phenylene.

135. A compound of claim 134, wherein R^m is selected from the group consisting of alkoxy diyl and polyalkoxydiyl wherein one or more carbons are optionally replaced by O, NR^9 , $N^+R^9R^{10}$, S, SO, SO_2 , $S^+R^9R^{10}$, PR^9 , $P^+R^9R^{10}$, phenylene, amino acid, peptide, polypeptide, carbohydrate, or polyalkyl.

136. A compound of claim 135, wherein R^1 , R^2 , R^3 , and R^m are independently selected from the group consisting of H and alkyl.


137. A compound of claim 136, wherein R^1 , R^2 , R^3 , and R^m are independently selected from the group consisting of H and OR^1 .

138. A compound of claim 137, wherein R^1 is H.

139. A compound of claim 138, wherein R^1 , R^2 , R^3 , and R^m are each H.


140. A compound of claim 139, wherein d and e are independently 1 or 2.

141. A compound of claim 140, having the formula:



PEG = 3400 molecular weight polyethylene glycol polymer chain

142. A compound of formula (V)



wherein :

q is an integer from 0 to 4;

r is an integer from 0 to 3;

d and e are independently integers from 0 to 2;

```
t is an integer from 0 to 4;
```

u is an integer from 0 to 5;

R^1 , R^2 , R^3 , and R^4 are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl, dialkylamino, alkylthio, (polyalkyl)aryl, and cycloalkyl.

wherein alkyl, alkenyl, alkenyl, alkynyl, haloalkyl, alkylaryl, arylalkyl, alkoxy, alkoxyalkyl,

dialkylamino, alkylthio, (polyalkyl)aryl, and

cycloalkyl optionally are substituted with one or more substituent selected from the group consisting of OR⁹, NR⁹R¹⁰, N^aR^bR^cR^d, SR⁹, S^aR^b-P^a^bR^cR^dA^e, S(O)R⁹, SO₂R⁹,

SO_3R^9 , CO_2R^9 , CN, halogen, oxo, and $\text{CONR}^9\text{R}^{10}$,
wherein alkyl, alkenyl, alkynyl, alkylaryl, alkylaryl,

alkoxy, alkoxyalkyl, polyalkyl, aryl, and cycloalkyl optionally have one or more carbons replaced by O, NR⁹, + 9 10 + 9 + 9 10

wherein R^9 , R^{10} , and R^W are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, acyl, heterocycle,

ammoniumalkyl, alkylammoniumalkyl, and arylalkyl; or R' and R' taken together with the carbon to which they are attached form C₃-C₆ cycloalkylidene, or

R^{1a} and R^{2a} taken together with the carbon to which they are attached form C-C₁₀ cycloalkylidene;

R' , R'' , R' , and R'' are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl,

acyloxy, aryl, heterocycle, OR^9 , NR^9 , R^9 , SR^9 , S(O)R^9 ,

SO_2R^y , and SO_3R^y , wherein R^y and R^n are as defined above; or

$$R^3 \text{ and } R^4 \text{ together form } O, =NOR^{11}, =S, =NNR^{11}R^{12}, \\ =NR^9, \text{ or } =CR^{11}R^{12} \text{ or}$$

3913

394

R^3A and R^4A together form =O, =NOR¹¹, =S,
=NNR¹¹R¹², =NR⁹, or =CR¹¹R¹²,

wherein R¹¹ and R¹² are independently selected
from the group consisting of H, alkyl, alkenyl,
alkynyl, aryl, arylalkyl, alkenylalkyl, alkynylalkyl,
heterocycle, carboxyalkyl, carbalkoxyalkyl,

cycloalkyl, cyanoalkyl, OR⁹, NR⁹R¹⁰, SR⁹, S(O)R⁹,
SO₂R⁹, SO₂R⁹, CO₂R⁹, CN, halogen, oxo, and CONR⁹R¹⁰,

wherein R⁹ and R¹⁰ are as defined above, provided that
both R³ and R⁴ cannot be OH, NH₂, and SH, or

R¹¹ and R¹² together with the nitrogen or carbon
atom to which they are attached form a cyclic ring;

wherein A⁻ is a pharmaceutically acceptable anion
and M is a pharmaceutically acceptable cation;

R⁷, R⁸, R⁹, and R¹⁰ are independently selected from
the group consisting of hydrogen and alkyl; and

one or more R^x and R^y are independently selected
from the group consisting of H, alkyl, alkenyl,
alkynyl, polyalkyl, acyloxy, aryl, arylalkyl, halogen,
haloalkyl, cycloalkyl, heterocycle, heterocycle,
polyether, quaternary heterocycle, quaternary

heteroaryl, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, S(O)₂R¹³,
SO₂R¹³, S⁺R¹³R¹⁴-, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³,
CN, OM, SO₂OM, SO₂NR¹³R¹⁴, NR¹³C(O)R¹⁴, C(O)NR¹³R¹⁴,
NR¹⁴C(O)R¹³, C(O)OM, COR¹³, OR¹⁸, S(O)NR¹⁸, NR¹³R¹⁸,
NR¹⁸OR¹⁴, N⁺R⁹R¹¹R¹²A⁻, P⁺R⁹R¹¹R¹²A⁻, amino acid,
peptide, polypeptide, and carbohydrate,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl,
polyalkyl, heterocycle, acyloxy, arylalkyl, haloalkyl,
polyether, quaternary heterocycle, and quaternary
heteroaryl can be further substituted with OR⁹, NR⁹R¹⁰,
N⁺R⁹R¹¹R¹²A⁻, SR⁹, S(O)R⁹, SO₂R⁹, SO₂R⁹, oxo, CO₂R⁹,

395

CN, halogen, CONR⁹R¹⁰, SO₂OM, SO₂NR⁹R¹⁰, PO(OR¹¹)OR¹²,
P⁺R⁹R¹¹R¹²A⁻, SR⁹R¹⁴, or C(O)OM, and

wherein R¹⁸ is selected from the group consisting
of acyl, arylalkoxy, carbonyl, arylalkyl, heterocycle,
heterocycle, alkyl,

wherein acyl, arylalkoxy, carbonyl, arylalkyl,
heterocycle, heterocycle, alkyl quaternary heterocycle,
and quaternary heteroaryl optionally are substituted
with one or more substituent selected from the group
consisting of OR⁹, NR⁹R¹⁰, N⁺R⁹R¹¹R¹²A⁻, SR⁹, S(O)R⁹,
SO₂R⁹, SO₂R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰, SO₂R⁹,
SO₂OM, SO₂NR⁹R¹⁰, PO(OR¹⁶)OR¹⁷, and C(O)OM,

wherein in Rⁱ and R^j, one or more carbons are
optionally replaced by O, NR¹³, N⁺R¹³R¹⁴A⁻, S, SO, SO₂,
S⁺R¹³A⁻, RR¹³, P(O)R¹³, P⁺R¹³R¹⁴A⁻, phenylene, amino
acid, peptide, polypeptide, carbohydrate, polyether, or
polyalkyl,

wherein in said polyalkyl, phenylene, amino acid,
peptide, polypeptide, and carbohydrate, one or more
carbons are optionally replaced by O, NR⁹, N⁺R⁹R¹⁰A⁻,
S, SO, SO₂, S⁺R⁹A⁻, PR⁹, P⁺R⁹R¹⁰A⁻, or P(O)R⁹;

wherein quaternary heterocycle and quaternary
heteroaryl are optionally substituted with one or more
groups selected from the group consisting of alkyl,
alkenyl, alkynyl, polyalkyl, polyether, aryl,
haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen,
oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³,
NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM,
SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴,
P⁺R¹³R¹⁴R¹⁵A⁻, P(OR¹³)OR¹⁴, S⁺R¹³R¹⁴A⁻, and N⁺R⁹R¹¹R¹²A⁻,
R¹⁸ is selected from the group consisting of alkene
diyl, alkene diyl, alkene diyl, polyalkene diyl, alkoxy
diyl, polyether diyl, polyalkoxy diyl, carbohydrate,

396

amino acid, and peptide, polypeptide, wherein alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl, polyalkoxy diyl, carbohydrate, amino acid, and peptide polypeptide, can optionally have one or more carbon replaced by O, NR⁷, N⁺R⁷R⁸, S, SO, SO₂, S⁺R⁷R⁸, PR⁷, P⁺R⁷R⁸, phenylene, heterocycle, quaternary heterocycle, quaternary heteroaryl, or aryl, wherein alkane diyl, alkene diyl, alkyne diyl, polyalkane diyl, alkoxy diyl, polyether diyl,

polyalkoxy diyl, carbohydrate, amino acid, peptide, and polypeptide can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, SO₃R¹³, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴, P⁺R¹³R¹⁴, R¹⁵A⁻, P(OR¹³)OR¹⁴, S⁺R¹³R¹⁴, and N⁺R¹³R¹⁴R¹²A⁻;

wherein one or more R⁷ and R⁸ are independently selected from the group consisting of H, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, quaternary heterocycle, OR⁹, SR⁹, S(O)R⁹, SO₂R⁹, and SO₃R⁹,

wherein alkyl, alkenyl, alkynyl, aryl, cycloalkyl, and heterocycle can be substituted with one or more substituent groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, heterocycle, arylalkyl, halogen, oxo, OR¹³, NR¹³R¹⁴, SR¹³, S(O)R¹³, SO₂R¹³, SO₃R¹³, NR¹³OR¹⁴, NR¹³NR¹⁴R¹⁵, NO₂, CO₂R¹³, CN, OM, SO₂OM, SO₂NR¹³R¹⁴, C(O)NR¹³R¹⁴, C(O)OM, COR¹³, P(O)R¹³R¹⁴, P⁺R¹³R¹⁴, R¹⁵A⁻, P(OR¹³)OR¹⁴, S⁺R¹³R¹⁴, and N⁺R¹³R¹⁴R¹²A⁻,

397

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can be further substituted with one or more substituent groups selected from the group consisting of OR⁷, NR⁷R⁸, SR⁷, S(O)R⁷, SO₂R⁷, SO₃R⁷, CO₂R⁷, CN, oxo, COMR⁷R⁸, N⁺R⁷R⁸A⁻, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heterocycle, arylalkyl, quaternary heterocycle, quaternary heteroaryl, P(O)R⁷R⁸, P⁺R⁷R⁸A⁻, and P(O)(OR⁷)OR⁸, and

wherein said alkyl, alkenyl, alkynyl, polyalkyl, polyether, aryl, haloalkyl, cycloalkyl, and heterocycle can optionally have one or more carbons replaced by O, NR⁷, N⁺R⁷R⁸A⁻, S, SO, SO₂, S⁺R⁷A⁻, PR⁷, P(O)R⁷, P⁺R⁷R⁸A⁻, or phenylene.

143. A compound of claim 142, wherein R¹, R², R³, and R⁴ are independently selected from the group consisting of H and alkyl.

144. A compound of claim 143, wherein R¹, R², R³, and R⁴ are independently selected from the group consisting of H and C₁-C₁₀ alkyl.

145. A compound of claim 144, wherein said alkyl is a C₁-C₁₀ alkyl.

146. A compound of claim 145, wherein R¹, R², R³, and R⁴ are independently C₁-C₁₀ alkyl.

147. A compound of claim 146, wherein R¹, R², R³, and R⁴ are independently selected from the group consisting of ethyl, n-propyl, n-butyl, and isobutyl.

148. A compound of claim 142, wherein R¹, R², R³, and R⁴ are independently selected from the group consisting of H and OR⁷.

398

149. A compound of claim 148, wherein R' is H.

150. A compound of claim 142, wherein R', R'', R', and R'' are H.

151. A compound of claim 142, wherein d and e are independently 1 or 2.

152. A compound of claim 151, wherein d and e are both 2.

153. A compound of claim 142, wherein one or more R^x and one or more R'' are independently selected from the group consisting of alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, polyether, halogen, OR¹³, NR¹³R¹⁴, NR¹³NR¹⁴R¹⁵, N⁺R⁹R¹¹R¹²A⁻, SR¹³, S⁺R¹³R¹⁴, CO₂R¹³, NR¹³C(O)R'', and NR¹³C(O)R'',

wherein alkyl, aryl, cycloalkyl, heterocycle, polyalkyl, acyloxy, and polyether, can be further substituted with OR⁹, NR⁹R¹⁰, N⁺R⁹R¹¹R¹²A⁻, SR⁹, S(O)R⁹, SO₂R⁹, SO₃R⁹, oxo, CO₂R⁹, CN, halogen, CONR⁹R¹⁰, SO₂OM, SO₂NR⁹R'', PO(OR⁹)OR'', P⁺R⁹R¹¹R¹²A⁻, S⁺R⁹R¹⁴A⁻, or C(O)OM, and

wherein in R^x, one or more carbons are optionally replaced by O, NR¹³, N⁺R¹³R¹⁴A⁻, S, SO, SO₂, S⁺R¹³A⁻, PR¹³, P(O)R'', P⁺R¹³R¹⁴A⁻, phenylene, amino acid, peptide, polypeptide, carbohydrate, polyether, or polyalkyl, and

wherein in said polyalkyl, phenylene, amino acid, peptide, polypeptide, and carbohydrate, one or more carbons are optionally replaced by O, NR⁹, N⁺R⁹R¹⁰A⁻, S, SO, SO₂, S⁺R⁹A⁻, PR⁹, P⁺R⁹R¹⁰A⁻, or P(O)R'.

154. A compound of claim 142, wherein one or more R^y and one or more R'' are independently selected from

39

the group consisting of alkyl, polyether, fluoride, chloride, bromide, iodide, NR¹³R¹⁴, NR¹³C(O)R'', and OR'', wherein alkyl and polyether can be further substituted with SO₃R⁹, N⁺R⁹R¹¹R¹²A⁻, and quaternary heteroaryl.

155. A compound of claim 142, wherein R'' is selected from the group consisting of alkane diyl, polyalkane diyl, alkoxy diyl, and polyalkoxy diyl, wherein alkane diyl and polyalkane diyl can optionally have one or more carbon replaced by O, NR⁷, N⁺R⁷R⁸, S, SO, SO₂, S⁺R⁷R⁸, PR⁷, P⁺R⁷R⁸, or phenylene.

156. A compound of claim 155, wherein R'' is selected from the group consisting of alkoxy diyl and polyalkoxydiyl wherein one or more carbons are optionally replaced by O, NR⁹, N⁺R⁹R¹⁰, S, SO, SO₂, S⁺R⁹R¹⁰, PR⁹, P⁺R⁹R¹⁰, phenylene, amino acid, peptide, polypeptide, carbohydrate, or polyalkyl.

157. A compound of claim 156, wherein R', R'', R', and R'' are independently selected from the group consisting of H and alkyl.

158. A compound of claim 157, wherein R', R'', R', and R'' are independently selected from the group consisting of H and OR'.

159. A compound of claim 158, wherein R' is H.

160. A compound of claim 159, wherein R', R'', R', and R'' are each H.

161. A compound of claim 160, wherein d and e are independently 1 or 2.


162. A compound of claim 161, having the formula:

400

a patient in need thereof a composition of claim 164 in unit dosage form.

167. A method for the prophylaxis or treatment of an atherosclerotic condition comprising administering to a patient in need thereof a composition of claim 165 in unit dosage form.

168. A method for the prophylaxis or treatment of hypercholesterolemia comprising administering to a patient in need thereof a composition of claim 166 in unit dosage form.



PEG = 3400 molecular weight polyethylene glycol polymer chain

163. A pharmaceutical composition comprising an anti-hyperlipidemic condition effective amount of a compound of formula (I) of claim 1, and a pharmaceutically acceptable carrier.

164. A pharmaceutical composition comprising an anti-atherosclerotic effective amount of a compound of formula (I) of claim 1, and a pharmaceutically acceptable carrier.

165. A pharmaceutical composition comprising an anti-hypercholesterolemia effective amount of a compound of formula (I) of claim 1, and a pharmaceutically acceptable carrier.

166. A method for the prophylaxis or treatment of a hyperlipidemic condition comprising administering to

Intern. . and Application No
PCT/US 97/04076

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D337/08 C07D409/10 C08G65/329 A61K31/38

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data have consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Quotation of document, with indication, where appropriate, of the relevant passages	Reference to claim No.
X	GB 1 211 258 A (BOEHRINGER) 4 November 1970 see page 1; claims; example 5	1, 163-165
P, X	WD 96 08404 A (MONSANTO) 21 March 1996 see the whole document -----	1-30, 163-165

Further documents are listed in the continuation of box C.

✕ Patient family members are listed in annex

* Special categories of cited documents

- | | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>1. document relating to personal data of the art which is not considered to be of general interest</p> <p>2. earlier document first published on or after the international filing date</p> <p>3. document which may throw doubts on priority claim(s) or which is used to establish the publication date of another document or to determine the date of publication of another document or relating to an oral disclosure, use, exhibition or other means</p> <p>4. document published prior to the international filing date but does not bear the priority date claimed</p> | <p>1. later document published after the international filing date which is not considered to be of general interest</p> <p>2. document published after the international filing date but is considered the principle or theory underlying an invention</p> <p>3. document of particular relevance the claimed invention, capable to substantiate novel or cannot be considered to be of general interest</p> <p>4. document of particular relevance the claimed invention, capable to substantiate novel or cannot be considered to be of general interest</p> <p>5. document is considered to involve an inventive step when the document is combined with one or more other such documents</p> <p>6. document combinations being different to a person skilled in the art</p> <p>7. document number of the same nature (family)</p> |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

Date of the actual completion of the international search

Date of mailing of the international search report

29 July 1997

0 4. 08. 97

Name and mailing address of the USA

Authorized official

European Patent Office, P.O. Box 5818 Patentland 2
NL - 2210 HV Rijswijk
Tel. (+31-70) 340-2000, Tx. 31 651 epo nl,
Fax. (+31-70) 340-2100

Francois, J

Form PCT/ISA/210 (second sheet) (July 1992)

In: national application No.
PCT/US 97/ 04076

DUX 1 Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 1(2)(a) for the following reasons:

1. ☐ **Claims Nos.:**
because they relate to subject matter not required to be searched by this Authority, namely:
- Remark: Although claim(s) 166-168**

body, the search has been carried out and based on the alleged effects of the compound/composition.

2. ☐ **Claims Not:**
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. ☐ **Claims Not:**
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box 11 Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims No.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention *litera* mentioned in the claims; it is covered by claims Nos:

Remark on Protests

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA.210 (continuation of first sheet (1)) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Appl. No.		PCT/US 97/04976	
Parent document cited in search report	Publication date	Parent family member(s)	Publication date
GB 1211258 A	04-11-70	CH 512429 A	15-09-71
		CH 512502 A	15-09-71
		CH 513111 A	30-09-71
		DE 1593760 A	08-06-72
		FR 8052 M	29-06-70
		FR 1603343 A	05-04-71

WO 9608484 A	21-03-96	AU 3373695 A	29-03-96
		EP 0781278 A	02-07-97

THIS PAGE BLANK (USPTO)